

PUBLIC HEALTH REPORTS

VOL. 34

AUGUST 15, 1919

No. 33

THE EPIDEMIOLOGY OF INFLUENZA.¹

By W. H. FROST, Surgeon, United States Public Health Service.

The history of influenza, so far as it is known, that is, for several centuries, comprises a series of long cycles in which great pandemics alternate with periods of relative quiescence, the length of cycles as measured by the intervals between pandemics being usually a matter of decades. The special characteristics of influenza pandemics are their wide and rapid extension, their high attack rates, and their great effect upon general mortality rates. Since these cycles are undoubtedly of fundamental significance in the natural history of influenza, any proper discussion of the epidemiology of the disease should cover at least one full cycle, preferably the last, from 1889 to the present. The material for such a discussion must, however, be collected from many and diverse sources and laboriously fitted together, since there is no concrete, specific, and continuous record of the prevalence or mortality of influenza during such a period of years.

Lack of Specific Records.

During great epidemics there are abundant, if not exact, records of prevalence, and the resulting mortality can be determined with fair precision, even though a large proportion of the deaths are classified under diagnoses other than influenza. In the intervals between epidemics influenza becomes inextricably confused with other respiratory diseases, having a general clinical resemblance but no definite etiological entity, so that the record of prevalence and even of mortality is virtually lost. The first requisites for epidemiological study, namely, clear differential diagnosis and systematic records of occurrence, are therefore lacking in influenza.

In the absence of these essential records, statistics of mortality from the group comprising influenza and all forms of pneumonia afford, perhaps, the nearest approximation to a record of influenza. It is not intended to suggest that the mortality from this group of

¹ Read at joint meeting of sections on Pharmacology and Therapeutics, Pathology and Physiology, and Preventive Medicine and Public Health, of the American Medical Association, at Atlantic City, June 13, 1919. This paper was published in the Journal of the American Medical Association, vol. 73, No. 5, Aug. 2, 1919, pp. 313-318.

diseases furnishes in any sense a *measure* of the prevalence of influenza, but only that it furnishes an *index*, since it is well established that the epidemic prevalence of influenza markedly affects the mortality from this group of diseases, and since it is at least probable that even in nonepidemic periods there may be some intimate and constant relation between the prevalence of influenza and the mortality from pneumonia.

The following discussion, which is necessarily confined to a few broad outlines, is, accordingly, based on records of mortality from influenza and pneumonia for a series of years, statistics of general mortality during the recent epidemic, and limited morbidity statistics.

Influenza and Pneumonia Mortality in Massachusetts, 1887-1916.

Table I and Chart 1 show for Massachusetts, a State in which continuous records are available, the number of deaths and the death rates per 100,000 from influenza and from all forms of pneumonia, by months, from 1887 to 1916, inclusive.

TABLE I.—*Death rates per 100,000 of population from pneumonia (all forms) and from influenza in Massachusetts, 1887-1916, inclusive.*

PNEUMONIA.

Year.	Total per year.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
1887.....	138.8	19.5	16.7	1.9	22.6	16.1	7.7	5.5	4.1	5.6	9.0	14.8	15.3
1888.....	172.7	24.7	25.1	26.4	21.3	16.4	8.0	5.5	4.0	6.2	11.4	9.8	13.9
1889.....	156.6	17.9	16.3	21.1	19.8	14.9	7.5	5.3	5.2	5.4	10.6	13.2	16.3
1890.....	180.0	47.8	17.7	17.6	20.1	12.7	8.5	6.2	4.8	5.0	9.4	11.8	18.3
1891.....	185.3	20.6	16.4	20.6	25.1	23.9	10.1	5.9	4.0	4.0	7.7	13.8	36.4
1892.....	213.0	61.5	25.2	23.1	21.4	17.0	8.3	5.4	4.2	6.8	8.5	12.5	19.0
1893.....	225.8	27.3	25.4	28.6	33.1	27.3	11.1	8.0	5.0	6.1	9.8	14.3	31.9
1894.....	164.0	32.8	19.9	22.7	18.5	14.0	8.5	5.1	5.1	6.8	8.2	11.3	13.6
1895.....	184.1	19.1	34.4	30.8	20.9	14.4	7.6	5.6	5.3	4.7	10.5	13.4	17.8
1896.....	182.0	19.5	20.7	24.9	25.9	18.7	10.3	7.6	4.5	7.6	10.7	13.8	18.8
1897.....	181.6	21.1	24.9	31.5	19.7	15.2	10.2	7.0	4.5	6.0	11.7	13.1	16.9
1898.....	156.0	18.8	17.1	18.5	18.7	15.7	6.6	6.0	4.8	5.6	10.6	12.8	21.0
1899.....	181.3	37.5	25.8	20.6	18.4	14.0	8.9	5.0	4.8	5.8	8.2	18.2	19.0
1900.....	188.3	22.7	21.1	42.0	30.5	17.6	8.7	5.3	4.3	5.3	6.2	9.9	14.7
1901.....	167.7	22.6	26.6	26.2	10.9	18.9	7.7	3.4	3.4	5.7	9.1	14.2	15.1
1902.....	158.9	15.7	18.9	19.9	18.8	15.5	8.0	6.2	5.3	5.7	11.8	14.5	20.6
1903.....	172.5	25.2	25.8	25.4	18.1	16.8	8.2	7.0	4.5	4.6	7.7	13.3	18.9
1904.....	172.1	22.6	22.7	24.1	21.2	14.2	6.7	5.9	4.4	6.5	9.4	13.0	16.5
1905.....	178.3	24.7	27.7	23.6	17.1	15.5	8.5	5.4	4.8	6.2	8.9	15.8	20.0
1906.....	174.1	22.5	21.9	24.1	21.7	14.9	5.1	6.2	5.0	6.0	9.4	13.8	21.0
1907.....	190.4	26.5	24.4	23.4	18.3	14.3	9.5	5.1	5.1	7.1	8.3	12.7	26.9
1908.....	165.8	26.6	22.2	21.1	19.4	13.5	6.6	4.9	5.8	6.4	9.2	12.4	17.6
1909.....	170.3	22.1	20.0	26.1	20.1	16.0	9.7	5.1	5.0	5.4	8.9	13.6	18.3
1910.....	197.6	24.1	20.7	27.5	23.8	16.9	9.7	7.3	6.3	9.1	12.3	17.0	23.6
1911.....	174.4	22.6	27.1	23.9	20.2	16.9	7.2	6.7	6.1	6.7	9.0	11.8	16.9
1912.....	152.0	19.8	20.2	21.2	16.4	12.8	6.2	5.0	4.1	5.8	8.5	10.5	19.9
1913.....	172.2	23.5	22.8	24.9	19.3	17.1	10.7	6.2	5.4	6.7	8.3	10.4	16.8
1914.....	166.0	22.9	20.1	23.2	20.4	15.2	8.2	5.1	5.5	6.0	10.1	12.5	16.8
1915.....	176.0	17.8	19.3	23.5	27.7	13.6	6.6	7.1	5.8	6.0	8.8	9.8	22.1
1916.....	176.6	35.6	25.5	23.0	17.3	14.0	7.4	5.8	4.1	5.9	7.6	12.8	18.8

TABLE 1.—*Death rates per 100,000 of population from pneumonia (all forms) and from influenza in Massachusetts, 1887—1916, inclusive—Continued.*

INFLUENZA.

Year.	Total for year.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
1887	0.90	0.05	0.14	0.28	0.14	0.05	0.05	0.10	0.05	0.05
1888	.51	.14	.05	.050505	.1405
1889	1.23	.09	.09	.14	.14	.05	0.05	0.050955
1890	18.31	10.96	2.90	1.43	.94	.40	.36	.22	.04	.18	.22	.27	.40
1891	25.74	.39	.39	1.00	3.87	4.65	1.57	.52	.52	.52	.22	.26	10.08
1892	41.06	24.33	7.64	3.02	1.83	1.10	.72	.51	.17	.21	.38	.68	.47
1893	12.27	.58	.70	1.20	2.57	2.07	.46	.17	.12	.0446	3.90
1894	14.97	7.41	3.16	1.38	1.09	.40	.20	.16	.08	.16	.20	.16	.36
1895	18.10	.79	4.50	6.97	3.29	1.11	.36	.24	.08	.08	.04	.24	.36
1896	5.81	.93	.62	1.16	1.05	.46	.39	.15	.12	.04	.27	.19	.43
1897	14.40	.83	2.42	5.68	2.50	1.02	.27	.15	.23	.15	.27	.27	.61
1898	8.56	.59	.59	1.56	1.30	1.48	.37	.11	.22	.07	.19	.26	1.82
1899	20.72	9.34	5.81	2.18	1.05	.51	.29	.07	.04	.07	.33	.18	.84
1900	25.27	1.21	1.50	9.20	9.23	1.85	.68	.21	.07	.11	.11	.32	.78
1901	18.94	4.32	6.36	4.08	1.51	.84	.46	.14	.04	.21	.18	.25	.77
1902	5.89	.59	.87	.90	.69	.73	.24	.21	.14	.03	.24	.45	.82
1903	12.65	1.57	3.08	3.66	1.61	.82	.38	.14	.10	.03	.21	.34	.70
1904	10.26	1.45	2.06	2.80	1.38	.78	.27	.20	.07	.10	.17	.34	.64
1905	14.69	2.75	4.58	3.48	1.76	.88	.30	.07	.07	.03	.03	.20	.56
1906	6.57	1.13	1.07	1.29	.78	.36	.06	.19	.13	.06	.16	.16	1.17
1907	16.13 ¹	3.61	3.48	2.28	1.04 ¹	.70	.22	.03	.03	.03	.19	.54	3.99
1908	10.42	4.48	2.10	1.51	.86	.31	.12	.06	.03	.19	.15	.22	.37
1909	7.16	.76	1.93	2.30	.88	.36	.24	.0303	.15	.21	.27
1910	7.40	1.39	1.01	1.75	1.51	.53	.09	.0903	.21	.35	.44
1911	4.63	.87	.96	.90	.79	.41	.03	.0903	.12	.06	.49
1912	6.93	1.46	1.35	1.06	.60	.26	.20	.17	.09	.09	.26	.17	1.23
1913	5.75	2.00	.76	.85	.51	.34	.1417	.14	.27	.25	.39
1914	3.58	.75	.39	.75	.64	.36	.14	.0606	.03	.17	.25
1915	6.34	.79	.60	.98	1.88	.49	.16	.08	.05	.08	.11	.22	.87
1916	14.50	6.32	3.04	1.67	.75	.56	.32	.16	.13	.05	.16	.43	.89


Compiled from number of deaths as given in Registration Reports, State of Massachusetts, 1887-1916, except data for 1912, which are taken from Mortality Statistics, Bureau of the Census.

Chart 1 shows clearly the effect of the epidemic of 1889-1892, developing in three distinct phases; the first culminating in January 1890, the second in April and May 1891, and the third in January 1892. It is noted that the mortality was higher in 1891 than in 1890, still higher in 1892; and that in 1893, although no distinct epidemic occurred, the pneumonia mortality for the year was still higher than in 1892. This corresponds to the experience in England according to Parsons,¹ and apparently represents the general experience in other countries, with some differences in chronology and relative severity of the several phases of the epidemic.

As comparable statistics for Massachusetts in 1917 and 1918 are not now at hand, it is necessary to turn to other sources for the record of these years. The Bureau of the Census has very kindly furnished to the writer, from tables prepared for publication in the 1917 Mortality Statistics, the number of deaths by months during 1917 from pneumonia and influenza in certain registration States,

¹ Parsons, H. Franklin. A Further Report on the Influenza Epidemics of 1889-90, 1891 and 1891-92. Local Govt. Board. London, 1893.

and cities of over 100,000 population, thus continuing through that year the records of monthly mortality available from published Mortality Statistics from 1910 to 1916, inclusive; and has also furnished records of deaths* in certain of these States and cities by months during 1918. The latter figures were compiled and submitted by State and local registrars in compliance with a special request from the Director of the Census. They are merely provisional and the Census Bureau in no way vouches for their accuracy; but they are believed to be sufficiently exact for present purposes.



Owing to the limitations of space the statistics of only three cities, New York, Cleveland, and San Francisco, are presented here. These are, however, fairly representative of the whole group of cities and States for which records are at hand. As may be seen from Table II and Chart 2 the mortality from influenza and pneumonia in these cities was fairly regular from 1910 to 1915, as was the case in Massachusetts (See Chart 1) and generally throughout the registration area of the United States.

August 15, 1919.

**ANNUAL DEATH RATES PER 100,000 FROM INFLUENZA AND PNEUMONIA (ALL FORMS)
BY MONTHS, 1910-1918**




CHART 2.

TABLE II.—Annual death rates per 100,000 of population from influenza and pneumonia (all-forms) by months for the years 1910-1918, inclusive, in New York, San Francisco, and Cleveland.

NEW YORK CITY.

Years.	Total.	Months.											
		Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dcc.
1910.....	229.3	327.8	300.2	343.3	303.5	225.4	162.1	125.5	108.0	126.9	154.7	211.6	365.9
1911.....	217.7	321.3	325.5	359.0	338.7	237.2	137.2	115.8	97.2	107.6	145.0	195.0	240.8
1912.....	198.6	266.5	294.5	287.9	254.6	238.3	153.5	112.3	99.0	94.9	157.8	178.5	251.6
1913.....	199.7	265.0	353.0	342.2	242.7	225.1	158.9	113.5	101.9	100.6	118.4	172.9	213.3
1914.....	184.3	247.3	293.0	319.4	241.8	211.1	131.6	92.5	86.1	88.1	110.0	162.9	234.3
1915.....	202.7	253.7	232.2	325.4	344.5	212.1	172.0	103.2	89.2	90.8	120.4	167.1	324.4
1916.....	189.7	402.5	261.7	252.0	210.9	194.6	133.1	113.7	100.5	87.7	116.4	154.0	242.1
1917.....	191.9	417.9	316.3	237.3	221.2	202.4	122.8	72.6	78.4	89.5	127.6	185.5	237.7
1918.....	636.4	348.4	339.0	548.0	432.1	201.4	105.4	81.3	71.8	3,516.0	351.6	1,259.6	549.8

SAN FRANCISCO.

1910.....	128.6	199.9	134.0	168.9	148.2	132.3	101.6	109.8	112.6	87.3	118.2	110.5	118.2
1911.....	135.7	207.3	150.0	135.6	68.6	91.2	150.0	88.5	83.0	135.0	132.7	144.0	270.9
1912.....	126.0	236.3	191.6	182.0	131.9	86.9	47.7	51.6	67.9	92.6	84.2	137.5	198.3
1913.....	139.6	261.7	192.1	194.9	143.5	122.8	91.0	66.8	72.1	99.3	125.5	110.4	194.9
1914.....	122.6	191.7	130.8	178.5	132.9	107.6	84.1	78.8	63.0	103.1	70.9	97.7	231.0
1915.....	133.2	214.6	137.4	118.8	125.4	113.6	101.4	90.4	111.0	138.7	116.2	117.4	211.8
1916.....	133.2	243.9	165.6	162.6	144.3	101.6	84.0	88.9	71.1	76.1	88.9	144.3	221.0
1917.....	127.5	219.0	185.4	130.0	121.4	125.0	72.2	80.0	77.5	118.8	100.0	144.6	167.5
1918.....	713.0	231.3	204.3	278.1	310.2	152.6	139.8	78.7	73.8	117.0	3,534.0	2,069.8	1,331.0

TABLE II.—*Annual death rates per 100,000 of population from influenza and pneumonia (all forms) by months for the years 1910–1918, inclusive, in New York, San Francisco, and Cleveland—Continued.*

CLEVELAND.

Years.	Total.	Months.											
		Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
1910....	136.3	198.3	180.0	223.0	213.5	148.3	107.9	54.2	48.2	51.8	86.6	138.0	190.0
1911....	106.5	174.2	166.9	164.1	156.0	103.4	77.5	40.5	50.6	67.1	81.0	100.9	93.2
1912....	114.6	163.6	141.2	159.7	142.7	126.2	91.7	55.2	75.0	30.6	88.8	110.1	185.4
1913....	131.9	179.8	205.1	217.5	152.5	147.4	107.5	68.1	60.6	74.3	115.3	130.9	128.6
1914....	117.2	160.1	175.2	197.0	201.8	139.9	53.3	51.6	46.3	59.0	66.4	116.0	143.7
1915....	182.7	166.5	233.1	252.6	201.8	109.3	77.4	55.5	78.8	59.2	121.8	170.2	336.8
1916....	198.5	545.0	209.2	263.8	266.9	211.2	86.6	57.6	82.1	75.7	153.6	175.0	242.9
1917....	210.9	344.0	386.0	415.0	233.8	208.0	148.8	90.2	66.4	72.1	160.0	182.9	267.0
1918....	548.0	161.3	178.5	190.4	366.3	142.6	60.0	62.5	43.5	48.0	1,812.0	1,355.0	1,376.0

Data for years 1910–1916 compiled from Mortality Statistics, Bureau of the Census, for those years.
Data for 1917 from tables prepared for publication in 1917 Mortality Statistics; advance sheets furnished by the Bureau of the Census.

Data for 1918 from special reports rendered to Director of the Census by city registrars.
Populations used are Bureau of the Census mid-year estimates.

Significant Rise in Mortality in December, 1915.

In December, 1915, and January, 1916, there occurred in New York and Cleveland a sudden sharp rise in mortality. This, though not shown distinctly in San Francisco, was almost universal and synchronous over the whole of the registration area. While this may have no direct significance in relation to the 1918 pandemic, it is at least of interest as indicating the operation of some definite and widespread factor and suggesting in this group of diseases an epidemic tendency which is perhaps not sufficiently appreciated. In January, 1916, influenza was reported to be epidemic in 22 States, including all sections of the United States (see Public Health Reports, Jan. 7, 1916). As this epidemic was of mild type it attracted but little attention at the time and has generally been forgotten.

Another Sharp and General Rise in Mortality in the Spring of 1918.

In the spring of 1918 there was another sharp and general rise in mortality from these diseases, clearly evident in Chart 2. In the larger cities on the Atlantic seaboard this increase occurred generally during January, February, and March, when pneumonia mortality normally reaches its maximum, and the increase was not so evident in all these cities as it was in New York City. In the rest of the country, especially in the central and western States, the increase occurred in April, a month during which pneumonia mortality is generally on the decline; and was sufficient to constitute an unmistakable departure from the normal. The increased mortality rate extended quite generally into May and in some areas still longer.

This occurrence has, I believe, a definite significance in relation to a coincident prevalence of influenza, and to the subsequent development of the influenza pandemic. Although there is no definite record

August 15, 1919.

of a generalized epidemic of influenza in the United States in the spring of 1918, definite local outbreaks were observed at that time. For example, Vaughan and Palmer¹ have described a mild epidemic prevailing in the Oglethorpe camps in March. The Commissioner of Health of Chicago² states, in his special report on the autumn epidemic in Chicago, that in March, 1918, distinct epidemics resembling influenza were observed in certain groups in that city. Stanley³ in a recent publication describes three epidemics of influenza occurring in April, October, and November, 1918, respectively, in San Quentin prison, California. These three epidemics were quite similar and that of April apparently developed from a newly received inmate, presumably infected in Los Angeles, indicating the presence of the infection there at that time.

Opie, Freeman, and others of the Army Commission studying pneumonia at Camp Funston,⁴ found that recurrent outbreaks of pneumonia observed there in March, April, and May, 1918, were definitely associated with coincident epidemics of a mild type of influenza.

The rise in mortality from this group of etiologically heterogeneous diseases in the spring of 1918 is so sudden, so marked, and so general throughout the United States as to point very clearly to the operation of a single definite and specific cause, something largely independent of meteorologic and other local conditions. The observed occurrence of local epidemics of influenza at that time in widely scattered localities, the intimate association established at Camp Funston between the epidemic of influenza and pneumonia, and the subsequent development of the influenza pandemic, all indicate that the increased pneumonia mortality of March and April, 1918, was the consequence of a beginning and largely unnoticed epidemic of influenza, the beginning in this country of the great pandemic which developed in the autumn.

Mild Epidemics of Influenza in Europe in April and May, 1918.

From various localities in western Europe, mild epidemics of influenza were reported in April and May, 1918; and perhaps there may be earlier reports. During June and July extensive epidemics were reported from Great Britain, various parts of Continental Europe,

¹ Vaughan, Col. Victor C., and Palmer, Capt. Geo. T. The Communicable Diseases in the National Guard and the National Army of the United States During the Six Months from Sept. 29, 1917, to Mar. 29, 1918. Military Surgeon, Vol. XLIII, No. 4, October, 1918.

² Robertson, John Dill. A Report on an Epidemic of Influenza in the City of Chicago in the Fall of 1918. Ed. Series No. 15, Dept. of Health, City of Chicago, 1918.

³ Stanley, L. L. Influenza at San Quentin Prison, California. Public Health Reports, vol. 34, No. 19, May 9, 1919.

⁴ Opie, Eugene L., Freeman, Allen W., Blake, Francis G., Small, Jas. C., and Rivers, Thomas M. Pneumonia at Camp Funston; Report to the Surgeon General, Journal Am. Med. Ass'n, vol. 72, No. 2, Jan. 11, 1919.

China, India, the Philippine Islands, and Brazil, with marked effect on general mortality rates in such of these countries for which records are available.

Course of the 1918 Epidemic.

The course of the epidemic in England and Wales, and in the United States, respectively, is illustrated by Chart 3, which shows the annual death rates from all causes by weeks in 96 great towns of England and Wales¹ with an aggregate population (1917) of 16,577,844 and in 45 large American cities² aggregating about 22,950,000 population (1918).

ANNUAL DEATH RATES FROM ALL CAUSES

BY WEEKS, MAR. 2, 1918 - APR 5, 1919

- ♦—♦ 45 AMERICAN CITIES.
POPULATION (1910) 22,672,630
- ♦—♦ 96 GREAT TOWNS OF ENGLAND AND WALES
q/a POPULATION (1917) 16,577,844




CHART 3.

In the British cities the epidemic has so far manifested three distinct waves—the first and slightest in point of mortality occurring in June and July, the second and most severe in November, the third in February and March. Data which need not be cited here in detail indicate that the course of the epidemic in western Europe generally was similar. In cities of India the sequence was similar but the mortality far greater. In the United States the epidemic developed more largely in a single wave during September, October, and November. If, however, the epidemic already mentioned as occurring in the spring be considered the first phase and the explosive outbreak of the autumn the second, a third phase or recrudescence is quite evident in many areas, though not shown distinctly in Chart 3, in which are combined a number of widely separated cities whose curves overlap. In general, this winter recrudescence was less marked in those cities which suffered most severely in the autumn epidemic.

¹ Weekly Return of Births and Deaths Registered in London and Ninety-five other Great Towns of England and Wales, London.

² Weekly Health Index, Bureau of the Census, Washington; D. C.

August 15, 1919.

The course of the epidemic in the United States since September is already well known and need not be discussed here in detail. The prevalence of a serious epidemic in the civilian population was first recognized in and around Boston about the first of September. Within about two weeks the epidemic was general in cities on the Atlantic seaboard, developing a little later among the cities farther west. Rural districts were usually attacked somewhat later than large cities in the same sections. The sequence of the epidemic in three cities—Boston, Washington, and San Francisco—is shown in Chart 4. The interval between the peaks of the epidemic in Boston and San Francisco is four weeks, which is approximately the maximum difference between any two large cities in this phase of the epidemic.




CHART 4.

The curves of mortality differed widely in different cities, as illustrated in Chart 5, showing in Philadelphia a very explosive and abrupt curve, observed more frequently among cities on the eastern seaboard; in New York a less explosive curve, with a more prolonged high mortality rate, as observed more commonly in middle western cities, often with a more pronounced recrudescence; in St. Louis a still more gradual curve, declining only slightly after the first wave, and with a severe recrudescence, curves of this general character being more common in the Middle West.

Total Number of Deaths Among the Civil Population.

Based on incomplete statistics, which represent, however, all the main geographic divisions of the country, including both urban and rural populations, the deaths in the civil population of this country directly attributable to the epidemic are estimated at not less than




CHART 5.

450,000, or more than 4 per thousand. This rate exceeds even a liberal estimate of the net excess mortality rate in Massachusetts during the four years, 1890-1893, following the influenza outbreak in 1889-90.

Difference in Mortality in Various Geographic Groups.

Classifying the 45 cities of the Weekly Health Index into three broad geographic groups there is a noticeable difference in mortality. In the cities east of the line of the Appalachians the excess mortality from pneumonia and influenza during the weeks ended September 14, 1918, to March 1, 1919,¹ was approximately 5.6 per 1,000; in cities between the Rocky Mountains and the Appalachians 4.35; and in those of the Pacific coast 5.55 per 1,000.

Notwithstanding this general geographic relation, there are notably wide differences in the mortality rates of individual cities in the same section, even between cities close together, differences which are not as yet explained on the basis of climate, density of population, character of preventive measures exercised, or any other determined environmental factor.

More intimate details of epidemiology can be given only the briefest mention here. In order to secure reliable statistics of morbidity the Public Health Service has made special house-to-house surveys in a number of localities,² ascertaining the number of persons affected, the dates of onset, and a few other simple facts in accurately enumerated groups representative of the general population. Partial analysis of the results of these surveys in eight localities,³ giving an aggregate of 112,958 persons canvassed, shows the following as the chief facts of interest:

Results of House to House Surveys.

The percentage of the population attacked varied from 15 per cent in Louisville to 53.3 per cent in San Antonio, Texas, the aggregate for the whole group being about 28 per cent.⁴ This agrees with scattered observations in the first phase of the 1889-90 epidemic, when the attack rate seems to have varied within about these limits.

The case incidence was found to be uniformly highest in children from 5 to 14 years old, and progressively lower in each higher age group. It was slightly higher in females than in males of corresponding age; usually higher in the white than the colored population.

The ratio of pneumonia cases to total population varied from 5.3 cases per 1,000 in Spartanburg, S. C., to 24.6 per 1,000 in the smaller towns of Maryland. The pneumonia rate showed little correlation with the influenza attack rate.

The ratio of deaths to population varied from 1.9 per 1,000 in Spartanburg to 6.8 in Maryland towns. The death rate was by no

¹ United States Census Bureau, *Weekly Health Index*, Mar. 1, 1919. "Summary, Influenza Epidemic."

² For a description of methods and more detailed statistics of certain localities see *Public Health Reports*, Volume 34, No. 11, pp. 491 to 504.

³ Data from surveys of population groups as follows: New London, Conn., 7,093; Baltimore, Md. 38,361; certain smaller towns and rural districts in Maryland, 12,689; Spartanburg, S. C., 5,257; Louisville, Ky., 12,602; Little Rock, Ark., 9,920; San Antonio, Tex., 12,534; San Francisco, Calif., 18,682.

⁴ The surveys were made mostly during December, after the first wave of the epidemic had subsided, but before the second wave. Later figures would add somewhat to the incidence.

means parallel to the influenza attack rate, but was correlated closely with the pneumonia rate. In other words, the case fatality of pneumonia tended to be fairly constant, around 30 per cent, except in San Antonio, where it was only 18.5 per cent. The death rate was notably high in children under one year old, in adults from 20 to 40, and in persons over 60; higher in males than in females of comparable ages; higher in the white than in the colored.

The case fatality was likewise higher in these age groups, under one year, 20 to 40, and over 60 years; and it is this fact, rather than the incidence rates, which determines the death rates in different age groups.




CHART 6.

Chart 6, shows the attack rates, death rates and case fatality rates in various age groups.

The Important Question of Immunity.

Concerning the important question of the immunity conferred by an attack of influenza, the evidence is not conclusive, chiefly because of the uncertain differentiation between influenza and other infections. Parsons, from his study of the last epidemic,¹ inclines to the view that an attack of influenza in the earlier years of the epidemic conferred a considerable but not absolute immunity in the later outbreaks.

¹ Loc. cit.

In Baltimore, where our first canvass of 33,776 people was made between November 20 and December 11, 1918, a second canvass of the same population was made in January to determine the extent of the recrudescence reported in December. Among 32,600 people included in this canvass, 724 cases of influenza were found to have occurred since the previous survey. Of this number, 121 cases were reported as second attacks, but on investigation through the attending physicians or by a medical officer, the clinical diagnosis of both attacks as influenza was confirmed in only 26 cases, or 0.37 per cent of the total, and even in these cases the diagnosis is necessarily uncertain. Considering that 23 per cent of the population had had influenza prior to December 11, the proportion of second attacks should have been much greater if no immunity had been acquired. A second canvass in San Francisco gave generally similar results. The data collected for a study of this question from other angles have not yet been analyzed.

General Characteristics of Epidemic.

In general this epidemic has been quite similar to that of 1889-90 in its early development, first in mild scattered outbreaks, later in a severe world-wide epidemic; in the rapidity of its spread and in its high case incidence. It has been notably different in a much higher frequency of pneumonia and consequently much higher mortality, especially among young adults.

Such evidence as has been collected confirms the conclusion previously reached that the infection is transmitted directly by "contact" in the broad sense. It appears probable, however, that the infection was already widely disseminated in this country some time before a serious epidemic was recognized.

Probability of Recurrence in the Near Future.

The question of most practical and immediate interest is the probability of recurrence in the near future. Recurrences are characteristic of influenza epidemics; and the history of the last pandemic and previous ones would seem to point to the conclusion that this one has not yet run its full course. On the other hand this epidemic has already shown three more or less distinct phases and has been more severe, at least in mortality, than the three-year epidemic of 1889-92, facts which may justify the hope, though not the conclusion, that it has run its course already.

It seems probable, however, that we may expect at least local recurrences in the near future, with an increase over the normal mortality from pneumonia for perhaps several years; and certainly we should be, as far as possible, prepared to meet them by previous organization of forces and measures for attempted prevention, treatment, and scientific investigation.

Future Study and Control.

As regards preventive measures, the efficacy of those carried out in recent months is not proved, and we can only continue to follow the apparently sound principles already applied. It seems hardly logical to expect that any measure short of effective specific immunization will afford lasting protection to the general population; but we may perhaps hope to delay the spread of infection, thus affording better facilities for treatment of the sick, and this is an achievement well worth while.

With reference to scientific investigations, these should not be conditioned on recurrence of the epidemic. Now is the time to inaugurate comprehensive laboratory and field investigations of influenza and pneumonia, to be continued not for a few months, but for a series of years, since a knowledge of influenza during the intervals between epidemics is essential to the understanding of epidemics. In this connection one of the most essential requisites for a better understanding of the disease is better differential diagnosis of endemic influenza with more careful observation and recording of the relatively mild indefinitely diagnosed epidemics of supposed influenza which are noted from time to time in the intervals between definite epidemics. Even in the absence of a definite diagnosis of individual cases, much would be added to the history of influenza by describing such outbreaks more fully and making them matters of more general record.

A RECENT PAMPHLET ON CANCER.

In response to repeated requests the United States Public Health Service has just issued a popular bulletin entitled "Cancer: Facts Which Every Adult Should Know." This 32-page, pocket-size pamphlet, constituting No. 6 of the "Keep Well Series" published by the United States Public Health Service, was prepared in cooperation with the foremost American authorities on cancer, and may therefore be accepted as representing the best scientific information on this important subject.

The bulletin gives in simple language the main facts regarding the prevalence of cancer, discusses three common causes of needless worry, tells how cancer develops, and describes the causative factors and the chief danger signals of the disease. Special attention is devoted to cancers of the breast, uterus, and digestive organs, and to the various forms of external cancer. An emphatic warning is given regarding alleged "cancer cures," and against quacks and patent nostrums. The use of radium and X-rays is discussed. Following is the summary which concludes the bulletin:

1. Cancer at the beginning is usually painless and its onset for this reason is especially insidious and dangerous.

2. Cancer is at first a small local growth which can be safely and easily removed by competent surgical or other treatment.
3. Cancer is not a constitutional or "blood" disease.
4. Cancer is not contagious.
5. Cancer is, practically speaking, not hereditary.
6. Every lump in the breast should be examined by a competent doctor.
7. Persistent abnormal discharge or bleeding is suspicious.
8. Sores, cracks, lacerations, lumps, and ulcers which do not heal, and warts, moles, or birthmarks which change in size, color, or appearance, may turn into cancer unless treated and cured.
9. Probably 60 per cent of cancers of the rectum are first regarded as piles. Insist on a thorough medical examination.
10. Continued irritation in some form is the usual cause of cancer. It rarely results from a sudden injury.
11. A doctor who treats a suspicious symptom without making a thorough examination does not know his business.

THE NOTIFIABLE DISEASES.

PREVALENCE DURING 1918 IN STATES.

ANTHRAX, CEREBROSPINAL MENINGITIS, DENGUE, DIPHTHERIA, INFLUENZA, MALARIA MEASLES, PNEUMONIA (ALL FORMS), POLIOMYELITIS, RABIES IN ANIMALS, RABIES IN MAN, ROCKY MOUNTAIN SPOTTED FEVER, SCARLET FEVER, SEPTIC SORE THROAT SMALLPOX, TUBERCULOSIS (PULMONARY AND ALL FORMS), TYPHOID FEVER, AND TYPHUS FEVER—CASES AND DEATHS REPORTED, CASE RATES, FATALITY AND MORTALITY RATES, 1918; AVERAGE NUMBER OF CASES REPORTED DURING RECENT YEARS (1913 TO 1917) AND CASE RATES.

The tables which appear on the following pages have been compiled from data furnished by the health officers of the several States in response to requests from the United States Public Health Service.

The following table shows States for which morbidity and mortality summaries were received:

Morbidity.	Mortality.	Morbidity.	Mortality.
Alabama.	Alabama.	Montana.	Missouri. ¹
Arizona.	Arizona.	Nebraska.	Nebraska.
Arkansas.	Arkansas.	New Jersey.	New Jersey.
California.	California.	New York.	New York.
Colorado.		North Carolina.	
Connecticut.	Connecticut.	Ohio.	Ohio.
District of Columbia.	District of Columbia.	Oregon.	Oklahoma.
Florida.	Florida.	Pennsylvania.	Oregon.
	Hawaii.	Porto Rico.	Pennsylvania.
Illinois.	Illinois.	Rhode Island.	Porto Rico.
Indiana.	Indiana.	South Carolina.	Rhode Island.
Iowa.	Iowa.	South Dakota.	South Carolina.
Kansas.	Kansas.	Texas.	South Dakota.
Louisiana.	Kentucky.	Utah.	Texas.
Maine.	Louisiana.	Vermont.	Utah.
Maryland.	Maine.	Virginia.	Vermont.
Massachusetts.	Maryland.	Washington.	Washington.
Michigan.	Massachusetts.	West Virginia.	
Minnesota.	Michigan.	Wisconsin.	
Mississippi.	Minnesota.		
	Mississippi.	Wyoming.	Wisconsin.
			Wyoming.

¹ Data not given by months.

The populations used in computing the case rates and death rates were those estimated by the Bureau of the Census as of July 1, 1918, except for the population of Iowa. For this State the population in 1910, as ascertained by the census of that year, was used, as no estimate for 1918 was available.

For most of the diseases four separate tables are given: (1) Averages, (2) Morbidity, (3) Mortality, and (4) Rates. The averages were obtained by adding together the cases reported by months for the years 1913, 1914, 1915, 1916, and 1917 and dividing the sums by 5. Data for the full five years were not available for some of the States. The column headed "Years" shows the years and number of years for which the figures were obtainable. In all instances the averages are for years between 1913 and 1917, inclusive. The averages are given by months for the following diseases: Cerebrospinal meningitis, diphtheria, measles, poliomyelitis, scarlet fever, smallpox, tuberculosis (pulmonary and all forms), and typhoid fever. The total cases of anthrax reported in each State during 1916 and 1917 are included in the rate table for this disease under a different heading.

In the morbidity tables are given the number of cases of communicable diseases reported in the States, by months, except rabies in animals and rabies in man, for which only totals are given. The morbidity for influenza is not complete, as this disease was not made reportable until the epidemic began in the fall of 1918. Neither are the pneumonia reports complete.

The mortality tables show deaths from communicable diseases registered in States by months.

The rate table shows the average cases reported per annum for the years 1913 to 1917, inclusive, where the data were available, and the average case rate per 1,000 inhabitants. This was obtained by using the average population of the years for which cases were reported. It also shows the cases and deaths reported; case rates, and fatality and mortality rates in 1918.

In comparing the figures for 1918 with the averages for preceding years, it should be borne in mind that an increase in the number of cases reported may indicate that there were more cases of the particular disease in the State during 1918, or it may indicate that the cases were better reported by the physicians.

A relatively large number of reported cases of a communicable disease, as indicated by a high case rate (and more especially when accompanied by a relatively small number of deaths, as indicated by a low fatality rate), usually means that the health department of that State is active and that the cases of the disease are being well reported by the practicing physicians. It does not necessarily mean that the disease is more prevalent in that State than in other States. A high fatality rate may mean that the disease was unusually virulent in a State, that the physicians did not treat the disease in that

State with the success usual elsewhere, or that the practicing physicians did not report all of their cases. On the other hand, an unusually low fatality rate may be due to the fact that the disease in the State was unusually mild, that the physicians treated it with unusual success; that the practicing physicians reported their cases satisfactorily, or that the registration of deaths was incomplete, or the assignment of the causes of death inaccurate.

Summaries of reported cases and deaths from communicable diseases similar to the summary here presented have been issued by the United States Public Health Service for the years 1912 to 1917, inclusive (Reprints Nos. 163, 208, 298, 345, 426, and 505). A comparison of these summaries indicates that, while morbidity reports in the United States are far from complete, steady advance is being made in many of the States.

Cases reported and deaths registered in New York City, during the calendar year 1918, from certain diseases, with indicated morbidity, mortality, and fatality rates.

	Cases.	Deaths.		Cases.	Deaths.
Anthrax:			Influenza:		
January.....	2	0	January.....	72	
February.....	1	0	February.....	52	
March.....	1	1	March.....	133	
April.....	1	1	April.....	106	
May.....	2	1	May.....	36	
June.....	1	0	June.....	7	
July.....	0	0	July.....	6	
August.....	0	0	August.....	5	
September.....	1	0	September.....	2,131	114
October.....	1	0	October.....	110,903	8,384
November.....	2	1	November.....	16,766	2,731
December.....	3	0	December.....	6,149	916
Total.....	15	4	Total.....		12,562
Rate per 1,000 population.....	.003	.001	Rate per 1,000 population.....		2,408
Deaths per 100 cases, 26.7.....					
Cerebrospinal meningitis:			Malaria:		
January.....	22	11	January.....	0	1
February.....	33	28	February.....	1	0
March.....	62	46	March.....	1	1
April.....	94	33	April.....	3	0
May.....	80	36	May.....	8	2
June.....	30	16	June.....	14	1
July.....	30	17	July.....	10	1
August.....	36	17	August.....	5	0
September.....	29	19	September.....	4	0
October.....	14	14	October.....	5	0
November.....	23	8	November.....	4	2
December.....	24	17	December.....	0	0
Total.....	477	262	Total.....	55	8
Rate per 1,000 population.....	.091	.030	Rate per 1,000 population.....	.011	.002
Deaths per 100 cases, 54.9.....			Deaths per 100 cases, 14.5.....		
Diphtheria:			Measles:		
January.....	1,048	112	January.....	4,264	85
February.....	898	101	February.....	4,477	98
March.....	1,086	138	March.....	6,345	173
April.....	1,249	160	April.....	6,003	161
May.....	1,288	121	May.....	4,536	121
June.....	1,231	106	June.....	1,812	79
July.....	893	80	July.....	664	33
August.....	541	52	August.....	241	16
September.....	472	45	September.....	114	10
October.....	801	120	October.....	76	8
November.....	836	90	November.....	67	1
December.....	1,112	120	December.....	76	5
Total.....	11,455	1,245	Total.....	28,675	790
Rate per 1,000 population.....	2.196	.239	Rate per 1,000 population.....	5.498	1.515
Deaths per 100 cases, 10.9.....			Deaths per 100 cases, 2.8.....		

Cases reported and deaths registered in New York City, during the calendar year 1918; from certain diseases, with indicated morbidity, mortality, and fatality rates.—Continued.

	Cases.	Deaths.		Cases.	Deaths.
Poliomyelitis:			Tuberculosis (pulmonary):		
January.....	3	0	January.....	1,538	842
February.....	3	2	February.....	1,344	771
March.....	8	4	March.....	1,366	1,061
April.....	10	3	April.....	1,698	860
May.....	1	1	May.....	1,300	743
June.....	13	4	June.....	1,759	643
July.....	26	4	July.....	873	595
August.....	33	6	August.....	1,326	546
September.....	24	3	September.....	1,002	546
October.....	9	1	October.....	721	826
November.....	2	1	November.....	707	711
December.....	2	0	December.....	805	635
Total.....	134	29	Total.....	14,439	8,779
Rate per 1,000 population.....	.026	.006	Rate per 1,000 population.....	2.768	1.683
Deaths per 100 cases, 21.6.			Number of cases reported for each death registered, 1.6.		
Pneumonia (all forms):			Tuberculosis (all forms):		
January.....	1,472		January.....	973	
February.....	1,304		February.....	884	
March.....	2,296		March.....	1,198	
April.....	1,747		April.....	993	
May.....	855		May.....	868	
June.....	445		June.....	765	
July.....	354		July.....	700	
August.....	313		August.....	653	
September.....	350	470	September.....	632	
October.....	15,230	7,184	October.....	926	
November.....	4,325	2,669	November.....	783	
December.....	2,757	1,519	December.....	724	
Total.....	20,628		Total.....	10,099	
Rate per 1,000 population.....	3.955		Rate per 1,000 population.....	1.936	
Scarlet fever:			Typhoid fever:		
January.....	579	30	January.....	30	6
February.....	546	22	February.....	28	8
March.....	634	16	March.....	35	13
April.....	579	26	April.....	53	6
May.....	512	25	May.....	51	10
June.....	321	20	June.....	164	13
July.....	206	13	July.....	170	30
August.....	105	5	August.....	265	30
September.....	164	1	September.....	209	32
October.....	244	8	October.....	125	28
November.....	206	4	November.....	52	10
December.....	364	7	December.....	56	10
Total.....	4,460	177	Total.....	1,238	196
Rate per 1,000 population.....	.855	.034	Rate per 1,000 population.....	.237	.038
Deaths per 100 cases, 4.0.			Deaths per 100 cases, 15.8.		
Smallpox:			Typhus fever:		
January.....	1		January.....	0	
February.....	3		February.....	0	
March.....	8		March.....	0	
April.....	3		April.....	1	
May.....	3		May.....	3	
June.....	2		June.....	2	
July.....	1		July.....	0	
August.....	0		August.....	1	
September.....	0		September.....	1	
October.....	0		October.....	0	
November.....	0		November.....	0	
December.....	0		December.....	0	
Total.....	21		Total.....	8	
Rate per 1,000 population.....	.004		Rate per 1,000 population.....	.002	

Cases of anthrax reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
California.....	4				1	3	1		6	3	1		19	California,
Florida.....	1												1	Florida.
Kansas.....	1												1	Kansas.
Louisiana.....	1	1	2	0	0	7	1	0	1	0	0	0	13	Louisiana.
Maryland.....			1	2	1								1	Maryland.
Massachusetts.....	1	5	2	1		3	4	1		3			23	Massachusetts.
New Jersey.....								6		1	1		10	New Jersey.
New York.....	3	2	1	3	2	1	1	0	1	1	2	3	20	New York.
Ohio.....													2	Ohio.
Pennsylvania.....	1			1		2	1			1		1	11	Pennsylvania.
Porto Rico.....						1	1			1		1	2	Porto Rico.
Texas.....							1						3	Texas.

Deaths registered as due to anthrax, 1918.

State.	Jun.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
California.....		1							1	1	1	1	5	California,
Connecticut.....									1				1	Connecticut.
Florida.....			1										1	Florida.
Hawaii.....		1											2	Hawaii.
Indiana.....				1					1				3	Indiana.
Iowa.....									1	2			3	Iowa.
Kansas.....			1								1		2	Kansas.
Louisiana.....		1	1						1				5	Louisiana.
Maine.....		1											1	Maine.
Massachusetts.....		1			1				2		1		1	Massachusetts.
New Jersey.....								1					1	New Jersey.
New York.....		1			1				2				8	New York.
Ohio.....					1								1	Ohio.
Pennsylvania.....										1			7	Pennsylvania.
Porto Rico.....										1			4	Porto Rico.
South Carolina.....											1		1	South Carolina.
Texas.....							1				1		4	Texas.

Anthrax—Cases reported 1916 and 1917; cases reported, deaths registered, and indicated morbidity, mortality, and fatality rates, 1918.

Cerebrospinal meningitis—Monthly averages of cases reported during the years for which data are available, 1913-1917.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Average per year.	State.
Alabama.....	1913-1915-1917-4 years.....	10	11	8	14	10	6	4	2	2	3	3	6	70	Alabama.
Arizona.....	1914-1 year.....	1	2	3	0	0	0	0	0	0	1	1	1	8	Arizona.
Arkansas.....	1917-1 year.....	2	0	1	0	2	0	1	1	0	1	2	1	12	Arkansas.
California.....	1913-1917-5 years.....	8	5	10	7	6	5	9	5	2	5	6	6	8	California.
Connecticut.....	do.....	6	9	16	21	13	10	4	5	5	3	2	3	96	Connecticut.
District of Columbia.....	do.....	(1)	1	2	(1)	2	(1)	(1)	(1)	(1)	(1)	(1)	(1)	2	District of Columbia.
Hawaii.....	1913-1914-1917-3 years.....	20	14	25	30	31	25	29	21	23	19	18	10	8	Hawaii.
Illinois.....	1914-1917-4 years.....	8	14	12	10	7	6	2	3	4	3	2	5	274	Illinois.
Indiana.....	1913-1914-1916-1917-4 years.....	3	3	4	5	2	1	1	2	1	5	2	3	66	Indiana.
Iowa.....	1913-1917-5 years.....	5	6	8	8	7	3	2	4	3	4	3	4	32	Iowa.
Kansas.....	1913-1917-4 years.....	4	6	4	4	1	1	1	2	(1)	1	2	3	9	Kansas.
Louisiana.....	1914-1917-4 years.....	0	2	0	0	0	0	0	0	0	0	1	1	36	Louisiana.
Maine.....	1917-1 year.....	4	4	8	14	11	8	4	3	3	3	4	8	6	Maine.
Maryland.....	1915-1917-3 years.....	13	10	15	17	23	10	13	13	11	13	15	15	86	Maryland.
Massachusetts.....	1913-1916-1917-3 years.....	13	10	15	17	23	10	13	13	11	13	15	15	175	Massachusetts.
Minnesota.....	1913-1917-5 years.....	2	9	19	23	16	8	5	3	2	3	2	4	96	Minnesota.
Mississippi.....	1913-1917-4 years.....	4	6	6	3	5	2	3	4	4	5	4	5	49	Mississippi.
Montana.....	1913-1914-1916-1917-4 years.....	(1)	1	1	1	1	2	(1)	2	1	1	1	1	12	Montana.
Nevada.....	1913-1915-1917-3 years.....	2	1	0	1	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	8	Nevada.
New York.....	1913-1917-5 years.....	26	29	39	38	49	44	30	24	27	25	21	24	376	New York.
Ohio.....	1914-1917-4 years.....	13	22	38	39	40	25	19	16	11	17	15	17	292	Ohio.
Oregon.....	1914-1916-1917-3 years.....	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	6	Oregon.
Pennsylvania.....	1916-1917-2 years.....	17	24	76	116	93	70	26	33	20	20	16	20	51	Pennsylvania.
Porto Rico.....	1917-1 year.....	0	0	2	2	0	0	1	0	1	0	1	0	5	Porto Rico.
Rhode Island.....	1916-1917-2 years.....	3	2	2	8	5	3	8	2	1	4	2	2	42	Rhode Island.
South Carolina.....	1913-1917-5 years.....	2	1	2	2	1	4	1	1	1	1	1	1	44	South Carolina.
South Dakota.....	1913-1914-1917-3 years.....	0	1	1	1	1	1	1	1	1	1	2	1	13	South Dakota.
Texas.....	1915-1917-2 years.....	7	9	8	5	5	8	4	2	3	1	2	9	63	Texas.
Vermont.....	1916-1917-2 years.....	0	3	(1)	0	(1)	2	(1)	0	0	0	0	0	6	Vermont.
Virginia.....	1915-1917-3 years.....	20	17	25	22	14	11	19	12	15	10	10	14	198	Virginia.
Washington.....	1913-1917-5 years.....	1	1	1	2	3	2	1	1	1	1	2	1	18	Washington.
West Virginia.....	1917-1 year.....	7	6	5	10	2	1	5	1	1	1	0	2	40	West Virginia.
Wisconsin.....	1913-1915-1917-5 years.....	10	8	13	11	9	12	6	7	6	6	6	6	101	Wisconsin.
Wyoming.....	1913-1914-1916-1917-4 years.....	0	(1)	1	(1)	0	(1)	0	(1)	0	(1)	0	2	6	Wyoming.

¹ Less than one.

Cases of cerebrospinal meningitis reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	22	52	34	15	20	9	8	2	12	1	4	9	188	Alabama.
Arkansas.....	45	55	47	16	7	2	1	1	13	13	13	13	204	Arkansas.
California.....	27	26	32	29	24	10	18	17	18	11	6	7	225	California.
Colorado.....	3	1	2	2	2	17	17	17	3	3	3	3	9	Colorado.
Connecticut.....	21	10	12	15	11	9	7	5	5	9	5	3	118	Connecticut.
District of Columbia.....	37	31	11	8	16	5	1	0	1	4	4	0	118	District of Columbia.
Florida.....	5	29	14	3	5	6	3	1	1	3	3	3	75	Florida.
Illinois.....	48	53	69	62	53	19	20	14	16	8	4	12	389	Illinois.
Indiana.....	8	16	35	21	19	8	6	4	7	11	1	4	140	Indiana.
Iowa.....	8	12	9	5	2	3	3	1	1	5	0	1	50	Iowa.
Kansas.....	32	53	56	23	24	11	3	7	2	4	0	16	231	Kansas.
Louisiana.....	139	64	48	23	21	18	11	6	4	17	11	17	394	Louisiana.
Maine.....	0	0	4	0	0	1	0	0	7	0	2	2	12	Maine.
Maryland.....	18	29	35	43	27	12	17	13	7	22	14	17	244	Maryland.
Massachusetts.....	21	35	51	67	38	36	25	18	18	31	19	21	378	Massachusetts.
Michigan.....	11	14	11	8	8	9	9	7	4	8	2	4	95	Michigan.
Minnesota.....	10	10	13	10	5	6	1	6	3	1	1	2	68	Minnesota.
Mississippi.....	15	40	25	21	5	3	2	1	1	0	4	2	125	Mississippi.
Montana.....	3	2	2	2	1	3	3	3	1	2	1	1	18	Montana.
Nebraska.....	4	4	7	7	7	7	7	7	7	7	7	7	28	Nebraska.
New Jersey.....	22	24	33	47	35	17	20	21	6	18	15	7	265	New Jersey.
North Carolina.....	43	51	90	127	99	43	40	48	41	41	33	38	670	New York.
Ohio.....	13	25	21	8	7	5	7	6	4	5	3	11	123	North Carolina.
Oregon.....	49	51	57	44	44	23	21	18	31	14	14	8	32	Oregon.
Pennsylvania.....	27	42	79	92	95	29	28	31	19	15	21	13	491	Pennsylvania.
Rhode Island.....	7	7	8	13	11	6	4	1	3	1	2	1	64	Rhode Island.
South Carolina.....	108	107	82	31	11	6	5	5	5	2	1	1	359	South Carolina.
South Dakota.....	3	6	4	5	4	1	4	2	3	4	5	4	45	South Dakota.
Texas.....	46	65	34	15	4	7	3	4	2	2	2	4	188	Texas.
Vermont.....	1	2	2	2	2	1	1	1	1	1	1	1	5	Vermont.
Virginia.....	53	38	51	33	28	9	9	12	15	28	13	20	314	Virginia.
Washington.....	5	8	10	9	10	7	5	1	2	4	7	7	66	Washington.
West Virginia.....	3	0	4	3	3	0	1	4	0	2	4	0	24	West Virginia.
Wisconsin.....	14	14	25	27	26	17	16	10	6	11	4	11	191	Wisconsin.
Wyoming.....	1	4	1	1	1	2	1	1	1	1	1	4	15	Wyoming.

Deaths registered as due to cerebrospinal meningitis, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	13	9	18	11	6	5	4	3	0	1	6	6	82	Alabama.
Arizona.....	3	2	11	7	5	2	3	0	1	5	5	4	51	Arizona.
Arkansas.....	27	37	27	17	32	8	10	8	3	14	12	30	225	Arkansas.
California.....	13	19	18	24	12	10	10	10	3	17	17	2	11 ^a	California.
Connecticut.....	7	5	4	5	7	7	3	2	1	1	1	2	44	Connecticut.
District of Columbia.....	4	7	4	6	2	2	2	4	1	3	3	2	28	District of Columbia.
Florida.....	0	0	0	4	2	0	2	4	0	0	2	1	2	Florida.
Hawaii.....	29	16	28	18	14	12	15	8	4	5	4	11	11	Hawaii.
Illinois.....	3	3	6	11	4	3	5	4	1	2	1	2	164	Illinois.
Indiana.....	9	14	7	5	11	2	5	2	2	4	2	2	45	Indiana.
Iowa.....	14	26	22	15	5	3	9	4	4	4	12	8	69	Iowa.
Kansas.....	51	74	70	62	38	45	30	35	26	44	36	25	331	Kansas.
Kentucky.....	21	19	12	7	4	5	2	3	0	4	2	3	82	Kentucky.
Louisiana.....	2	1	5	5	5	2	2	3	2	5	2	1	30	Louisiana.
Maine.....	11	8	17	15	12	7	7	7	1	15	5	3	106	Maine.
Maryland.....	16	23	30	34	20	20	15	8	12	12	14	20	224	Maryland.
Massachusetts.....	28	30	21	33	36	28	21	12	15	18	11	15	151	Massachusetts.
Michigan.....	2	2	5	2	1	2	2	2	2	2	1	1	23	Michigan.
Minnesota.....	4	3	11	8	1	2	2	2	2	2	0	0	4	Minnesota.
Nebraska.....	3	1	4	7	4	7	2	1	3	4	1	1	37	Nebraska.
New Jersey.....	7	4	11	17	11	1	7	4	1	1	1	1	4	New Jersey.
New York.....	28	41	63	43	65	30	27	25	26	40	22	33	443	New York.
Ohio.....	13	21	32	21	17	16	10	12	11	5	5	4	161	Ohio.
Oklahoma.....	3	6	6	4	1	2	0	0	1	0	0	1	31	Oklahoma.
Oregon.....	0	0	0	3	1	2	0	0	1	0	0	1	8	Oregon.
Rhode Island.....	3	1	4	7	4	7	2	1	3	4	1	1	38	Rhode Island.
South Carolina.....	33	40	39	22	5	15	10	3	5	20	4	11	207	South Carolina.
South Dakota.....	2	6	4	5	4	4	2	2	3	4	5	4	43	South Dakota.
Texas.....	31	46	27	16	5	8	7	9	2	4	5	4	164	Texas.
Utah.....	1	2	1	1	1	1	1	1	1	1	1	1	7	Utah.
Vermont.....	9	4	10	9	11	3	1	1	2	1	1	1	3	Vermont.
Washington.....	31	33	37	42	35	26	23	7	1	1	15	3	81	Washington.
Wisconsin.....											30	51	365	Wisconsin.
Wyoming.....											1	4	12	Wyoming.

Cerebrospinal meningitis—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated morbidity, mortality, and fatality rates, 1918.

State.	Estimated Population July 1, 1918.	Average.		1918		State.		
		Years.	Cases reported per annum.	Cases reported per 1,000 inhabitants.	Total cases reported.	Cases per 1,000 inhabitants.	Deaths per 1,000 inhabitants.	Fatalities per 100 cases.
Alabama.....	2,385,270	1913, 1915-1917—4 years.....	79	0.034	188	0.078	82	0.034
Arizona.....	1,722,034	1914—1 year.....	8	.033	...	51	187	43.6
Arkansas.....	1,722,965	1917—1 year.....	12	.007	204	.225	125	...
California.....	3,119,412	1913-1917—5 years.....	76	.027	225	.072	160	.051
Colorado.....	1,014,581	1913-1917—5 years.....	96	.073	118	.092	44	...
Connecticut.....	1,286,268	1913-1917—5 years.....	11	.031	118	.284	28	0.04
District of Columbia.....	401,681	do.	96	.073	75	.080	37	0.70
Florida.....	938,877	1913-1917—5 years.....	8	.033	...	11	11	23.7
Hawaii.....	223,410	1913-1917—5 years.....	274	.045	389	.062	164	...
Illinois.....	6,177,734	1913, 1914, 1917—3 years.....	66	.024	140	.049	45	42.2
Indiana.....	2,884,167	1914-1917—4 years.....	32	.014	50	.104	69	0.016
Kansas.....	2,224,771	1913, 1914, 1916, 1917—4 years.....	62	.034	231	.123	131	32.1
Louisiana.....	1,874,195	1913-1917—5 years.....	408	.547	...	538	223	...
Kentucky.....	1,894,778	1914-1917—4 years.....	36	.020	394	.029	82	56.7
Maine.....	782,191	1917—1 year.....	6	.008	26	.30	30	...
Maryland.....	1,334,539	1915-1917—3 years.....	86	.063	244	.176	108	44.3
Massachusetts.....	3,832,790	1913, 1916, 1917—3 years.....	176	.048	375	.099	224	59.3
Michigan.....	3,133,678	1913-1917—5 years.....	96	.043	95	.029	268	...
Minnesota.....	3,345,287	1913-1917—5 years.....	49	.026	68	.020	23	0.086
Mississippi.....	2,001,466	1914-1917—4 years.....	49	.026	125	.062	6	0.010
Missouri.....	3,448,398	1913-1917—5 years.....	12	.027	18	.037	107	33.8
Montana.....	436,376	1913, 1914, 1916, 1917—4 years.....	26	4.8
Nebraska.....	1,296,877	1913, 1915, 1917—3 years.....	8	.078	...	37	.029	...
Nevada.....	114,742	1914, 1916, 1917—3 years.....	6
New Jersey.....	3,080,371	1913-1917—5 years.....	376	.087	265	.086	69	0.022
New York.....	10,646,089	1914-1917—4 years.....	531	.062	491	.083	443	26.0
North Carolina.....	2,468,025	1914-1917—4 years.....	292	.057	342	.086	161	0.042
Ohio.....	5,273,814	1914-1917—4 years.....	5	31	.031	44.1
Oklahoma.....	2,377,629	1914, 1916, 1917—3 years.....	6	8	.009	...
Oregon.....	888,243	1914, 1916, 1917—3 years.....	6
Pennsylvania.....	8,758,067	1914, 1916, 1917—3 years.....	5
Puerto Rico.....	1,247,677	1917—1 year.....	5
Rhode Island.....	1,361,155	1916, 1917—5 years.....	64	.004
South Carolina.....	1,660,934	1913-1917—5 years.....	44	.027	359	.216	207	67.7

August 25, 1912.

Dengue—Cases reported and deaths registered, 1918.

Month	California:		Texas:		Deaths.
	Cases.	Deaths.	Louisiana:	Cases.	
October.....	1	0	February.....	1	13
November.....	1	0	May.....	72	1
December.....	1	0	June.....	1	3
Florida:			Porto Rico:	0	96
February.....	8	0	September.....	145	9
September.....	1	0	October.....	200	1
October.....	2	0	November.....	2
December.....	1	0	August.....	3

Diphtheria.—Monthly averages of cases reported during the years for which data are available, 1913-1917.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Average per year.	State.	
Alabama.....	1913-1915-1917-4 years.....	41	31	35	25	26	14	20	60	123	124	91	57	647	Alabama.	
Arizona.....	1914-1916-2 years.....	2	3	4	4	4	2	3	1	1	5	9	40	40	Arlington.	
Arkansas.....	1917-1 Year.....	10	13	7	6	4	4	14	40	53	33	75	288	Arkansas.		
California.....	1913-1917-5 years.....	29	26	191	20	200	133	142	142	250	321	323	2,731	California.		
Colorado.....	1914-1916-1917-3 years.....	272	239	148	41	31	30	30	449	66	65	65	70	566	Colorado.	
Connecticut.....	1913-1916-5 years.....	204	199	181	165	164	152	128	96	151	216	301	266	2,343	Connecticut.	
District of Columbia.....	do.....	67	40	55	45	37	20	18	16	36	121	128	98	681	District of Columbia.	
Hawaii.....	1913-1917-5 years.....	16	15	48	15	9	10	7	11	15	10	9	13	178	Hawaii.	
Idaho.....	1913-1917-5 years.....	2	6	1	3	0	5	4	4	0	2	1	1	29	Idaho.	
Iowa.....	1913-1914-1917-3 years.....	1,366	933	1,117	903	922	828	715	575	889	1,123	1,579	1,287	12,247	Iowa.	
Illinois.....	1913-1917-5 years.....	337	255	189	153	138	106	116	166	330	632	623	443	3,228	Illinois.	
Indiana.....	1913-1917-5 years.....	75	69	66	56	49	44	34	28	44	103	96	88	752	Indiana.	
Kansas.....	1913-1917-5 years.....	130	90	85	65	63	52	47	56	86	303	354	233	1,864	Kansas.	
Louisiana.....	1914-1917-4 years.....	113	78	71	53	38	33	32	32	155	186	126	87	1,054	Louisiana.	
Maine.....	1916-1917-2 years.....	31	20	15	19	12	19	21	21	25	48	63	62	354	Maine.	
Maryland.....	1915-1917-3 years.....	235	175	163	89	90	68	72	89	167	283	319	251	2,007	Maryland.	
Massachusetts.....	1913-1916-1917-3 years.....	771	638	714	638	668	612	474	464	516	790	827	918	8,113	Massachusetts.	
Michigan.....	1912-1914-5 years.....	500	378	426	365	368	333	296	305	389	681	775	694	5,480	Michigan.	
Minnesota.....	1911-1912-4 years.....	314	255	247	233	297	220	204	227	231	396	377	295	3,306	Minnesota.	
Mississippi.....	1911-1917-4 years.....	78	54	37	29	23	34	42	42	93	153	212	173	1,057	Mississippi.	
Montana.....	1913-1914-1916-1917-4 years.....	16	16	20	15	17	15	20	13	17	23	22	22	228	Montana.	
Nevada.....	1913-1915-1917-3 years.....	1	(1)	0	1	1	(1)	2	1	1	2	1	1	10	Nevada.	
New Jersey.....	1913-1917-5 years.....	630	527	517	444	520	456	368	300	404	658	763	674	6,261	New Jersey.	
New York.....	do.....	2,631	1,921	1,924	1,740	1,020	1,850	1,493	989	1,118	1,719	1,780	1,964	20,361	New York.	
North Dakota.....	1917-1 Year.....	33	36	49	24	23	33	46	34	61	27	78	95	639	North Dakota.	
Ohio.....	1914-1917-4 years.....	882	649	600	472	466	409	350	436	49	154	276	1,324	939	8,551	Ohio.
Oklahoma.....	1913-1915-1916-1917-3 years.....	112	72	58	47	31	22	22	49	154	276	230	220	2,739	Oklahoma.	
Oregon.....	1913-1915-1917-2 years.....	32	21	24	21	21	22	15	17	16	25	25	39	38	291	Oregon.
Pennsylvania.....	1916, 1917-2 years.....	1,451	1,239	1,169	971	998	870	829	909	1,374	1,948	1,929	1,550	15,297	Pennsylvania.	
Puerto Rico.....	1914, 1915, 1917-3 years.....	15	12	12	7	13	8	13	16	9	18	11	9	143	Puerto Rico.	
Rhode Island.....	1913-1914-1916-1917-3 years.....	113	103	101	75	63	61	74	66	125	110	97	97	1,053	Rhode Island.	
South Carolina.....	1913-1917-5 years.....	97	89	65	71	50	51	56	160	316	348	242	202	1,747	South Carolina.	
South Dakota.....	1913-1914-1916-1917-3 years.....	33	23	24	24	17	20	16	16	13	36	36	36	254	South Dakota.	
Texas.....	1915, 1917-2 years.....	156	110	74	62	46	41	69	53	139	355	456	277	1,871	Texas.	
Utah.....	1913-1915, 1917-4 years.....	33	24	23	23	17	22	17	20	29	42	38	336	Utah.		
Vermont.....	1914-1917-4 years.....	42	27	25	33	33	28	25	24	25	66	67	52	447	Vermont.	
Virginia.....	1913-1917-5 years.....	262	244	120	86	64	53	225	445	659	618	357	3,188	Virginia.		
Washington.....	do.....	48	40	37	38	35	28	27	33	35	45	45	45	4,333	Washington.	
West Virginia.....	1917-1 year.....	82	55	43	43	36	29	18	70	125	173	73	927	927	West Virginia.	
Wisconsin.....	1913-1917-5 years.....	25	167	198	180	160	160	158	140	248	250	250	217	2,251	Wisconsin.	
Wyoming.....	do.....	3	4	5	1	(1)	2	1	6	6	6	6	6	46	46	Wyoming.

¹Less than one.

Cases of diphtheria reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sep.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	53	39	40	26	16	17	60	126	150	120	57	43	747	Alabama.
Arizona.....	10	1	3	1	3	0	6	7	1	2	1	1	38	Arizona.
Arkansas.....	17	19	15	16	19	5	5	12	13	14	11	27	173	Arkansas.
California.....	354	280	306	284	290	246	222	188	187	381	169	210	3,117	California.
Colorado.....	68	36	48	62	86	62	53	36	58	46	23	222	620	Colorado.
Connecticut.....	322	173	203	208	132	159	99	204	248	209	259	259	2,348	Connecticut.
District of Columbia.....	57	37	49	55	72	44	23	13	21	31	48	67	747	District of Columbia.
Florida.....	25	31	32	8	10	19	13	14	36	43	55	43	329	Florida.
Illinois.....	1,028	649	751	655	585	586	487	382	604	990	703	705	8,075	Illinois.
Indiana.....	382	334	301	284	201	155	176	183	222	173	216	230	2,839	Indiana.
Iowa.....	88	78	73	74	68	52	62	51	78	115	98	76	911	Iowa.
Kansas.....	214	88	128	138	92	57	59	48	116	118	99	113	1,293	Kansas.
Louisiana.....	84	84	51	52	96	293	194	145	79	42	41	35	1,201	Louisiana.
Maine.....	20	15	54	15	9	15	17	30	15	29	29	14	262	Maine.
Maryland.....	165	125	92	93	91	59	54	58	88	152	122	162	1,261	Maryland.
Massachusetts.....	950	678	744	667	663	473	453	337	430	410	532	584	6,921	Massachusetts.
Michigan.....	593	498	482	379	368	342	454	302	455	683	637	700	5,912	Michigan.
Minnesota.....	314	211	339	362	303	255	232	226	221	411	321	365	3,600	Minnesota.
Mississippi.....	72	54	35	35	49	50	49	103	105	109	104	82	1,138	Mississippi.
Montana.....	58	25	21	30	51	19	23	15	14	16	21	11	309	Montana.
Nebraska.....	55	34	21	57	104	69	51	36	80	80	32	32	706	Nebraska.
New Jersey.....	413	361	413	485	312	303	341	204	324	374	422	483	4,465	New Jersey.
New York.....	1,543	1,268	1,554	1,649	1,690	1,587	1,226	821	819	1,290	1,300	1,754	16,501	New York.
North Carolina.....	75	51	49	44	40	38	38	143	143	233	171	97	1,327	North Carolina.
North Dakota.....	39	38	49	52	61	31	30	24	38	11	9	9	391	North Dakota.
Ohio.....	618	619	607	385	333	361	330	338	459	549	509	389	5,807	Ohio.
Oregon.....	7	15	10	8	14	22	11	13	13	17	21	25	12,182	Oregon.
Pennsylvania.....	1,003	1,049	1,046	1,002	850	824	779	752	1,085	1,568	1,118	1,291	12,378	Pennsylvania.
Porto Rico.....	7	8	11	11	17	7	10	19	21	10	10	7	138	Porto Rico.
Rhode Island.....	93	65	79	95	70	74	55	29	70	63	72	130	885	Rhode Island.
South Carolina.....	99	45	68	93	51	8	50	84	221	79	147	94	1,039	South Carolina.
South Dakota.....	26	15	17	24	7	16	30	27	14	44	26	27	273	South Dakota.
Texas.....	126	100	107	61	76	50	48	76	70	38	42	39	853	Texas.
Utah.....	46	41	52	45	21	47	48	45	44	86	69	51	594	Utah.
Vermont.....	17	11	11	18	8	9	8	38	38	38	19	35	241	Vermont.
Virginia.....	121	111	98	77	37	38	42	132	113	272	199	129	1,367	Virginia.
Washington.....	93	173	58	45	57	38	33	38	33	124	53	57	882	Washington.
West Virginia.....	51	28	19	28	13	13	31	20	45	25	36	32	345	West Virginia.
Wisconsin.....	241	156	138	130	171	133	146	100	138	160	125	125	1,773	Wisconsin.
Wyoming.....	20	50	18	18	25	13	6	11	24	7	6	7	292	Wyoming.

August 15, 1918.

1850

Deaths registered as due to diphtheria, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sep.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	30	10	11	8	2	3	12	26	27	38	18	10	185	Alabama.
Arizona.....	1	2	3	4	1	4	4	8	9	18	1	1	17	Arizona.
Arkansas.....	19	7	3	7	3	4	4	8	9	18	5	5	17	Arkansas.
California.....	21	18	19	18	18	17	13	9	10	18	16	16	204	California.
Connecticut.....	19	12	25	8	13	10	9	10	18	29	21	21	201	Connecticut.
Delaware.....	5	0	2	1	2	1	0	1	2	8	3	3	27	Delaware.
District of Columbia.....	4	1	1	1	4	4	1	1	1	1	3	4	14	District of Columbia.
Florida.....	10	6	8	3	2	2	5	5	6	12	14	9	86	Florida.
Hawaii.....	3	3	3	2	1	2	1	2	1	2	1	1	17	Hawaii.
Illinois.....	144	105	103	100	82	65	69	61	77	154	103	89	1,152	Illinois.
Indiana.....	63	45	42	25	14	21	23	23	23	35	40	39	417	Indiana.
Iowa.....	15	11	16	14	13	8	13	12	26	33	25	25	211	Iowa.
Kansas.....	10	9	14	18	10	4	1	1	1	8	13	16	120	Kansas.
Kentucky.....	27	14	20	8	9	7	5	19	36	37	35	35	234	Kentucky.
Louisiana.....	16	12	5	16	1	5	4	6	9	7	15	6	102	Louisiana.
Maine.....	11	8	5	5	1	1	1	4	6	3	3	1	60	Maine.
Maryland.....	15	13	8	14	9	2	5	4	5	34	13	15	137	Maryland.
Massachusetts.....	90	54	55	68	40	30	34	25	35	57	34	72	594	Massachusetts.
Michigan.....	78	67	67	38	41	29	46	25	36	78	72	81	658	Michigan.
Minnesota.....	16	16	26	18	33	21	19	16	23	42	26	18	274	Minnesota.
Nebraska.....	17	6	8	8	9	8	6	6	9	11	6	9	103	Nebraska.
New Jersey.....	54	43	46	40	33	28	27	18	36	56	50	54	485	New Jersey.
New York.....	172	150	173	206	160	134	105	79	83	189	140	181	1,772	New York.
North Dakota.....	4	6	8	2	2	2	2	2	2	2	2	2	35	North Dakota.
Ohio.....	61	55	50	36	34	22	19	23	42	56	49	50	497	Ohio.
Oklahoma.....	7	15	13	7	7	3	3	7	7	21	14	19	124	Oklahoma.
Oregon.....	1	2	0	1	1	1	0	2	2	4	5	6	27	Oregon.
Pennsylvania.....	192	129	159	131	108	112	94	73	133	257	194	108	1,770	Pennsylvania.
Porto Rico.....	8	3	3	6	3	1	0	11	3	5	3	3	45	Porto Rico.
Rhode Island.....	13	8	12	13	10	2	3	3	9	9	11	11	100	Rhode Island.
South Carolina.....	3	2	5	6	4	3	4	8	11	20	10	5	81	South Carolina.
South Dakota.....	2	2	2	2	2	2	3	2	3	4	1	4	25	South Dakota.
Texas.....	31	30	25	12	12	6	7	9	10	20	15	20	197	Texas.
Utah.....	7	7	6	7	5	5	4	4	6	5	7	5	68	Utah.
Vermont.....	3	3	1	1	3	2	2	2	2	4	18	1	23	Vermont.
Washington.....	28	18	4	5	8	4	6	3	5	4	18	1	88	Washington.
Wisconsin.....	25	25	14	23	25	17	25	20	14	27	22	20	250	Wisconsin.
Wyoming.....	2	5	4	2	2	2	1	1	1	3	1	2	24	Wyoming.

Diphtheria—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated mortality, morbidity, and fatality rates, 1918.

State.	Estimated population July 1, 1918.	Average.		1918		Deaths per 1,000 inhabitants.	Fatalities per 100 cases.	State.	
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.				
Alabama.....	2,385,270	1913, 1915-1917—4 years.....	647	0.28	.747	0.31	.195	26.1 Alabama.	
Arizona.....	2,272,034	1914, 1916—2 years.....	40	.16	.38	.14	.17	44.7 Arizona.	
Arkansas.....	1,792,965	1917—1 year.....	288	.16	.173	.10	.104	60.1 Arkansas.	
California.....	3,119,412	1913-1917—5 years.....	2,731	.96	.3,117	1.00	.204	6.5 California.	
Colorado.....	1,014,581	1914, 1916, 1917—3 years.....	506	.59	.620	.61	6.5 Colorado.	
Connecticut.....	1,286,268	1913-1917—5 years.....	2,343	1.91	2,348	1.83	201	3.6 Connecticut.
Delaware.....	216,041	1913-1917—3 years.....	681	1.90	517	1.29	27	12 Delaware.
District of Columbia.....	401,081	1913-1917—3 years.....	681	1.90	329	.35	40	10	7.8 District of Columbia.
Florida.....	938,877	1913-1917—5 years.....	178	.94	17	08 Florida.
Hawaii.....	223,410	1913-1917—5 years.....	12,247	2.03	8,075	1.28	1,102	18 Hawaii.
Idaho.....	461,766	1913, 1914, 1915-1917—3 years.....	752	1.26	2,839	.99	417	15 Idaho.
Illinois.....	6,317,734	1913-1917—5 years.....	3,528	1.06	911	.99	211	16	14.3 Illinois.
Indiana.....	5,843,167	1913-1917—5 years.....	3,528	1.06	2,839	.99	128	10	23.2 Indiana.
Iowa.....	2,224,771	1913, 1914, 1916, 1917—4 years.....	752	.34	.91	.41	234	07	10.0 Iowa.
Kansas.....	1,874,195	1913-1917—5 years.....	1,564	.87	1,293	.69	102	06 Kansas.
Kentucky.....	2,408,547	1913-1917—5 years.....	1,054 Kentucky.	
Louisiana.....	1,884,778	1914-1917—4 years.....	1,054	.58	1,201	.64	102	06	6.5 Louisiana.
Maine.....	782,191	1916, 1917—2 years.....	1,354	.16	2,262	.33	60	08	22.9 Maine.
Maryland.....	1,384,539	1915, 1917—3 years.....	2,007	1.47	1,261	.91	137	10	10.9 Maryland.
Massachusetts.....	3,832,790	1913, 1916, 1917—3 years.....	8,113	2.20	6,921	1.81	594	15	8.6 Massachusetts.
Michigan.....	3,133,678	1913-1917—5 years.....	5,380	1.82	5,912	1.89	658	21	11.1 Michigan.
Minnesota.....	2,345,287	do.....	3,306	1.47	3,600	1.53	274	12	7.6 Minnesota.
Mississippi.....	2,001,466	1914-1917—4 years.....	1,057	.55	1,138	.57	137	07	12.0 Mississippi.
Missouri.....	3,445,408	1913, 1914, 1916, 1917—4 years.....	228	.51	309	.64	364	11 Missouri.
Montana.....	1,298,877	1913, 1915, 1917—3 years.....	10	706	.64	103	.08	14.6 Montana.
Nevada.....	114,742	1913-1917—5 years.....	6,261	2.17	4,485	1.45	485	16 Nevada.
New Jersey.....	3,080,371	1913-1917—5 years.....	20,361	2.02	16,501	1.55	1,772	17	10.9 New Jersey.
New York.....	10,616,989	do.....	1,327	1,327	10.7 New York.	
North Carolina.....	2,468,025	1917—1 year.....	1,327	1,327 North Carolina.	
North Dakota.....	709,427	1914-1917—4 years.....	8,339	1.70	5,807	1.04	35	04	9.0 North Dakota.
Oklahoma.....	5,775,814	1913, 1915, 1917—3 years.....	8,351	1.67	124	9.0 Oklahoma.	
Oregon.....	2,371,629	1913-1917—5 years.....	1,279	.60	182	.20	27	05 Oregon.
Pennsylvania.....	8,798,067	1916, 1917—2 years.....	15,267	1.78	12,378	1.41	1,770	.20	14.3 Pennsylvania.

Diphtheria—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated mortality, morbidity, and fatality rates, 1918—Continued.

State.	Estimated population July 1, 1918.	Average.		1918			State.
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.	Deaths per 1,000 inhabitants.	
Porto Rico.....	1,247,877	1914, 1915, 1916, 1917—3 years.....	143	12	.11	.55	.04
Rhode Island.....	415	1911, 1916, 1917—3 years.....	1,063	1,77	1.40	.10	39.9
South Carolina.....	637,934	1913—1917—5 years.....	1,447	1,66	.83	.06	11.2
South Dakota.....	733,434	1913, 1914, 1917—3 years.....	254	.38	.37	.25	7.8
Texas.....	4,601,279	1915, 1917—2 years.....	1,371	.32	.18	.197	23.6
Utah.....	455,648	1913—1915, 1917—4 years.....	336	.80	.594	1.31	11.5
Vermont.....	369,192	1914—1917—4 years.....	447	1.23	.241	.66	9.5
Virginia.....	2,237,620	1913—1917—6 years.....	3,188	1.47	1,387	.61
Washington.....	1,660,678	do.....	453	.31	.862	.92	9.6
West Virginia.....	1,438,165	1917—1 year.....	927	.66	.345	.88
Wisconsin.....	2,553,983	1913—1917—5 years.....	2,261	.91	1,774	.24	14.1
Wyoming.....	190,380	do.....	46	.26	1,202	1.06	11.9

Cases of influenza and pneumonia (all forms) reported, 1918.

	Jan.	Feb.	Mar.	Apr.	May.	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Total.
Alabama:													
Influenza.	81	54	27	37	29	7	2	1	15	122,127	13,507	8,148	141,035
Pneumonia (all forms).	1,007	565	1,061	269	340	158	291	108	415	8,229	775	452	13,670
Arkansas:													
Influenza.													
California:													
Influenza.													
Pneumonia (all forms).	440	306	370	331	319	190	185	140	132	1,644	301	273	4,631
Colorado:													
Influenza.													
Pneumonia (all forms).	18	1	2	3	1				2	1,183	548	201	1,956
Connecticut:													
Influenza.													
Pneumonia (all forms).	7	28	75	57	30	8	17	11	8,970	82,057	6,773	8,041	2,256
Florida:													
Influenza.													
Pneumonia (all forms).	47	77	36	27	75	22	14	21	16	9,295	1,722	598	982
Illinois:													
Influenza.													
Pneumonia (all forms).	602	580	1,242	1,409	612	163	142	68	541	145,067	40,906	43,304	20,298
Indiana:													
Influenza.													
Pneumonia (all forms).													
Iowa:													
Influenza.													
Kansas:													
Influenza.													
Pneumonia (all forms).	249	310	306	212	117	23	10	8	277	45,971	40,875	46,354	2,922
Louisiana:													
Influenza.													
Pneumonia (all forms).	487	454	334	292	233	347	120	126	108	137,156	21,384	25,816	3,742
Maine:													
Influenza.													
Pneumonia (all forms).													
Maryland:													
Influenza.													
Pneumonia (all forms).	69	80	83	38	18	7	10	5	1,702	47,928	16,534	13,780	80,554
Massachusetts:													
Influenza.													
Pneumonia (lobar).	719	925	1,252	860	462	165	135	65	5,377	7,311	11,688	325	10,939
Michigan:													
Influenza.													
Pneumonia (all forms).	223	187	267	445	168	83	55	42	78	43,540	26,862	45,693	4,835

Cases of influenza and pneumonia (all forms) reported, 1918—Continued.

Texas:	Influenza.....	1,153	748	1,049	46	19	6	7	445	45,943	9,094	13,412	71,849	Texas: Influenza (all forms).	
Pneumonia (all forms).....	0	325	286	261	46	19	6	7	42	1,781	4,461	1,471	4,461	Utah:	
Utah:	Influenza.....	135	115	109	92	35	18	15	31	16,632	10,615	9,842	1,811	1,811	Influenza (all forms).
Pneumonia (all forms).....	228	135	115	109	92	35	18	15	320	452	291	1,811	1,811	Pneumonia (all forms).	
Vermont:	Influenza.....	14	12	9	11	0	1	0	5,451	25,724	1,437	3,342	1,773	1,773	Vermont: Influenza (all forms).
Pneumonia (all forms).....	16	14	12	9	11	0	1	0	374	1,292	17	27	27	27	Pneumonia (all forms).
Virginia:	Influenza.....	512	512	512	512	512	512	512	512	139,439	46,563	54,302	54,302	54,302	Virginia: Influenza (all forms).
Washington:	Influenza.....	126	126	126	126	126	126	126	126	824	824	847	847	847	Washington: Influenza (all forms).
Pneumonia (all forms).....	54	65	65	65	65	65	65	65	78	134	20,707	19,075	15,575	15,575	Influenza (all forms).
West Virginia:	Influenza.....	138	138	138	138	138	138	138	138	16,323	16,323	4,849	3,528	3,528	West Virginia: Influenza (all forms).
Wisconsin:	Influenza.....	6	6	6	6	6	6	6	6	15,122	15,122	9,886	15,312	15,312	Wisconsin: Influenza (all forms).

148,366 cases reported during October and November.

Influenza, pneumonia (all forms), and both diseases combined—Deaths registered, 1918.

	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Total.		
Alabama:															
Influenza.....	81	54	27	37	29	7	2	1	15	2,478	1,446	1,269	5,446	Alabama: Influenza, Pneumonia (all forms).	
Pneumonia (all forms).....	730	731	462	567	275	111	62	75	110	1,225	775	836	5,689		
Combined.....	811	786	489	684	304	118	64	76	125	3,703	2,221	1,906	11,135	Combined.	
Arizona:															
Influenza.....	2	1	2	7	3	1	1	1	230	246	84	577	577	Arizona: Influenza, Pneumonia (all forms).	
Pneumonia (all forms).....	81	76	84	126	66	68	23	26	19	618	762	292	2,226		
Combined.....	83	77	86	133	69	63	24	26	20	848	1,008	376	2,803	Combined.	
Arkansas:															
Influenza.....	28	24	33	15	12	1	1	1	20	1,066	814	1,015	3,029	Arkansas: Influenza, Pneumonia (all forms).	
Pneumonia (all forms).....	294	286	269	132	130	53	55	50	71	842	207	330	2,389		
Combined.....	322	310	242	147	142	54	56	50	91	1,708	1,021	1,345	5,428	Combined.	
California:															
Influenza.....	32	21	20	26	20	11	7	2	8	4,420	5,702	2,959	13,288	California: Influenza, Pneumonia (all forms).	
Pneumonia (all forms).....	457	415	492	441	304	215	135	171	195	961	803	659	5,268		
Combined.....	489	436	512	467	324	226	162	173	203	6,381	6,505	3,688	18,556	Combined.	
Connecticut:															
Influenza.....	33	47	42	49	15	5	2	3	300	5,158	1,022	742	7,418	Connecticut: Influenza, Pneumonia (all forms).	
Pneumonia (all forms).....	332	310	429	356	165	72	69	69	148	605	275	380	5,294		
Combined.....	365	357	471	405	180	78	64	72	448	5,823	1,267	1,002	10,622	Combined.	
Delaware:															
Influenza.....	82	86	80	50	26	8	14	18	75	359	18	19	426	Delaware: Influenza, Pneumonia (all forms).	
Pneumonia (all forms).....	Combined.....	82	86	80	50	26	8	14	18	75	1,223	97	110	1,869	Combined.
District of Columbia:															
Influenza.....	21	11	15	10	3	1	1	1	37	1,471	131	327	2,028	District of Columbia: Influenza, Pneumonia (all forms).	
Pneumonia (all forms).....	149	128	180	104	61	26	24	18	51	338	68	149	1,295		
Combined.....	170	139	195	114	64	28	25	19	88	1,800	109	476	3,328	Combined.	

Florida:		Influenza.											
Influenza.		Pneumonia (all forms).										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	
Hawaii:													
Influenza.		Influenza.										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	
Illinois:													
Influenza.		Influenza.										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	
Indiana:													
Influenza.		Influenza.										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	
Iowa:													
Influenza.		Influenza.										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	
Kansas:													
Influenza.		Influenza.										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	
Kentucky:													
Influenza.		Influenza.										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	
Louisiana:													
Influenza.		Influenza.										Influenza.	
Pneumonia (all forms).		Pneumonia (all forms).										Pneumonia (all forms).	
Combined.												Combined.	

Influenza, pneumonia (all forms), and both diseases combined—Deaths registered, 1918—Continued.

	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	
Maine:														Maine: Influenza, Pneumonia (all forms).
Influenza.....	10	17	27	54	24	8	5	105	1,315	451	850	2,546		
Pneumonia (all forms).....	141	133	201	233	59	33	18	90	626	150	270	1,972		
Combined.....													4,518	Combined.
Maryland:														Maryland: Influenza, Pneumonia (all forms).
Influenza.....	81	18	40	22	9	3	4	2	85	4,154	465	405	5,244	
Pneumonia (lobar).....	412	436	407	316	174	83	66	38	122	6,535	335	384	6,211	
Combined.....													11,655	Combined.
Massachusetts:														Massachusetts: Influenza, Pneumonia (lobar).
Influenza.....	437	475	490	585	243	113	106	66	2,268	8,274	1,022	1,804	9,354	
Pneumonia (lobar).....														
Combined.....														Combined.
Michigan:														Michigan: Influenza.
Influenza.....	51	56	39	125	54	11	13	5	22	2,451	1,866	2,104	6,797	
Minnesota:														Minnesota: Influenza, Pneumonia (all forms).
Influenza.....	200	172	210	426	193	98	67	30	48	2,084	3,166	1,884	1,965	
Pneumonia (all forms).....									72	163	163	130	1,965	
Combined.....														Combined.
Nebraska:														Nebraska: Influenza, Pneumonia (all forms).
Influenza.....	132	147	165	188	108	87	27	27	4	279	266	465	587	
Pneumonia (all forms).....									39	1,336	931	1,012	4,169	
Combined.....														Combined.
New Jersey:														New Jersey: Influenza, Pneumonia (all forms).
Influenza.....	132	147	165	188	108	57	27	27	48	1,616	1,197	1,477	4,050	
Pneumonia (all forms).....														
Combined.....														Combined.
New York:														New York: Influenza, Pneumonia (all forms).
Influenza.....	148	139	268	216	81	27	20	10	273	17,990	5,800	2,499	26,676	
Pneumonia (all forms).....	2,326	2,129	3,308	2,344	1,416	713	601	487	913	18,864	4,105	2,653	38,080	
Combined.....														Combined.
	2,474	2,268	3,576	3,060	1,497	740	621	497	1,185	31,246	9,411	5,181	61,726	

North Dakota:		North Dakota:			
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
53	37	55	64	35	32
Combined.....	14	23
				1	1
				51	199
				75	559
				788	326
				559	2,345
				788	2,681
				559	Combined.
Ohio:		Ohio:		Ohio:	
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
1,114	94	93	196	67	13
1,053	864	1,002	1,322	514	208
Combined.....	14	14
				14	14
				30	5,778
				284	4,691
				3,441	3,877
				2,175	14,988
				1,823	13,042
				559	Combined.
Oklahoma:		Oklahoma:		Oklahoma:	
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
25	10	16	14	9	3
190	200	259	264	132	20
Combined.....	14	14
				29	587
				8	1,943
				30	90
				313	4,047
				2,446	2,446
				570	Combined.
Oregon:		Oregon:		Oregon:	
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
3	4	5	5	8	5
55	48	84	82	65	47
Combined.....	14	14
				18	18
				25	384
				285	539
				279	390
				190	1,315
				190	Combined.
Pennsylvania:		Pennsylvania:		Pennsylvania:	
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
182	188	195	364	147	35
2,334	2,096	2,470	3,170	1,331	637
Combined.....	20	14
				658	28,505
				404	13,967
				846	6,778
				843	50,951
				3,334	28,056
				3,334	Combined.
Porto Rico:		Porto Rico:		Porto Rico:	
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
8	7	3	2	2	1
187	152	171	131	141	79
Combined.....	10	1
				211	67
				202	1,841
				100	3,516
				147	5,474
				661	822
				661	3,004
				214	Combined.
Rhode Island:		Rhode Island:		Rhode Island:	
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
8	13	16	24	1	2
120	97	142	162	71	38
Combined.....	20	20
				140	243
				113	147
				600	2,306
				101	1,639
				147	Combined.
South Carolina:		South Carolina:		South Carolina:	
Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)	Influenza	Pneumonia (all forms)
29	33	15	19	7	4
339	283	242	255	209	107
Combined.....	79	44
				155	354
				3,611	1,710
				1,615	976
				1,330	6,967
				8,377	Combined.

August 15, 1919.

1860

Influenza, pneumonia (all forms), and both diseases combined—Deaths registered, 1918—Continued.

	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	
South Dakota:														South Dakota:
Influenza.....	6	4	5	7	2	1			5	442	796	579	1,847	Influenza,
Pneumonia (all forms).....	35	40	48	55	36	26	9	8	22	77	101	56	544	Pneumonia (all forms).
Combined.....	41	44	54	62	38	27	9	8	27	519	897	665	2,391	Combined.
Texas:														Texas:
Influenza.....	195	192	154	106	24	27	14	4	21	4,690	3,477	2,950	11,854	Influenza,
Pneumonia (all forms).....	1,204	1,202	840	880	379	139	123	87	95	1,319	948	613	7,686	Pneumonia (all forms).
Combined.....	1,399	1,394	994	786	403	166	186	91	119	6,006	4,425	3,583	19,460	Combined.
Utah:														Utah:
Influenza.....	6	2	5	13	6	1	1	2	4	404	626	387	1,459	Influenza,
Pneumonia (all forms).....	80	63	48	86	69	30	12	13	19	194	198	74	798	Pneumonia (all forms).
Combined.....	86	55	53	99	76	81	13	15	23	588	788	461	2,255	Combined.
Vermont:														Vermont:
Influenza.....	9	9	15	15	7	3	2	2	99	1,224	158	185	1,728	Influenza,
Pneumonia (all forms).....	67	51	63	73	35	16	19	6	5	44	24	30	433	Pneumonia (all forms).
Combined.....	76	60	78	88	42	19	21	8	104	1,268	162	215	2,161	Combined.
Washington:														Washington:
Influenza.....	10	8	19	13	8	7	2	3	8	4,072	1,419	1,585	4,154	Influenza,
Pneumonia (all forms).....	126	120	172	205	113	90	54	44	78	517	1,153	157	1,636	Pneumonia (all forms).
Combined.....	136	128	191	218	121	97	56	47	86	1,389	1,671	1,742	5,732	Combined.
Wisconsin:														Wisconsin:
Influenza.....	24	26	26	48	17	13	8	1	45	2,091	890	1,210	4,339	Influenza,
Pneumonia (all forms).....	324	324	325	551	341	179	118	101	105	625	1,440	1,772	6,126	Pneumonia (all forms).
Combined.....	348	340	351	679	368	193	126	102	104	2,616	2,360	2,662	10,468	Combined.
Wyoming:														Wyoming:
Influenza.....	20	21	22	54	23	14	7	5	9	42	95	32	219	Influenza,
Pneumonia (all forms).....	29	21	22	64	22	14	7	6	9	203	264	227	519	Pneumonia (all forms).
Combined.....	29	21	22	64	22	14	7	6	9	203	264	227	519	Combined.

Influenza, pneumonia (all forms), and influenza and pneumonia combined—Deaths registered and indicated mortality rates, 1918.

State.	Estimated population July 1, 1918.	Influenza.		Pneumonia (all forms).		Influenza.		Pneumonia (all forms).		Combined..	
		Total deaths registered.	Deaths per 1,000 inhabitants.	Total deaths registered.	Deaths per 1,000 inhabitants.	State.	Estimated population July 1, 1918.	Total deaths registered.	Deaths per 1,000 inhabitants.	Total deaths registered.	Deaths per 1,000 inhabitants.
Alabama.....	2,395,270	5,446	2.3	5,080	2.4	11,135	4.6	Mississippi.....	2,001,466	9,677	1.4
Arizona.....	272,034	2.1	2,228	8.2	2,803	10.3	Missouri.....	3,445,488	7,377	2.1	
Arkansas.....	1,792,965	3,029	1.7	2,399	1.3	5,288	3.0	Nebraska.....	1,296,877	7,014	5.183
California.....	3,119,412	288	4.3	5,288	1.7	18,536	5.9	New York.....	10,646,989	26,876	3.2
Connecticut.....	1,286,268	7,415	5.8	3,204	2.5	5,922	8.3	North Dakota.....	791,337	2,355	2.681
Delaware.....	216,941	426	2.0	1,443	6.7	1,869	8.6	Ohio.....	5,273,814	14,988	2.5
District of Columbia.....	401,681	2,028	5.0	1,298	3.2	3,936	8.3	Oklahoma.....	2,377,029	4,047	2.7
Florida.....	938,877	3,113	3.3	1,675	1.8	4,788	5.1	Oregon.....	886,243	1,315	6,493
Hawaii.....	223,419	165	1.6	176	4.92	657	2.2	Pennsylvania.....	50,951	5.8	2.507
Illinois.....	6,317,734	17,879	2.8	14,445	2.3	32,324	5.0	Porto Rico.....	1,247,677	5,474	3.2
Indiana.....	2,834,167	6,219	2.2	5,609	2.0	11,828	4.1	Rhode Island.....	637,415	2,306	2.4
Iowa.....	2,224,771	6,280	2.8	3,818	1.7	10,078	4.5	South Carolina.....	1,660,834	1,710	8,377
Kansas.....	1,874,195	2,639	1.4	6,052	3.2	8,691	4.6	Texas.....	735,434	1,847	2.5
Kentucky.....	2,408,547	9,181	3.8	5,302	2.2	14,483	6.0	Tennessee.....	4,601,279	11,854	1.7
Louisiana.....	1,884,778	5,925	3.1	4,026	2.1	9,951	5.3	Utah.....	455,648	1,459	3.2
Maine.....	782,191	2,546	3.3	1,972	2.5	4,518	5.8	Vermont.....	366,192	1,723	4.7
Maryland.....	1,384,559	5,244	3.8	6,311	4.6	18,555	8.3	Washington.....	1,660,678	4,154	2.5
Massachusetts.....	3,832,790	11,100	2.9	1,9354	2.4	120,454	5.3	Wisconsin.....	2,353,983	4,339	2.4
Michigan.....	3,133,678	6,797	2.2	—	—	—	—	Wyoming.....	190,350	169	4.3
Minnesota.....	2,345,287	7,182	3.1	1,955	.8	9,137	4.0			.9	5.2

Includes lobar pneumonia only.

Cases of malaria reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	38	18	41	100	105	182	370	354	388	81	83	30	1,790	Alabama.
Arkansas.....	173	323	409	505	621	525	903	469	128	102	171	4,709	Arkansas.	
California.....	20	11	23	26	53	76	144	92	121	83	8	4	681	California.
Colorado.....	1	1	1	1	1	1	1	1	1	1	1	1	2	Colorado.
Connecticut.....	4	Connecticut.
Florida.....	89	44	43	50	44	74	98	161	236	71	50	21	981	Florida.
Illinois.....	1	1	1	1	1	2	2	1	1	8	9	1	7	Illinois.
Kansas.....	19	30	103	152	359	7	348	26	362	105	41	16	28	Kansas.
Louisiana.....	3	2	2	0	29	29	22	22	35	8	6	10	2,125	Louisiana.
Maryland.....	213	Maryland.
Massachusetts.....	1	1	2	5	4	5	10	31	9	4	2	3	82	Massachusetts.
Minnesota.....	0	0	0	3	6	6	1	1	2	0	0	0	7	Minnesota.
Mississippi.....	3,492	4,659	6,094	8,707	11,477	14,103	16,214	15,712	12,172	5,996	4,265	2,742	105,783	Mississippi.
New Jersey.....	8	1	18	15	35	36	32	32	34	15	6	47	21	New Jersey.
Ohio.....	3	16	6	6	6	6	6	9	4	16	7	84	Ohio.
Pennsylvania.....	1	1	1	1	1	1	1	1	1	1	1	5	Pennsylvania.
Porto Rico.....	148	233	204	101	77	75	77	63	93	74	128	69	1,471	Porto Rico.
South Carolina.....	2	35	3	64	113	53	197	50	50	16	4	6	1,691	South Carolina.
Texas.....	0	2	19	23	34	53	55	52	58	45	31	9	562	Texas.
Virginia.....	193	262	270	335	367	577	560	701	415	360	184	99	4,821	Virginia.

1863

August 15, 1919.

Deaths registered as due to malaria, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	11	9	14	18	19	55	51	88	63	29	8	16	376	Alabama.
Arizona.....	24	20	27	15	24	44	52	68	46	45	19	36	404	Arizona.
Arkansas.....	3	1	3	2	6	3	8	3	8	12	3	2	34	Arkansas.
California.....	1	1	1	1	3	Connecticut.
Connecticut.....
Florida.....	6	4	11	18	17	32	25	29	26	33	9	8	217	Florida.
Illinois.....	3	3	12	5	7	37	12	18	10	6	2	2	77	Illinois.
Indiana.....	1	1	1	1	1	4	7	15	3	2	2	4	31	Indiana.
Iowa.....	0	0	0	1	1	0	0	0	0	0	0	0	2	Iowa.
Kansas.....	1	0	0	3	5	1	3	2	4	6	2	1	28	Kansas.
Kentucky.....	4	7	5	9	9	11	11	15	15	10	8	5	109	Kentucky.
Louisiana.....	7	16	16	42	64	89	83	73	71	23	15	11	610	Louisiana.
Maine.....	0	0	0	0	0	0	0	0	0	1	1	0	2	Maine.
Maryland.....	0	0	1	1	1	0	1	0	0	0	1	0	1	Maryland.
Massachusetts.....	1	0	0	0	0	0	0	0	0	0	0	0	4	Massachusetts.
Minnesota.....	0	0	0	0	0	0	0	0	0	0	0	0	1	Minnesota.
Nebraska.....	3	2	1	1	1	1	1	1	1	1	Nebraska.
New Jersey.....	2	1	3	1	5	2	2	3	2	1	2	1	12	New Jersey.
New York.....	3	1	3	2	1	4	22	26	26	26	3	2	24	New York.
Ohio.....	3	1	11	7	18	22	26	26	26	10	9	3	20	Ohio.
Oklahoma.....	1	1	1	1	1	1	1	1	1	168	Oklahoma.
Pennsylvania.....	1	107	121	104	94	61	120	129	113	150	213	171	1,634	Pennsylvania.
Porto Rico.....	151	107	121	104	94	61	120	129	113	150	213	171	1	Porto Rico.
Rhode Island.....	5	4	6	6	16	14	38	20	45	50	53	13	28	Rhode Island.
South Carolina.....	18	20	32	37	38	52	80	50	53	30	15	9	435	South Carolina.
Texas.....	1	1	Texas.
Vermont.....	1	1	Vermont.

August 15, 1919.

1864

Malaria—Cases reported, deaths registered, and indicated morbidity, mortality, and fatality rates, 1918.

State.	Estimated population, July 1, 1918.	Total cases reported.	Cases per 1,000 inhabitants.	Total Deaths registered.	Deaths per 1,000 inhabitants.	Fatalities per 100 cases.	State.	Estimated population, July 1, 1918.	Total cases reported.	Total Deaths registered.	Cases per 1,000 inhabitants.	Estimated population, July 1, 1918.	Total cases reported.	Total Deaths registered.	Fatalities per 100 cases.
Alabama.....	2,395,270	1,790	0.747	376	0.157	31.0	Massachusetts.....	3,822,730	32	0.021	4	0.001	4.9	14.3	
Arizona.....	2,272,034	4,709	1	1	.001		Minnesota.....	2,315,287	7	.003	1		305	.7	
Arkansas.....	1,792,965	651	2.626	404	.225	8.6	Mississippi.....	2,001,666	105,733	52,853	791		365	.64	
California.....	3,119,412	651	2.12.	54	.017	8.2	Missouri.....	3,418,198			222		222	.004	
Colorado.....	1,014,581	2	.002				Nebraska.....	1,286,877			1		1		
Connecticut.....	1,286,288	4	.003	3	.002	75.0	New Jersey.....	3,050,371	281	.091	12	.004	4.3		
Florida.....	938,877	931	.092	217	.231	23.3	New York.....	10,646,989			24		24		
Illinois.....	6,317,734	7	.013	77	.013		Ohio.....	5,273,314	34	.016	20	.004	23.8		
Indiana.....	2,854,167			31	.01		Oklahoma.....	2,377,329			163		163		
Iowa.....	2,224,771			2	.001		Pennsylvania.....	8,795,907			12		12		
Kansas.....	1,874,195	28	.015	28	.015		Porto Rico.....	1,247,677	6		1,471		1,471		
Kentucky.....	2,408,547			109	.045		Rhode Island.....	637,415			1		1		
Louisiana.....	1,884,775	2	1.127	510	.271	24.0	South Carolina.....	1,660,924	591	.356	289	.174	48.9		
Maine.....	1,752,101	213	.154	2	.003		Texas.....	4,601,270	562	.122	455	.065	77.4		
Maryland.....	1,384,539			6	.004		Vermont.....	2,380,182			1		1		
							Virginia.....	2,231,030	4,321	1.934					

Measles—Monthly averages of cases reported during the years for which data are available, 1913-1917.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Aver. age per year.	State.	
Alabama.....	1915-1917—3 years.....	265	1,318	2,042	1,542	767	229	85	62	27	58	143	357	6,915	Alabama.	
Arkansas.....	1917—1 year.....	891	1,517	1,765	1,143	644	116	33	5	67	542	1,260	1,260	7,945	Arkansas.	
California.....	1915-1916, 1917—3 years.....	804	1,552	1,668	1,523	1,170	614	338	227	262	101	491	381	6,319	California.	
Colorado.....	1914, 1916, 1917—3 years.....	385	983	1,630	1,622	679	883	494	46	46	41	47	145	6,313	Colorado.	
Connecticut.....	1913-1917—5 years.....	631	692	429	564	1,441	925	379	79	37	113	113	267	7,967	Connecticut.	
District of Columbia.....	1913-1917—5 years.....	214	75	60	48	53	348	153	31	11	18	55	75	3,010	District of Columbia.	
Hawaii.....	1913-1917—5 years.....	3,564	4,574	6,600	5,133	2,826	856	105	16	3	12	82	81	642	Hawaii.	
Illinois.....	1913, 1914, 1917—3 years.....	1,481	2,268	3,459	4,073	2,268	1,989	617	83	46	148	422	690	32,159	Illinois.	
Indiana.....	1914-1917—4 years.....	1,044	1,188	1,188	2,632	2,965	2,938	760	178	41	32	152	257	18,211	Indiana.	
Louisiana.....	1914-1917—4 years.....	1,044	501	297	179	139	75	48	42	25	145	538	779	11,002	Louisiana.	
Maine.....	1916-1917—2 years.....	165	194	194	193	193	118	19	184	48	43	58	241	519	2,082	Maine.
Maryland.....	1915-1917—3 years.....	638	844	1,407	1,661	1,300	1,385	466	140	96	116	270	663	9,356	Maryland.	
Massachusetts.....	1913, 1916, 1917—3 years.....	2,820	2,848	3,819	4,254	3,625	1,532	404	183	183	391	707	1,461	26,822	Massachusetts.	
Michigan.....	1913-1917—5 years.....	6,696	8,924	1,289	1,504	1,861	1,252	522	100	100	276	344	517	9,076	Michigan.	
Minnesota.....	1913-1917—6 years.....	323	1,193	1,193	1,193	1,193	1,193	672	109	80	60	26	60	1,170	6,864	Minnesota.
Mississippi.....	1914-1917—4 years.....	1,043	2,610	3,847	2,786	1,901	857	377	188	182	368	1,465	2,614	18,008	Mississippi.	
Montana.....	1913, 1914, 1916, 1917—4 years.....	443	391	620	488	632	203	68	24	17	63	49	121	3,026	Montana.	
Nevada.....	1913-1917—2 years.....	80	205	333	128	75	34	28	2	2	4	20	29	29	Nevada.	
New York.....	1913-1917—5 years.....	4,675	5,813	8,386	9,195	11,210	8,945	3,837	1,077	521	984	1,861	3,245	60,697	New York.	
North Dakota.....	1917—1 year.....	1,131	1,187	2,511	1,921	2,244	1,825	2,27	2	22	11	11	11	1,123	North Dakota.	
Ohio.....	1914-1917—4 years.....	2,352	3,688	5,445	6,284	5,185	2,977	1,045	227	138	294	693	1,061	2,779	Ohio.	
Oklahoma.....	1913, 1915, 1917—3 years.....	233	902	1,370	1,515	1,650	1,179	69	10	15	45	45	149	6,039	Oklahoma.	
Oregon.....	1913-1917—5 years.....	309	441	327	327	229	129	61	34	23	41	74	168	2,283	Oregon.	
Pennsylvania.....	1916, 1917—2 years.....	7,223	9,100	13,357	12,716	9,635	6,041	2,186	628	813	619	1,319	2,277	65,420	Pennsylvania.	
Puerto Rico.....	1917—1 year.....	112	181	387	504	1,005	1,005	464	345	540	861	705	638	7,001	Puerto Rico.	
Rhode Island.....	1914-1916, 1917—3 years.....	122	92	198	225	222	149	76	12	5	21	65	39	1,223	Rhode Island.	
South Carolina.....	1918-1917—5 years.....	416	522	577	459	156	73	34	21	16	29	151	258	2,762	South Carolina.	
South Dakota.....	1912, 1914, 1915, 1917—3 years.....	192	170	174	269	269	94	23	10	0	15	33	33	1,227	South Dakota.	
Texas.....	1917—1 year.....	874	1,715	1,144	269	269	1	6	0	66	84	909	4,746	Texas.		
Utah.....	1918-1915, 1917—4 years.....	1,988	1,623	1,210	1,001	664	244	62	22	12	30	62	179	7,011	Utah.	
Vermont.....	1914-1917—4 years.....	2,267	443	721	506	559	346	247	77	62	60	64	73	3,442	Vermont.	
Virginia.....	1915-1917—3 years.....	1,668	2,311	3,197	3,199	1,486	494	494	175	134	204	548	548	17,749	Virginia.	
Washington.....	1912-1917—5 years.....	1,080	1,207	1,250	1,445	1,341	815	272	88	54	256	454	694	9,056	Washington.	
West Virginia.....	1912—1 year.....	1,197	921	1,455	1,624	1,624	1,620	459	29	8	17	54	21	3,922	West Virginia.	
Wisconsin.....	1912-1917—5 years.....	822	907	1,273	1,080	1,150	914	389	81	48	211	314	516	7,069	Wisconsin.	
Wyoming.....	1913-1917—5 years.....	41	71	101	101	61	61	16	4	6	20	20	20	5,830	Wyoming.	

August 15, 1919.

1866

Cases of measles reported, 1918.

State.	State.											Total.	State.		
	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.			
Alabama.....	2,327	1,451	1,194	417	149	81	41	66	229	93	51	111	6,220	Alabama.	
Arizona.....	5	38	73	27	6	8	2	2	2	1	0	0	6,162	Arizona.	
Arkansas.....	1,242	1,913	1,031	527	362	99	34	5	8	13	15	66	5,335	Arkansas.	
California.....	2,688	5,392	5,288	3,041	2,989	1,474	715	219	314	494	61	29	23,194	California.	
Colorado.....	491	482	498	289	181	99	16	5	6	3	3	29	2,080	Colorado.	
Connecticut.....	705	701	773	851	932	550	304	110	86	204	89	245	5,450	Connecticut.	
District of Columbia.....	576	1,084	2,038	2,023	2,023	214	79	22	16	12	10	17	1,001	District of Columbia.	
Florida.....	590	2,207	1,127	151	76	62	111	69	33	64	185	2,187	Florida.		
Illinois.....	1,150	829	1,227	1,278	1,501	641	273	121	56	141	67	171	7,575	Illinois.	
Indiana.....	820	1,370	1,211	633	634	304	62	60	62	44	113	5,983	Indiana.		
Iowa.....	440	218	250	118	63	30	15	6	2	4	8	1,198	Iowa.		
Kansas.....	2,170	2,391	1,504	2,119	1,615	481	74	77	20	66	46	90	10,798	Kansas.	
Louisiana.....	1,002	887	680	222	163	81	122	83	33	9	72	151	3,225	Louisiana.	
Maine.....	217	217	208	65	75	74	146	25	13	52	41	4	1,647	Maine.	
Maryland.....	953	1,240	2,631	3,943	3,316	1,287	554	183	95	91	163	174	14,660	Maryland.	
Massachusetts.....	2,950	3,471	4,018	5,210	6,335	3,653	1,802	500	263	318	179	343	28,212	Massachusetts.	
Michigan.....	675	558	1,018	1,263	1,047	455	127	60	243	106	63	63	5,906	Michigan.	
Minnesota.....	458	549	615	415	532	707	388	59	18	19	21	26	3,727	Minnesota.	
Mississippi.....	15,452	10,985	7,743	4,011	1,884	757	277	161	191	17	123	166	286	42,046	Mississippi.
Montana.....	134	190	195	218	287	150	124	14	24	2	16	2	1,288	Montana.	
Nebraska.....	280	283	214	480	155	43	20	11	69	16	17	17	1,584	Nebraska.	
New Jersey.....	2,457	3,925	5,432	6,502	5,950	2,786	1,121	103	84	27	48	26	4,427	New Jersey.	
New York.....	7,749	8,470	13,446	14,842	14,310	7,829	4,804	1,832	446	588	414	345	74,385	New York.	
North Dakota.....	1,449	1,700	1,887	1,268	946	384	111	77	100	127	34	38	8,090	North Dakota.	
Ohio.....	30	58	287	69	94	35	27	4	8	1	2	2	615	North Dakota.	
Oregon.....	1,172	1,754	2,288	495	2,750	1,445	728	237	24	322	168	452	14,065	Ohio.	
Pennsylvania.....	3,050	4,411	5,577	3,322	810	3,280	3,725	18	34	46	16	34	4,742	Pennsylvania.	
Puerto Rico.....	4,432	8,146	10,583	11,138	6,611	3,440	1,004	616	1,021	520	505	51,886	Puerto Rico.		
Rhode Island.....	311	658	834	85	70	62	301	5	10	1	1	2	1,746	Rhode Island.	
South Carolina.....	449	754	257	1,434	1,174	20	26	8	11	50	4	4	3,947	South Carolina.	
South Dakota.....	108	170	140	289	283	164	79	30	13	12	10	22	1,320	South Dakota.	
Texas.....	1,509	722	1,110	500	265	136	9	8	30	10	8	49	4,346	Texas.	
Utah.....	630	630	758	1,079	1,171	601	184	100	92	78	27	17	5,274	Utah.	
Vermont.....	42	33	37	112	112	2,450	2,288	114	34	57	68	83	1,038	Vermont.	
Virginia.....	2,565	3,808	4,685	4,750	2,801	807	391	288	96	102	38	219	20,546	Virginia.	
Washington.....	545	768	1,088	991	668	535	276	163	102	36	12	24	6,166	Washington.	
West Virginia.....	116	112	2,266	4,034	4,060	4,037	618	51	20	6	16	10	1,488	West Virginia.	
Wisconsin.....	1,112	2,266	4,034	4,060	4,037	700	181	137	433	63	59	59	19,376	Wisconsin.	
Wyoming.....	68	124	326	150	218	117	11	1	1	1	1	4	1,033	Wyoming.	

Deaths registered as due to measles, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	80	118	76	40	9	8	4	4	8	6	4	3	470	Alabama.
Arizona.....	1	6	8	4	5	5	6	6	1	7	14	16	37	Arizona.
Arkansas.....	124	65	46	27	25	13	6	1	1	7	3	2	146	Arkansas.
California.....	17	24	26	29	22	23	4	4	3	4	7	5	164	California.
Connecticut.....	13	7	12	9	9	6	4	3	4	6	7	5	85	Connecticut.
Delaware.....	2	1	2	0	0	3	1	1	1	1	0	0	14	Delaware.
District of Columbia.....	5	11	7	12	10	10	7	7	7	1	0	0	48	District of Columbia.
Hawaii.....	10	26	35	30	17	17	1	1	1	1	1	1	3	Hawaii.
Illinois.....	37	28	43	61	76	13	7	5	5	5	13	15	341	Illinois.
Iowa.....	7	13	29	52	51	26	13	15	15	1	1	2	120	Iowa.
Indiana.....	10	12	20	67	51	20	13	17	17	1	1	3	214	Indiana.
Kansas.....	20	44	64	67	51	20	13	17	17	1	1	4	5	Kansas.
Kentucky.....	26	46	66	68	88	88	17	17	17	3	3	5	313	Kentucky.
Louisiana.....	47	46	35	38	37	28	13	13	13	4	4	1	341	Louisiana.
Maine.....	20	17	19	10	11	12	7	4	4	2	1	0	254	Maine.
Maryland.....	10	11	34	29	41	11	7	3	3	1	3	1	168	Maryland.
Massachusetts.....	39	48	59	79	100	80	52	21	15	21	15	4	519	Massachusetts.
Michigan.....	17	33	31	38	69	30	19	6	2	7	10	4	259	Michigan.
Minnesota.....	2	3	13	18	18	14	14	14	14	1	2	5	89	Minnesota.
Nebraska.....	1	1	1	2	6	1	1	1	1	0	0	0	0	Nebraska.
New Jersey.....	69	50	97	112	84	61	31	8	5	4	5	2	523	New Jersey.
New York.....	123	150	251	262	191	129	73	37	21	29	14	13	1,283	New York.
North Dakota.....	1	1	3	3	3	3	3	3	3	1	1	1	7	North Dakota.
Ohio.....	11	23	35	68	44	37	15	10	3	9	7	8	270	Ohio.
Oklahoma.....	16	63	64	59	39	11	6	3	1	3	0	0	254	Oklahoma.
Oregon.....	3	1	10	17	19	19	9	4	2	0	0	0	57	Oregon.
Pennsylvania.....	48	56	101	142	160	65	59	53	21	37	43	17	802	Pennsylvania.
Porto Rico.....	63	64	70	45	65	32	41	18	14	10	3	3	425	Porto Rico.
Rhode Island.....	2	13	14	20	24	20	13	9	5	2	5	1	128	Rhode Island.
South Carolina.....	14	16	15	9	8	1	2	1	1	3	3	3	81	South Carolina.
South Dakota.....	2	2	5	2	9	4	2	1	1	3	3	3	33	South Dakota.
Texas.....	123	268	287	176	45	47	19	6	1	13	3	2	1,009	Texas.
Utah.....	2	4	7	10	2	1	1	1	1	3	2	2	34	Utah.
Vermont.....	2	2	5	18	15	12	1	1	1	3	3	3	15	Vermont.
Washington.....	6	15	18	13	20	16	7	2	1	2	3	3	102	Washington.
Wisconsin.....	1	1	5	1	1	1	1	1	1	2	1	2	100	Wisconsin.
Wyoming.....													6	Wyoming.

August 15, 1919.

1868

Measles—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated mortality, morbidity, and fatality rates, 1918.

State.	Estimated population July 1, 1918.	Average.		1918				State.
		Years.	Cases reported per annum.	Cases reported per 1,000 inhabitants.	Total cases reported.	Cases per 1,000 inhabitants.	Total deaths registered.	
Alabama.....	2,395,270	1915-1917-3 years.	6,915	3.0	6,220	2.6	470	0.196
Arizona.....	272,634	1911-1 year	7,445	4.4	5,335	3.0	37	.136
Arkansas.....	1,762,865	1913-1917-5 years	9,907	3.5	22,194	7.4	386	.187
California.....	3,119,412	1914, 1916, 1917-3 years	6,319	6.6	2,080	2.1	164	.049
Colorado.....	1,014,581	1913-1917-5 years	6,518	5.3	5,450	4.2	85	...
Connecticut.....	1,286,268	1913-1917-5 years	3,010	8.4	7,001	17.4	14	.066
Delaware.....	216,941	1913-1917-5 years	3,010	8.4	2,197	2.3	46	...
District of Columbia.....	401,681	1913-1917-5 years	643	3.0	7,575	1.2	137	119
Florida.....	938,677	1915-1917-5 years	32,190	5.3	6,583	2.0	111	.146
Hawaii.....	223,419	1913, 1914, 1917-3 years	18,211	6.6	1,198	5.5	341	...
Illinois.....	6,317,734	1914-1917-4 years	11,002	6.1	10,783	5.8	120	.042
Indiana.....	2,854,107	1914-1917-4 years	11,002	6.1	10,783	5.8	214	2.1
Iowa.....	2,224,771	1915-1917-5 years	3,866	2.7	3,925	1.9	313	.096
Kansas.....	1,874,195	1915-1917-5 years	2,082	2.7	1,647	2.1	341	17.9
Kentucky.....	2,406,547	1914-1917-2 years	2,082	2.7	1,647	2.1	103	1.42
Louisiana.....	1,684,778	1916-1917-3 years	9,383	6.9	14,690	10.6	132	7.2
Maine.....	782,191	1913, 1916, 1917-3 years	26,822	7.0	29,212	7.6	519	1.35
Maryland.....	1,384,539	1913, 1916, 1917-3 years	9,076	5.906	5,906	1.9	112	1.1
Massachusetts.....	3,832,750	1913-1917-5 years	5,864	3.0	5,727	1.6	259	1.8
Michigan.....	3,133,678	1913-1917-5 years	18,008	9.3	42,045	21.0	89	0.83
Minnesota.....	2,345,287	1914-1917-4 years	1,025	1.2	1,025	1.6	732	4.4
Mississippi.....	2,001,466	1914-1917-4 years	1,025	1.2	1,025	1.6	89	2.4
Missouri.....	3,443,498	1913, 1914, 1916, 1917-4 years	1,025	1.2	1,025	1.6	732	1.7
Montana.....	480,376	1913, 1914, 1916, 1917-4 years	6,8	1,288	6,8	2.6	60	.017
Nebraska.....	1,296,977	1913, 1917-2 years	949	9.2	1,584	1.2	13	.010
Nevada.....	114,742	1913, 1917-2 years	949	9.2	1,584	1.2	13	.010
New Jersey.....	3,084,371	1913-1917-5 years	60,697	6.0	26,487	6.9	523	.110
New York.....	10,694,869	1913-1917-5 years	60,697	6.0	74,365	7.0	1,293	1.8
North Carolina.....	2,468,025	1917-1 year	1,135	1.5	8,060	8.3	8	1.7
North Dakota.....	791,437	1917-1 year	20,379	5.7	14,065	2.7	7	...
Ohio.....	5,273,814	1914-1917-4 years	5,009	2.4	2,700	2.7	270	.051
Oklahoma.....	2,377,629	1913, 1915, 1917-3 years	1,258	4.72	2,289	5.3	254	1.9
Oregon.....	588,243	1913-1917-5 years	65,450	2.8	51,886	5.9	57	.064
Pennsylvania.....	8,785,067	1916, 1917-2 years	1,247	5.7	802	5.9	1,247	1.5
Porto Rico.....	1,247,677	1917-1 year	1,247	5.7	425	1.4	425	24.3

Rhode Island.....	637,415	1914-1916, 1917-3 years.....	1,223	2.0	3,647	5.7	128	.201	3.5
South Carolina.....	1,660,834	1913-1917-5 years.....	2,792	1.7	2,245	1.4	81	.049	3.6
South Dakota.....	735,434	1913-1914, 1917-3 years.....	1,877	2.0	1,820	1.8	33	.045	2.5
Texas.....	4,001,270	1917-—year.....	1,356	1.1	1,356	1.9	1,000	.219	23.2
Utah.....	455,648	1913-1915, 1917-4 years.....	16,6	5,274	11.6	34	15	.073	.6
Vermont.....	366,192	1914-1917-—years.....	2,422	9.5	1,086	3.0	15	.041	1.4
Virginia.....	2,234,080	1915-1917-3 years.....	11,749	8.1	20,846	9.3	102	.061	2.0
Washington.....	1,660,578	1913-1917-5 years.....	6,036	6.1	5,146	3.1	10
West Virginia.....	1,439,165	1917-—year.....	3,922	2.8	1,485	1.0	103	.059	.5
Wisconsin.....	2,553,982	1918-1917-5 years.....	7,689	3.1	19,376	7.4	6	.032	.6
Wyoming.....	180,38000.....	589	3.1	1,083	6.4	6

August 15, 1919.

1870

Poliomyelitis (infantile paralysis)—Monthly averages of cases reported during the years for which data are available, 1913-1917.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Average per year.	State.		
Alabama.....	1913-1915-1917—4 years.....	1	1	1	2	2	4	23	18	5	3	2	2	64	Alabama.		
Arizona.....	1914-1916—2 years.....	(1)	0	0	0	0	0	(1)	1	1	0	0	0	3	Arizona.		
Arkansas.....	1917—1 year.....	2	0	0	0	0	0	1	2	0	0	0	0	8	Arkansas.		
California.....	1913-1915—5 years.....	5	3	2	3	2	5	1	9	10	8	12	12	81	California.		
Colorado.....	1916-1917—5 years.....	0	1	1	2	2	0	(1)	1	1	3	4	2	0	Colorado.		
Connecticut.....	1913-1917—5 years.....	1	1	1	1	2	2	34	76	58	24	8	3	21	Connecticut.		
District of Columbia.....	1913-1917—5 years.....	(1)	0	0	0	0	0	(1)	0	0	(1)	0	1	13	District of Columbia.		
Hawaii.....	1913-1915-1917—4 years.....	0	0	1	0	0	0	0	0	0	(1)	0	0	3	Hawaii.		
Idaho.....	1913-1914-1917—3 years.....	10	12	11	13	8	9	0	3	1	1	1	0	10	Idaho.		
Illinois.....	1913-1917—5 years.....	3	3	3	2	1	2	12	19	19	20	7	2	38	Illinois.		
Indiana.....	1913-1917—5 years.....	3	3	3	2	1	1	2	12	19	20	7	2	38	Indiana.		
Iowa.....	1913-1914-1916-1917—4 years.....	2	2	1	1	1	1	2	11	31	44	15	8	121	Iowa.		
Kansas.....	1913-1917—5 years.....	1	1	1	1	1	1	1	5	11	11	13	7	4	68	Kansas.	
Louisiana.....	1914-1917—4 years.....	1	1	1	1	1	1	1	2	7	3	2	1	26	Louisiana.		
Maine.....	1916-17—2 years.....	0	0	1	0	1	0	(1)	8	8	28	22	16	3	80	Maine.	
Maryland.....	1915-1917—3 years.....	4	1	1	2	2	2	16	38	43	44	14	2	169	Maryland.		
Massachusetts.....	1913-1916-1917—3 years.....	8	13	6	4	6	6	9	51	117	254	267	76	20	821	Massachusetts.	
Michigan.....	1913-1917—5 years.....	4	4	3	3	3	2	3	4	14	44	50	33	9	173	Michigan.	
Minnesota.....	1914-1917—4 years.....	3	2	2	2	2	3	7	13	19	52	77	26	9	31	Minnesota.	
Montana.....	1913-1914-1916-1917—4 years.....	6	6	9	12	7	10	1	1	1	9	9	7	9	31	Montana.	
Nevada.....	1912-1915—2 years.....	0	0	0	0	0	0	(1)	2	4	9	9	7	3	35	Nevada.	
New Jersey.....	1915-1917—3 years.....	4	5	3	4	5	4	5	132	130	199	199	0	0	3	857	New Jersey.
New York.....	1913-1916-1917—3 years.....	14	19	9	12	12	12	33	85	1,168	491	171	44	16	2,886	New York.	
North Dakota.....	1910-11-12-13 years.....	10	0	0	0	0	0	0	0	0	7	5	11	10	2	30	North Dakota.
Oklahoma.....	1914-1917—4 years.....	8	8	8	5	7	7	7	11	11	47	94	43	18	8	367	Oklahoma.
Oregon.....	1913-1914-1916-1917—4 years.....	1	1	1	2	0	0	1	1	1	0	1	2	0	0	20	Oregon.
Pennsylvania.....	1916-1917—2 years.....	9	7	14	8	0	0	(1)	0	0	43	462	226	60	26	1,368	Pennsylvania.
Rhode Island.....	1911—1 year.....	0	0	0	0	0	0	11	88	1	3	2	1	0	9	35	Rhode Island.
South Carolina.....	1915-1917—4 years.....	0	0	2	0	0	1	1	1	1	15	15	7	2	21	South Carolina.	
South Dakota.....	1910-11-12-13 years.....	0	1	0	1	0	0	1	1	1	1	1	4	6	4	2	South Dakota.
Texas.....	1914-1917—4 years.....	10	10	8	8	5	7	7	11	11	47	94	43	18	8	367	Texas.
Vermont.....	1917—1 year.....	1	1	1	2	0	0	1	1	1	0	1	2	0	0	4	Vermont.
Virginia.....	1913-1915-1917—4 years.....	1	1	1	1	1	1	1	1	1	1	1	1	0	0	3	Virginia.
West Virginia.....	1916-1917—2 years.....	1	1	1	1	1	1	1	1	1	1	1	1	0	0	3	West Virginia.
Washington.....	1912-1913—4 years.....	15	15	12	17	17	17	15	35	45	40	30	10	6	33	Washington.	
Wisconsin.....	1913-1914—5 years.....	89	13	13	1	1	0	1	2	64	72	59	16	6	20	Wisconsin.	
Wyoming.....	1916, 1917—2 years.....	3	3	0	0	0	0	1	1	1	5	5	2	1	4	13	Wyoming.

¹ Less than one.

Cases of poliomyelitis (infantile paralysis) reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	2	0	1	3	2	0	1	0	1	2	0	2	2	15
Arkansas.....	0	1	1	4	7	6	5	0	5	5	0	0	0	6
California.....	10	2	1	4	7	6	5	10	5	5	9	8	2	75
Colorado.....	1	2	1	0	2	2	1	14	21	2	7	2	2	6
Connecticut.....	0	0	0	0	0	0	0	0	2	0	0	0	0	53
District of Columbia.....	10	4	17	21	18	16	43	73	79	18	2	2	2	303
Florida.....	16	2	14	1	3	5	15	11	16	17	1	1	1	68
Illinois.....	2	2	4	0	1	7	15	39	18	29	1	1	1	176
Indiana.....	1	4	4	1	2	1	3	4	5	19	2	2	2	30
Iowa.....	2	0	2	1	0	2	5	1	9	3	2	2	2	39
Kansas.....	0	0	0	1	1	0	0	1	2	1	0	0	0	5
Louisiana.....	2	0	0	0	1	4	12	15	16	20	12	3	1	90
Maine.....	0	0	0	1	3	6	6	6	6	10	20	7	3	99
Maryland.....	0	5	3	6	3	3	3	12	12	30	26	9	3	103
Massachusetts.....	5	2	1	4	3	3	2	8	11	38	13	3	1	82
Michigan.....	1	2	2	0	3	2	3	3	12	2	5	6	3	67
Minnesota.....	3	5	2	2	3	3	3	3	12	2	5	6	3	8
Mississippi.....	3	3	2	2	2	1	1	1	1	1	1	1	1	10
Montana.....	0	1	2	1	2	2	2	1	1	1	1	1	1	1
Nebraska.....	1	4	5	3	2	2	2	4	9	10	2	0	1	43
New Jersey.....	14	5	10	13	4	19	38	46	70	16	4	4	4	233
New York.....	1	1	3	3	3	2	3	3	3	1	3	2	2	22
North Carolina.....	1	1	1	1	1	1	1	7	9	3	1	1	1	132
North Dakota.....	7	6	8	11	6	16	31	51	6	6	6	3	1	4
Ohio.....	0	0	0	2	0	0	1	0	0	0	0	1	0	302
Oregon.....	0	7	11	7	5	13	40	83	100	25	5	3	3	12
Pennsylvania.....	3	0	1	0	0	0	0	0	4	7	0	0	0	4
Rhode Island.....	0	1	1	0	1	2	1	1	1	1	1	1	1	1
South Carolina.....	1	1	1	1	1	1	1	1	1	1	1	1	1	1
South Dakota.....	0	2	1	2	4	9	12	6	3	1	5	1	1	46
Texas.....	0	14	1	0	0	3	5	6	41	47	8	2	2	288
Utah.....	1	4	20	4	1	1	2	12	12	12	4	5	5	82
Vermont.....	1	3	4	4	4	2	1	8	2	2	1	1	2	24
Virginia.....	2	1	0	0	3	5	6	3	7	1	2	0	0	30
Washington.....	14	50	28	11	8	1	1	1	1	1	8	2	2	2
West Virginia.....	4	14	1	0	0	3	5	6	3	7	1	2	0	288
Wisconsin.....	2	1	1	0	0	3	5	6	41	47	8	2	2	2
Wyoming.....	1	14	50	28	11	1	1	1	1	1	8	2	2	2

Deaths registered as due to poliomyelitis (infantile paralysis), 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.	
Alabama.....	2	0	1	2	3	2	0	1	0	1	0	0	2	12	Alabama.
Arkansas.....	3	1	1	2	1	2	0	3	0	4	1	1	1	14	Arkansas.
California.....	5	1	1	2	1	2	1	4	1	2	1	2	3	18	California.
Connecticut.....														8	Connecticut.
Florida.....														7	Florida.
Illinois.....	13	8	9	9	9	16	13	21	24	10	7	1	1	144	Illinois.
Indiana.....	2	2	1	3	3	5	2	5	5	4	7	5	5	34	Indiana.
Iowa.....	0	1	3	0	0	0	2	13	8	6	5	3	0	42	Iowa.
Kansas.....	2	0	0	0	0	0	1	3	1	1	1	1	1	11	Kansas.
Kentucky.....	2	2	1	3	0	1	1	16	10	4	1	0	0	40	Kentucky.
Louisiana.....	2	1	0	1	1	1	2	0	1	3	1	0	0	12	Louisiana.
Maine.....														7	Maine.
Maryland.....	0	0	1	0	0	3	2	0	3	3	4	2	0	16	Maryland.
Massachusetts.....	3	1	3	2	2	3	1	2	2	12	6	1	2	33	Massachusetts.
Michigan.....	1	2	0	0	1	2	3	4	7	10	10	4	4	45	Michigan.
Minnesota.....	1	1	2	2	1	2	1	0	1	5	6	3	1	26	Minnesota.
Nebraska.....														8	Nebraska.
New Jersey.....														12	New Jersey.
New York.....	0	6	5	5	5	12	8	4	11	9	1	2	2	53	New York.
North Dakota.....														9	North Dakota.
Ohio.....	1	2	1	2	0	3	2	2	3	12	14	3	2	55	Ohio.
Oklahoma.....	0	0	0	0	0	1	0	0	0	0	0	1	0	10	Oklahoma.
Oregon.....														1	Oregon.
Pennsylvania.....	3	5	5	5	2	4	0	0	0	0	0	0	0	90	Pennsylvania.
Rhode Island.....	1	1	2											1	Rhode Island.
South Carolina.....														6	South Carolina.
South Dakota.....														3	South Dakota.
Texas.....	2	1	5	4	0	1	6	1	6	1	5	1	2	4	Texas.
Utah.....														1	Utah.
Vermont.....														1	Vermont.
Washington.....	0	1	0	2	1	0	0	0	0	0	1	0	1	7	Washington.
Wisconsin.....		2	18	8	4	5	7	18	15	6	1	3	1	87	Wisconsin.
Wyoming.....														1	Wyoming.

Poliomyelitis (infantile paralysis)—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, registered, and indicated morbidity, mortality, and fatality rates, 1918.

State.	Estimated population July 1, 1918.	Average.		1918				State.	
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.	Total deaths registered.	Deaths per 1,000 inhabitants.	Fatalities per 100 cases.	
Alabama.....	2,395,270	1913-1915-1917-4 years.....	.64	0.028	15	.006	12	.005	80.0
Arizona.....	272,034	1914-1916-2 years.....	.3	.012	6	0	
Arkansas.....	1,782,965	1917-1 year.....	.8	.005	6	14	.008	
California.....	3,116,412	1913-1917-5 years.....	.81	.028	75	.024	18	.006	24.0
Colorado.....	1,014,531	1916,1917-2 years.....	.12	.012	6	.008	
Connecticut.....	1,286,268	1913-1917-5 years.....	.21	.0172	53	.041	8	.006	15.1
District of Columbia.....	401,681	do.....	.13	.036	2	.005	
Hawaii.....	938,877	1913-1915, 1917-4 years.....	3	.014	7	.007	
Idaho.....	223,419	1913-1915-1 year.....	.10	.026	0	
Illinois.....	461,766	1913-1914, 1917-3 years.....	.374	.062	308	.048	144	.023	47.6
Indiana.....	6,317,734	1913-1917-4 years.....	.93	.033	68	.024	34	.012	50.0
Iowa.....	2,354,167	1913-1917-5 years.....	.68	.034	176	.019	42	.019	23.9
Kansas.....	2,408,547	1913, 1914-1916, 1917-4 years.....	.68	.033	30	.016	11	.006	36.7
Kentucky.....	1,874,105	1913-1917-5 years.....	26	.014	40	.017	Kentucky.
Louisiana.....	1,584,778	1914-1917-4 years.....	.80	.013	5	7	.006	30.8
Maine.....	732,194	1916, 1917-2 years.....	.169	.0124	90	.065	16	.009	Maine.
Massachusetts.....	1,384,539	1915-1917-3 years.....	.821	.023	99	.028	38	.012	17.8
Michigan.....	3,832,790	1913-1916, 1917-5 years.....	173	.057	103	.033	55	.018	38.4
Minnesota.....	3,133,678	1913-1917-5 years.....	241	.107	82	.035	26	.011	53.4
Mississippi.....	2,345,287	do.....	131	.068	57	.028	9	.004	31.7
Missouri.....	2,001,456	1914-1917-4 years.....	29	.008	15.8
Montana.....	3,448,498	1913, 1914, 1916, 1917-4 years.....	35	.078	8	.016	Montana.
Nebraska.....	486,376	1913-1917-2 years.....	11	.008	10	.008	8	.006	80.0
Nevada.....	1,286,877	1913-1915-2 years.....	3	.030	Nevada.
New Jersey.....	3,080,371	1913-1917-5 years.....	847	.297	43	.014	12	.004	27.9
New York.....	10,646,939	do.....	2,896	.287	233	.022	58	.005	24.9
North Carolina.....	2,468,025	1917-1 year.....	22	.009	North Carolina.
North Dakota.....	191,437	1914-1917-4 years.....	36	.047	26	.083	8	North Dakota.
Ohio.....	5,273,814	1917-1 year.....	357	.070	152	.029	55	.010	36.2
Oklahoma.....	2,377,628	1913-1915-1917-4 years.....	20	.009	10	.004	Oklahoma.
Oregon.....	888,243	1913-1917-2 years.....	15	.018	4	.005	1	.001	25.0
Pennsylvania.....	8,798,067	1916,1917-2 years.....	1,368	.159	302	.034	90	.010	26.8

Poliomyelitis (infantile paralysis)—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated mortality, and fatality rates, 1918—Continued.

State.	Estimated population July 1, 1918.	Average.		1918				State.
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.	Total deaths registered.	Fatalities per 100 cases.	
Rhode Island.....	637,415	1917-1 year	0	0.014	12	0.019	6	0.009
South Dakota.....	1,686,934	1914-1917—4 years	40	.025	4	.033	9	0.006
South Dakota.....	733,324	1913, 1914-2 years	21	.021	24	.010	3	0.004
Texas	4,601,279	1915, 1917-2 years	9	.002	46	.002	35	0.008
Utah	4,452,648				1		1	0.002
Vermont	386,192	1911-1917—4 years	145	39	.010	1	0.03
Virginia.....	2,238,520	1913, 1915-1917—4 years	271	.015	82	.037	7	0.004
Washington.....	1,686,578	1913-1916—4 years	20	.014	24	.014	7	0.004
West Virginia.....	1,431,165	1917-1918—4 years	265	.015	35	.021	1	0.003
Wisconsin.....	2,555,933	1913-1917—5 years	382	.018	248	.010	87	0.004
Wyoming.....	190,380	1916, 1917-2 years	4	.022	2	.011	1	0.005

Rabies in man—Cases reported and deaths registered, 1918.

State.	Estimated population July 1, 1918.	Cases.	Deaths.	State.		Estimated population July 1, 1918.	Cases.	Deaths.
				State.	Cases.			
Alabama.....	2,395,270	3	3	Ohio.....	5,273,814	3	10	10
Illinois.....	6,317,754	1	1	Pennsylvania.....	8,798,967	3	3	3
Kentucky.....	2,408,547	1	1	Porto Rico.....	1,247,677	1	1	1
Louisiana.....	1,884,778	5	5	South Carolina.....	1,660,834	2	2	2
Missouri.....	3,448,498	1	1	Texas.....	4,601,279	6	6	6
Nebraska.....	1,296,877	1	1	Utah.....	4,452,648	1	1	1
New Jersey.....	3,080,371	2	2	Wisconsin.....	2,558,933	1	1	1

<i>Cases of rabies in animals reported, 1918.</i>								
State.	Cases.	State.	Cases.	State.	Cases.	State.	Cases.	State.
Arkansas.....	30	Maryland.....	37	Mississippi.....	108	South Carolina.....	108	
Connecticut.....	96	Massachusetts.....	26	Oklahoma.....	78	Virginia.....	78	
District of Columbia.....	20	Minnesota.....	7	Oregon.....	11	Wisconsin.....	11	
Kansas.....	9							27

Scarlet fever—Monthly averages of cases reported during the years for which data are available, 1913-1917.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Average per year.	State.	
Alabama.....	1913, 1915-1917-4 years.....	48	87	67	34	31	24	28	51	89	118	98	65	681	Alabama.	
Arizona.....	1914, 1916-2 years.....	8	12	11	7	2	6	2	1	1	1	1	7	46	Arizona.	
Arkansas.....	1917-1 year.....	27	12	20	2	5	6	28	11	27	35	35	66	254	Arkansas.	
California.....	1913-1917-5 years.....	364	322	325	257	254	162	136	121	158	323	353	359	3,147	California.	
Colorado.....	1914, 1916-1917-3 years.....	322	155	166	101	51	88	36	20	28	39	84	126	359	Colorado.	
Connecticut.....	1913-1917-5 years.....	234	69	237	222	167	132	63	44	61	106	148	157	1,770	Connecticut.	
District of Columbia.....	do.....	69	67	73	62	46	34	17	11	24	42	45	86	553	District of Columbia.	
Hawaii.....	1913-1917-5 years.....	3	2	1	3	3	4	8	7	4	3	4	0	46	Hawaii.	
Illinois.....	1913-1914, 1917-3 years.....	31	860	1,884	2,320	2,481	924	214	8	1	1	27	207	14,161	Illinois.	
Indiana.....	1913, 1915-1916, 1917-4 years.....	562	554	518	443	312	195	110	89	194	388	627	745	14,927	Indiana.	
Iowa.....	1913-1917-5 years.....	129	168	214	169	144	52	35	22	37	82	107	141	1,268	Iowa.	
Kansas.....	1913-1917-5 years.....	189	217	262	209	198	71	49	48	59	168	222	224	1,917	Kansas.	
Louisiana.....	1914-1917-4 years.....	16	13	16	13	13	9	4	9	16	26	35	32	194	Louisiana.	
Maine.....	1916, 1917-2 years.....	10	13	9	14	9	5	16	20	27	14	32	31	180	Maine.	
Maryland.....	1915-1917-3 years.....	291	221	221	204	194	113	74	48	81	185	176	185	1,908	Maryland.	
Massachusetts.....	1913, 1916, 1917-3 years.....	860	771	872	720	684	455	27	166	239	418	570	792	6,761	Massachusetts.	
Michigan.....	1913-1917-5 years.....	693	705	867	778	653	358	178	143	448	571	617	6,638	Michigan.		
Minnesota.....	do.....	450	528	606	617	423	281	141	124	210	286	377	472	3,272	Minnesota.	
Mississippi.....	1914-1917-4 years.....	38	47	47	30	28	23	18	35	56	80	61	57	488	Mississippi.	
Montana.....	1913, 1914, 1916, 1917-4 years.....	76	59	85	97	56	34	34	34	89	81	112	108	867	Montana.	
Nevada.....	1913, 1915, 1917-3 years.....	23	17	20	18	7	9	4	4	10	32	170	170	1,921	Nevada.	
New Jersey.....	1913-1917-5 years.....	615	580	728	648	553	437	189	114	135	244	337	319	4,969	New Jersey.	
New York.....	do.....	1,746	1,833	2,290	2,214	1,612	1,249	649	352	422	601	884	1,228	15,600	New York.	
North Dakota.....	1917-1 year.....	37	97	87	48	33	15	22	12	35	35	71	71	528	North Dakota.	
Ohio.....	1914-1917-4 years.....	1,180	976	1,100	924	741	443	267	233	421	798	1,008	884	8,928	Ohio.	
Oklahoma.....	1913, 1915, 1917-3 years.....	165	196	139	74	54	30	35	73	126	161	267	206	3,161	Oklahoma.	
Oregon.....	1913, 1915-5 years.....	62	54	69	67	69	42	37	19	23	41	60	59	611	Oregon.	
Pennsylvania.....	1916, 1917-2 years.....	1,008	887	980	885	784	525	316	269	383	649	808	825	8,260	Pennsylvania.	
Porto Rico.....	1914, 1917-2 years.....	1	0	0	0	0	(1)	0	0	0	0	0	0	0	Porto Rico.	
Rhode Island.....	1914, 1916, 1917-3 years.....	1,164	1,164	1,164	1,164	1,164	1,164	1,164	1,164	1,164	1,164	1,164	1,164	1,164	Rhode Island.	
South Carolina.....	1913-1917-5 years.....	19	10	10	12	6	11	12	55	55	55	55	55	55	55	South Carolina.
South Dakota.....	1913, 1915, 1917-3 years.....	66	80	94	62	42	22	22	14	16	48	40	39	366	South Dakota.	
Texas.....	1915, 1917-2 years.....	122	126	77	61	37	63	51	28	71	200	291	337	1,464	Texas.	
Utah.....	1913, 1915, 1917-4 years.....	165	176	137	113	106	65	41	23	40	91	147	138	1,242	Utah.	
Vermont.....	1914-1917-4 years.....	58	83	66	55	57	57	52	33	24	18	36	49	570	Vermont.	
Virginia.....	do.....	207	177	114	94	57	52	33	63	120	212	267	164	1,900	Virginia.	
Washington.....	1913-1917-5 years.....	93	82	88	70	55	47	33	20	14	19	27	61	64	588	Washington.
West Virginia.....	1917-1 year.....	70	47	52	38	30	14	14	12	61	105	105	122	347	3,798	West Virginia.
Wisconsin.....	1913-1917-3 years.....	434	416	607	496	394	282	20	20	112	287	347	347	3,798	Wisconsin.	
Wyoming.....	do.....	14	21	20	16	12	8	8	9	16	12	22	22	183	Wyoming.	

1 Less than 1.

Cases of scarlet fever reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.	
Alabama.....	45	29	30	22	17	11	17	63	77	59	25	44	439	Alabama.	
Arizona.....	28	15	5	8	4	3	2	3	2	3	1	1	91	Arizona.	
Arkansas.....	79	67	46	19	63	6	24	19	17	9	16	24	389	Arkansas.	
California.....	566	438	382	285	440	236	141	133	120	198	122	126	3,196	California.	
Colorado.....	217	244	285	283	217	151	64	42	38	59	43	50	1,624	Colorado.	
Connecticut.....	173	190	178	175	160	103	50	37	68	118	97	187	1,450	Connecticut.	
District of Columbia.....	208	165	161	151	91	32	16	12	16	16	12	23	883	District of Columbia.	
Florida.....	40	56	9	4	3	8	4	7	5	6	16	11	138	Florida.	
Illinois.....	738	611	561	565	362	148	160	101	221	189	201	208	4,025	Illinois.	
Indiana.....	694	593	557	500	291	159	94	94	156	184	163	210	3,670	Indiana.	
Iowa.....	422	461	384	428	366	151	45	47	79	110	117	110	2,730	Iowa.	
Kansas.....	548	484	376	533	346	152	57	47	92	134	90	79	3,118	Kansas.	
Louisiana.....	32	8	13	9	9	7	5	17	9	10	21	10	150	Louisiana.	
Maine.....	27	22	31	22	20	20	43	26	83	13	5	10	60	Maine.	
Maryland.....	174	212	206	164	144	48	47	38	60	62	47	111	1,313	Maryland.	
Massachusetts.....	608	537	540	626	487	287	200	122	117	225	287	338	4,490	Massachusetts.	
Michigan.....	1,048	830	1,021	720	529	304	286	220	230	345	286	350	6,116	Michigan.	
Minnesota.....	325	546	815	606	545	338	131	94	86	102	53	127	3,768	Minnesota.	
Mississippi.....	113	23	24	82	41	23	28	22	57	50	48	55	568	Mississippi.	
Montana.....	32	313	224	158	177	104	56	54	47	87	22	42	1,609	Montana.	
Nebraska.....	171	217	153	192	199	152	292	26	30	33	25	40	1,429	Nebraska.	
New Jersey.....	396	378	348	288	159	124	64	129	129	177	217	217	2,817	New Jersey.	
New York.....	1,346	1,183	1,415	1,349	1,076	649	448	302	410	587	650	824	10,139	New York.	
North Carolina.....	85	32	22	21	21	23	32	86	109	135	114	74	71	804	North Carolina.
North Dakota.....	129	175	134	94	68	37	14	6	10	2	12	8	689	North Dakota.	
Oregon.....	1,010	1,167	954	876	701	400	263	213	367	571	382	395	7,298	Oregon.	
Pennsylvania.....	951	972	1,119	905	739	425	310	219	329	451	338	330	7,388	Pennsylvania.	
Rhode Island.....	47	48	63	78	75	44	32	21	31	37	46	47	71	688	Rhode Island.
South Carolina.....	10	6	9	2	3	...	50	19	9	8	15	6	4	91	South Carolina.
South Dakota.....	107	142	127	149	92	50	30	22	38	47	47	47	898	South Dakota.	
Texas.....	210	166	238	198	63	33	23	34	31	29	9	13	1,045	Texas.	
Utah.....	288	321	331	163	214	143	61	55	34	60	27	26	1,723	Utah.	
Vermont.....	48	35	17	26	31	17	18	23	23	11	27	27	312	Vermont.	
Virginia.....	102	114	82	82	66	31	15	29	31	68	49	33	707	Virginia.	
Washington.....	182	343	442	384	417	363	202	180	135	144	46	68	2,876	Washington.	
West Virginia.....	40	76	27	33	18	17	28	23	43	21	21	21	372	West Virginia.	
Wisconsin.....	715	645	609	475	472	241	209	128	185	354	180	188	4,401	Wisconsin.	
Wyoming.....	78	527	106	59	69	40	6	2	6	5	1	1	906	Wyoming.	

Deaths registered as due to scarlet fever, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sep.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	2	1	0	0	1	0	1	2	2	0	3	0	12	Alabama.
Arizona.....	1	2	1	0	2	3	1	0	1	1	1	0	2	Arizona.
Arkansas.....	5	6	0	1	2	3	0	0	1	1	1	0	17	Arkansas.
California.....	8	6	11	11	4	4	4	4	3	3	2	0	54	California.
Connecticut.....	1	3	0	4	5	0	1	1	1	1	1	0	86	Connecticut.
District of Columbia.....	0	0	1	0	1	0	1	1	0	0	1	0	3	District of Columbia.
Florida.....	3	1	5	5	1	1	1	1	1	1	1	1	16	Florida.
Illinois.....	32	19	21	22	9	6	4	4	3	3	6	7	154	Illinois.
Indiana.....	25	17	18	18	10	3	3	3	3	3	6	5	117	Indiana.
Iowa.....	8	16	23	20	18	6	6	6	6	6	6	7	123	Iowa.
Kansas.....	11	11	14	15	4	3	3	3	3	3	6	7	73	Kansas.
Kentucky.....	5	2	2	6	5	3	3	3	3	3	6	7	43	Kentucky.
Louisiana.....	2	0	0	0	0	0	0	0	0	0	0	0	3	Louisiana.
Maine.....	2	3	1	1	1	1	1	1	1	1	1	1	13	Maine.
Maryland.....	2	1	5	3	5	1	1	1	1	1	1	1	23	Maryland.
Massachusetts.....	14	9	10	9	4	6	3	3	3	3	4	5	76	Massachusetts.
Michigan.....	24	27	14	31	26	10	6	6	6	6	6	6	168	Michigan.
Minnesota.....	13	17	25	26	33	10	7	10	6	3	3	5	117	Minnesota.
Nebraska.....	2	6	8	8	8	5	4	4	3	2	2	0	40	Nebraska.
New Jersey.....	5	6	9	10	4	5	5	5	3	3	8	1	57	New Jersey.
New York.....	50	34	37	39	35	26	18	17	7	20	11	18	300	New York.
North Dakota.....	3	3	4	4	4	2	2	2	1	1	1	1	19	North Dakota.
Ohio.....	13	15	23	16	20	7	7	7	7	7	10	17	110	Ohio.
Oklahoma.....	2	4	1	3	3	1	1	1	1	1	0	1	16	Oklahoma.
Oregon.....	1	1	3	2	1	3	0	0	0	0	1	0	16	Oregon.
Pennsylvania.....	18	22	18	27	18	7	7	7	7	7	7	15	16	Pennsylvania.
Porto Rico.....	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Rhode Island.....	4	3	4	4	4	4	4	4	4	4	4	4	2	Rhode Island.
South Carolina.....	1	1	1	1	1	1	1	1	1	1	1	1	3	South Carolina.
South Dakota.....	3	4	8	3	4	3	4	4	4	4	4	4	26	South Dakota.
Texas.....	7	3	6	4	11	6	4	4	4	4	4	4	28	Texas.
Utah.....	4	6	4	1	1	1	1	1	1	1	1	1	47	Utah.
Vermont.....	1	1	1	1	1	1	1	1	1	1	1	1	1	Vermont.
Washington.....	2	4	17	15	15	15	15	15	15	15	15	15	71	Washington.
Wisconsin.....	22	22	24	24	24	24	24	24	24	24	24	24	178	Wisconsin.

Scarlet fever—Average number of cases reported per annum and average indicated morbidity rates, 1913–1917; cases reported, deaths registered, and indicated mortality, morbidity, and fatality rates, 1918.

State.	Estimated Population July 1, 1918.	Average.		1918		State.		
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.	Total deaths registered.	Deaths per 1,000 inhabitants.	Fatalities per 100 cases.
Alabama.....	2,395,270	1913, 1915–1917—4 years.	681	0.3	439	0.2	12	0.005
Arizona.....	272,034	1914, 1916–2 years.	46	.2	51	.3	2	2.7
Arkansas.....	1,762,965	1917–1 year.	254	.1	380	.2	17	.007
California.....	3,119,113	1913–1917—5 years.	3,147	1.1	3,194	1.0	54	.009
Colorado.....	1,014,381	1914, 1916, 1917—3 years.	950	1.0	1,924	1.6	54	.017
Connecticut.....	1,286,368	1913–1917—5 years.	1,770	1.4	1,850	1.2	35	.027
Delaware.....	216,041	1913–1917—5 years.	555	1.5	883	2.2	3	.014
District of Columbia.....	401,681	1913–1917—5 years.	555	1.5	138	.1	0	.040
Florida.....	938,377	1913–1917—5 years.	46	.5	4,026	.6	134	.024
Hawaii.....	223,119	1913–1917—5 years.	14,611	2.3	1,611	1.6	117	.32
Idaho.....	461,186	1913–1917—1 year.	4,327	1.6	3,870	1.3	123	.041
Illinois.....	6,317,734	1913, 1915, 1917—3 years.	1,928	1.6	2,998	2.0	1,177	.055
Indiana.....	2,854,167	1913–1917—5 years.	1,917	1.1	3,118	1.7	72	.038
Iowa.....	2,249,771	1913, 1914, 1916, 1917—4 years.	1,917	1.1	3,118	1.7	43	.018
Kansas.....	1,871,956	1913–1917—5 years.	1,406	.5	2,722	1.6	13	.002
Kentucky.....	2,406,547	1914–1917—4 years.	1,982	.7	1,904	.1	3	.002
Louisiana.....	1,982,778	1914–1917—4 years.	180	.1	150	.1	3	.002
Maine.....	782,191	1916, 1917—3 years.	1,908	1.4	1,813	.6	13	.017
Maryland.....	1,384,639	1915–1917—3 years.	1,908	1.4	1,813	.9	23	.017
Massachusetts.....	3,832,790	1913, 1916, 1917—3 years.	6,761	1.8	4,490	1.2	76	.020
Michigan.....	3,138,678	1913–1917—5 years.	5,688	1.9	6,116	2.0	168	.054
Minnesota.....	2,345,287	do.	4,272	1.9	3,788	1.6	157	.067
Mississippi.....	2,001,466	1914–1917—4 years.	4,488	.3	568	.3	7	.003
Missouri.....	3,446,493	1913, 1914, 1916, 1917—4 years.	807	1.9	1,606	3.3	70	.020
Montana.....	480,316	1913–1917—3 years.	170	.7	1,429	1.1	40	.031
Nebraska.....	1,286,877	1913, 1915, 1917—3 years.	4,966	1.7	2,817	.9	57	.019
Nevada.....	114,742	1913–1917—3 years.	15,660	1.5	10,130	1.0	300	.028
New Jersey.....	3,080,371	1913–1917—5 years.	10,646	.9	8,924	.9	30	.020
New York.....	10,646,989	do.	10,646	.9	10,646	.9	30	.020
North Carolina.....	2,468,025	1917—1 year.	529	.7	639	.9	19	.024
North Dakota.....	791,437	1917—1 year.	8,928	1.7	7,299	1.4	170	.032
Oklahoma.....	5,277,814	1914–1917—4 years.	1,516	.7	553	.6	16	.007
Oregon.....	885,243	1913, 1915, 1917—3 years.	6,611	1.0	7,305	.8	174	.017
Pennsylvania.....	8,793,067	1916, 1917—2 years.	8,360	1.0	7,305	.8	174	.020

Scarlet fever—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated mortality, morbidity, and fatality rates, 1918—Continued.

State.	Estimated population July 1, 1918.	Average.		1918		1918		Fatalities per 100 cases.	State.
		Years.	Cases reported per annum.	Cases per 1,000 inhabitants.	Total cases reported.	Cases per 1,000 inhabitants.	Total deaths registered.		
Porto Rico.....	1,247,677	1914, 1915—2 years.	2	593	0.9	28	0.001	Porto Rico.
Rhode Island.....	637,415	1914, 1916, 1917—3 years.....	867	1.4	91	1.1	3	.044	Rhode Island.
South Carolina.....	1,660,334	1913—1917—5 years.....	288	.8	898	1.2	25	.002	South Carolina.
South Dakota.....	725,334	1913, 1914, 1917—3 years.....	566	.3	1,044	.2	29	.033	South Dakota.
Texas.....	4,691,779	1913—1915, 1917—2 years.....	1,442	2.9	1,723	3.8	47	.006	Texas.
Utah.....	453,948	1915—1916, 1917—4 years.....	579	1.6	312	.9	10	.104	Utah.
Vermont.....	386,192	1914—1917—4 years.....	1,600	.7	707	.3	1	.027	Vermont.
Virginia.....	2,234,320	do.....	786	.5	2,876	1.7	71	.043	Virginia.
Washington.....	1,660,378	1913—1917—5 years.....	596	.4	372	.3	1	.043	Washington.
West Virginia.....	1,438,165	1913—1917—1 year.....	3,786	1.5	4,401	1.7	179	.07	West Virginia.
Wisconsin.....	2,552,983	1913, 1914—5 years.....	183	1.1	906	4.8	4.1	Wisconsin.
Wyoming.....	190,380	do.....	Wyoming.

Cases of septic sore throat reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Connecticut.....	6	6	1	2	86	1	1	1	1	2	1	6	113	Connecticut.
Illinois.....	9	6	10	5	3	2	1	6	10	4	10	72	Illinois.	
Kansas.....	2	5	1	1	3	2	0	1	3	8	6	32	Kansas.	
Louisiana.....	0	0	0	0	0	0	0	0	0	0	0	3	Louisiana.	
Maryland.....	9	1	2	0	0	1	3	0	0	1	1	3	Maryland.	
Massachusetts.....	16	12	13	10	12	3	0	0	2	4	4	107	Massachusetts.	
Minnesota.....	0	0	0	0	0	0	0	0	0	0	0	0	Minnesota.	
Nebraska.....	28	20	6	4	5	2	1	0	3	6	9	20	Nebraska.	
New York.....	141	New York.	
North Carolina.....	0	1	0	3	1	2	0	0	0	1	1	1	North Carolina.	
Ohio.....	22	0	10	35	1	23	1	22	1	13	2	11	Ohio.	
Rhode Island.....	10	Rhode Island.		
South Carolina.....	140	South Carolina.		

Deaths registered as due to septic sore throat, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Connecticut.....	7	6	1	1	6	4	3	6	10	19	3	1	76	Connecticut.
Illinois.....	2	3	2	5	2	5	7	2	6	3	2	1	44	Illinois.
Iowa.....	0	1	1	3	2	0	0	0	0	0	0	0	22	Iowa.
Kansas.....	5	2	2	5	3	3	3	4	2	2	2	1	25	Kansas.
Louisiana.....	3	1	0	0	0	0	0	0	0	0	0	0	3	Louisiana.
Maryland.....	3	2	2	1	0	0	0	0	0	0	0	0	3	Maryland.
Massachusetts.....	3	1	2	2	0	0	2	1	0	1	1	1	10	Massachusetts.
Nebraska.....	1	1	1	1	0	0	0	1	1	1	1	0	9	Nebraska.
New York.....	15	5	12	8	6	8	6	14	5	10	6	15	111	New York.
North Dakota.....	1	1	1	1	1	1	1	1	1	1	1	1	6	North Dakota.
Ohio.....	1	1	1	1	1	1	1	1	1	1	1	1	6	Ohio.
Rhode Island.....	1	1	1	1	1	1	1	1	1	1	1	1	6	Rhode Island.
South Carolina.....	1	1	1	1	1	1	1	1	1	1	1	1	2	South Carolina.

Sepic sore throat—Cases reported, deaths registered, and indicated morbidity, mortality, and fatality rates, 1918.

Smallpox—Monthly averages of cases reported during the years for which data are available, 1913-1917.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sep.	Oct.	Nov.	Dec.	Average per year.	State.		
Alabama.....	1913, 1915-1917—4 years.....	98	93	86	77	65	52	25	16	6	21	24	61	624	Alabama.		
Arkansas.....	1917—1 year.....	156	232	571	251	245	91	28	20	13	44	93	272	2,016	Arkansas.		
California.....	1913, 1915-1917—3 years.....	68	74	64	65	44	29	19	21	19	25	25	31	2,475	California.		
Colorado.....	1913, 1914, 1915, 1917—4 years.....	30	25	30	10	25	27	22	9	9	23	40	28	289	Colorado.		
Connecticut.....	1913-1917—3 years.....	23	20	25	11	15	15	3	1	3	23	15	27	181	Connecticut.		
District of Columbia.....	do.....	9	5	13	3	13	10	3	1	1	1	1	3	3	64	District of Columbia.	
Florida.....	1913—1 year.....	246	211	217	113	94	88	71	29	8	54	17	18	1,181	Florida.		
Idaho.....	1913, 1914, 1917—3 years.....	73	96	27	42	8	25	184	73	81	184	247	348	3,819	Idaho.		
Illinois.....	1913, 1914, 1916, 1917—4 years.....	438	520	580	434	390	386	273	133	99	69	117	379	3,271	Illinois.		
Indiana.....	1913, 1914, 1916, 1917—4 years.....	261	397	469	390	296	257	144	72	16	25	69	148	2,244	Indiana.		
Iowa.....	1913, 1914, 1916, 1917—4 years.....	308	306	364	321	295	257	168	67	22	27	74	202	2,225	Iowa.		
Kansas.....	1917-1917—5 years.....	174	212	306	234	234	185	165	22	23	9	5	16	230	1,999	Kansas.	
Louisiana.....	1914-1917—4 years.....	108	94	89	94	95	12	44	52	12	7	41	40	199	647	Louisiana.	
Maine.....	1917—1 year.....	0	0	5	0	5	0	5	0	0	0	0	8	28	600	Maine.	
Maryland.....	1915-1917—3 years.....	193	1916, 1917—3 years.....	9	7	4	18	16	3	4	0	0	0	8	87	Maryland.	
Massachusetts.....	1913-1917—3 years.....	4	3	11	11	13	21	2	3	2	0	4	7	4	85	Massachusetts.	
Michigan.....	1913-1917—5 years.....	188	154	171	173	155	134	91	37	36	108	108	204	274	1,725	Michigan.	
Minnesota.....	do.....	292	273	271	244	188	145	90	44	37	93	150	207	2,082	Minnesota.		
Mississippi.....	1914-1917—4 years.....	216	222	213	251	165	92	43	41	37	50	61	83	204	1,641	Mississippi.	
Montana.....	1913, 1914, 1915, 1917—4 years.....	130	111	93	95	81	85	57	51	31	32	47	116	116	1,991	Montana.	
Nevada.....	1913, 1915, 1917—3 years.....	35	15	9	13	33	11	5	1	0	1	1	1	2	122	Nevada.	
New Jersey.....	1913-1917—5 years.....	7	3	14	17	3	4	0	(1)	(1)	(1)	1	1	1	51	51	New Jersey.
New York.....	do.....	87	74	34	42	26	24	19	7	9	28	41	427	427	New York.		
North Dakota.....	1913-1917—2 years.....	22	30	49	76	27	44	30	10	18	49	38	87	41	480	North Dakota.	
Ohio.....	1913-1917—5 years.....	381	339	383	373	344	301	161	90	75	222	370	488	3,547	Ohio.		
Oklahoma.....	1913, 1915, 1917—3 years.....	483	369	480	359	277	197	118	66	64	91	111	425	3,703	Oklahoma.		
Oregon.....	1913-1917—5 years.....	68	68	59	57	41	35	22	18	7	14	7	10	47	460	Oregon.	
Pennsylvania.....	1916, 1917—2 years.....	11	8	20	29	35	22	18	7	9	11	34	34	238	Pennsylvania.		
Porto Rico.....	1917—1 year.....	1	11	13	40	8	1	4	0	3	1	3	0	85	85	Porto Rico.	
South Carolina.....	1913-1917—5 years.....	63	43	58	69	38	32	16	10	3	11	15	25	383	South Carolina.		
South Dakota.....	1913, 1914, 1917—3 years.....	117	141	176	135	102	78	42	24	20	31	64	103	1,064	South Dakota.		
Texas.....	1913, 1915, 1917—3 years.....	234	232	302	247	164	104	63	21	32	108	190	190	1,718	Texas.		
Utah.....	1913, 1915, 1917—3 years.....	212	190	144	124	106	79	56	34	50	63	99	1,234	Utah.			
Vermont.....	1913-1917—5 years.....	19	13	10	17	20	10	12	6	1	1	3	6	118	Vermont.		
Virginia.....	do.....	249	210	212	182	136	84	57	37	22	39	53	98	1,379	Virginia.		
Washington.....	1917—1 year.....	107	76	97	90	86	70	38	25	21	30	40	50	720	Washington.		
West Virginia.....	1913-1917—5 years.....	5	4	8	40	21	30	37	21	36	65	96	143	143	West Virginia.		
Wisconsin.....	do.....	285	217	211	207	158	104	78	62	35	93	151	237	1,838	Wisconsin.		
Wyoming.....	do.....	8	17	8	2	2	9	7	2	4	4	4	3	81	81	Wyoming.	

^aLess than one.

Cases of smallpox reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	537	1,000	761	408	255	182	85	75	45	31	22	286	3,647	Alabama.
Arizona.....	12	5	66	34	56	17	104	3	0	0	0	0	3,230	Arizona.
Arkansas.....	762	733	212	457	200	62	97	52	14	13	16	16	2,590	Arkansas.
California.....	42	59	111	81	172	97	43	52	119	97	102	102	1,069	California.
Colorado.....	175	176	250	239	266	133	92	43	82	63	80	102	1,680	Colorado.
Connecticut.....	3	2	21	12	11	32	1	1	1	1	1	1	94	Connecticut.
District of Columbia.....	7	2	17	5	4	6	5	6	5	5	4	4	48	District of Columbia.
Florida.....	6	11	10	3	8	6	6	6	5	5	5	5	55	Florida.
Illinois.....	742	744	645	557	571	189	103	73	26	42	36	114	3,852	Illinois.
Indiana.....	1,003	1,255	1,027	804	688	314	188	86	30	13	101	132	5,682	Indiana.
Iowa.....	650	569	559	414	399	225	121	54	47	79	109	109	3,807	Iowa.
Kansas.....	994	1,419	1,640	1,132	1,077	435	114	89	26	89	48	67	7,130	Kansas.
Louisiana.....	73	158	153	142	136	111	26	30	5	34	12	70	950	Louisiana.
Maine.....	92	103	124	29	23	182	15	2	19	7	20	55	55	Maine.
Maryland.....	16	12	17	83	19	12	0	12	18	9	15	7	219	Maryland.
Massachusetts.....	3	2	10	2	3	1	1	1	2	1	2	2	27	Massachusetts.
Michigan.....	1,083	921	689	441	421	247	119	92	87	137	111	111	4,417	Michigan.
Minnesota.....	210	285	355	304	217	213	107	44	52	68	207	180	2,232	Minnesota.
Mississippi.....	631	690	622	662	418	160	90	41	57	58	63	63	3,601	Mississippi.
Montana.....	246	172	157	118	125	103	64	17	28	50	18	14	1,104	Montana.
Nebraska.....	451	597	600	640	546	223	139	40	29	152	78	98	3,96	Nebraska.
New Jersey.....	1	3	32	0	2	16	2	2	0	0	0	0	3	New Jersey.
New York.....	140	68	52	95	35	33	25	6	7	5	7	7	535	New York.
North Carolina.....	45	159	147	105	80	48	68	40	34	34	41	98	899	North Carolina.
North Dakota.....	57	52	72	51	31	47	11	3	9	2	8	8	345	North Dakota.
Ohio.....	2,068	1,642	1,676	1,276	1,214	628	198	121	226	460	269	10,227	Ohio.	
Oregon.....	20	25	32	25	66	53	56	28	49	33	61	47	453	Oregon.
Pennsylvania.....	86	111	89	101	96	52	51	14	3	3	5	2	612	Pennsylvania.
Porto Rico.....	12	1	4	5	2	2	1	1	0	0	0	0	27	Porto Rico.
Rhode Island.....	0	0	0	1	0	0	1	0	0	0	0	0	2	Rhode Island.
South Carolina.....	4	16	12	8	28	3	10	4	5	94	33	8	214	South Carolina.
South Dakota.....	124	131	113	138	141	161	77	18	16	48	26	67	1,050	South Dakota.
Texas.....	431	875	948	556	729	280	101	67	125	125	50	152	4,338	Texas.
Utah.....	377	312	236	208	222	235	130	124	107	126	91	99	2,263	Utah.
Vermont.....	5	6	19	3	1	1	1	1	1	1	1	1	65	Vermont.
Virginia.....	144	172	209	348	206	120	73	31	39	88	78	78	1,605	Virginia.
Washington.....	112	62	198	190	171	190	137	113	116	116	116	116	1,676	Washington.
West Virginia.....	220	301	184	218	145	63	35	22	24	16	24	14	1,286	West Virginia.
Wisconsin.....	459	334	313	369	224	240	159	63	193	98	76	76	2,592	Wisconsin.
Wyoming.....	6	45	36	55	51	20	8	9	19	35	17	11	312	Wyoming.

Deaths registered as due to smallpox, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	2	4	5	1	1	2	0	0	0	0	1	0	16	Alabama.
Arizona.....	2	0	1	5	3	1	0	1	0	1	1	0	3	Arizona.
Arkansas.....	0	2	2	5	3	1	1	1	0	1	1	0	21	Arkansas.
California.....	2	1	4	1	1	1	1	1	1	1	1	1	3	California.
Illinois.....	4	1	7	2	3	3	1	3	1	1	1	1	24	Illinois.
Indiana.....	3	2	3	0	0	0	0	0	0	0	1	1	10	Indiana.
Iowa.....	2	3	4	7	8	3	0	0	0	0	0	0	26	Iowa.
Kansas.....	3	4	0	0	1	0	0	0	0	0	0	1	1	Kansas.
Kentucky.....	2	4	0	0	1	0	1	0	0	0	0	0	8	Kentucky.
Louisiana.....	2	1	0	2	0	1	0	0	0	2	1	1	16	Louisiana.
Michigan.....	0	0	4	2	2	1	0	1	0	0	0	1	11	Michigan.
Minnesota.....	0	0	0	0	0	3	1	1	0	0	0	1	0	Minnesota.
Nebraska.....	0	1	2	0	1	0	1	0	0	0	0	1	0	Nebraska.
New York.....	1	1	1	1	1	1	1	1	1	1	1	1	1	New York.
North Dakota.....	1	1	2	1	2	3	4	4	4	4	4	4	1	North Dakota.
Ohio.....	5	4	6	5	6	2	3	2	1	0	0	0	22	Ohio.
Oklahoma.....	2	4	6	5	6	1	1	1	0	0	0	0	0	Oklahoma.
Pennsylvania.....	1	1	1	1	1	1	1	1	1	1	1	1	2	Pennsylvania.
Rhode Island.....	1	1	1	1	1	1	1	1	1	1	1	1	1	Rhode Island.
South Dakota.....	25	22	11	6	9	11	3	3	3	8	11	8	124	South Dakota.
Texas.....	1	3	1	1	1	1	2	2	2	2	2	1	10	Texas.
Utah.....	1	1	1	1	2	1	1	1	1	1	1	1	2	Utah.
Washington.....	1	1	1	1	2	2	2	2	2	2	2	1	4	Washington.
Wisconsin.....	1	1	1	1	1	1	1	1	1	1	1	1	7	Wisconsin.
Wyoming.....	1	1	1	1	2	2	2	2	2	2	2	1	6	Wyoming.

Smallpox—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated morbidity, mortality, and fatality rates, 1918.

State.	Estimated population July 1, 1918.	Average.		1918.		State.	
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.		
Alabama.....	2,395,270	1913, 1915-1917-4 years.....	624	0.27	3,647	1.52	Alabama.
Arizona.....	272,134	1917-1 year.....	2,016	1.14	200	.007	Arizona.
Arkansas.....	1,792,965	1917-5 years.....	475	1.74	2,590	.011	Arkansas.
California.....	3,119,412	1913-1917-5 years.....	389	1.17	1,068	.012	California.
Colorado.....	1,014,581	1913, 1914-1916, 1917-4 years.....	389	1.17	1,680	.012	Colorado.
Connecticut.....	1,280,288	1913-1917-5 years.....	181	.94	1,151	.001	Connecticut.

District of Columbia.....		64	18	48	32	0	
Florida.....	1,166	1,41	56	36	0		
Idaho.....	1,133	777	1913-1917—do—year.....	1,609	1,61	1,61	
Illinois.....	491	766	do.....	3,819	63	3,842	
Indiana.....	6,317	734	1913-1914.....	3,271	1,17	5,582	
Iowa.....	2,854	167	1913-1917—5 years.....	2,224	1,00	3,307	
Kansas.....	2,224	771	1913-1914.....	2,223	1,00	1,49	
Kentucky.....	1,874	195	1913-1917—5 years.....	1,999	1,11	7,130	
Louisiana.....	1,884	778	1914-1917—4 years.....	647	.36	950	
Maine.....	752	191	1911—1 year.....	600	.78	555	
Maryland.....	1,384	539	1915-1917—3 years.....	87	.66	219	
Massachusetts.....	3,832	190	1912-1916, 1917—5 years.....	85	.02	27	
Michigan.....	3,183	678	1913-1917—5 years.....	1,725	.57	4,417	
Minnesota.....	2,345	287	do.....	2,082	.88	2,232	
Mississippi.....	2,001	466	1914-1917—4 years.....	1,641	.85	3,601	
Missouri.....	3,445	498	1913-1914, 1916, 1917—4 years.....	991	2.22	1,104	
Montana.....	488	376	1913, 1914, 1916, 1917—4 years.....	991	2.22	3,906	
Nebraska.....	1,286	877	1915, 1916, 1917—3 years.....	122	1.19	0	
Nevada.....	114	742	1913-1917—5 years.....	61	.02	65	
New Jersey.....	3,050	371	1913-1917—5 years.....	427	.04	525	
New York.....	10,646	389	do.....	427	.04	65	
North Carolina.....	2,468	125	1915, 1917—2 years.....	67	.67	860	
North Dakota.....	701	437	1913-1917—3 years.....	3,047	.70	345	
Ohio.....	5,273	314	1913-1917—5 years.....	3,070	1.45	10,227	
Oklahoma.....	2,377	629	1913-1915, 1917—3 years.....	460	.57	483	
Oregon.....	888	243	1913-1917—5 years.....	238	.03	612	
Pennsylvania.....	8,798	667	1916, 1917—2 years.....	85	.07	27	
Puerto Rico.....	1,247	677	1917—1 year.....	2	2	.02	
Rhode Island.....	637	415	1913-1917—5 years.....	383	.74	214	
South Carolina.....	1,660	834	1913-1914, 1917—3 years.....	1,064	1.38	1,050	
South Dakota.....	738	424	1913, 1915, 1917—3 years.....	1,718	.40	4,338	
Texas.....	4,601	270	1913-1915, 1917—4 years.....	1,234	2.62	2,268	
Utah.....	453	648	1913-1915, 1917—5 years.....	1,118	.33	4,99	
Vermont.....	399	192	1913-1917—5 years.....	1,234	.55	1,605	
Virginia.....	2,224	180	do.....	1,750	.64	1,670	
West Virginia.....	1,439	165	1917—1 year.....	413	.29	1,266	
Wisconsin.....	2,553	983	1913-1917—5 years.....	1,458	.75	2,592	
Wyoming.....	190	350	do.....	81	.47	313	

Tuberculosis (pulmonary)—Monthly averages of cases reported during the years for which data are available, 1916-17.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Average per year.	State.
Alabama.....	1916, 1917-2 years.	244	260	304	275	268	253	207	218	176	207	2,989	2,989	Alabama.	
Arizona.....	1916-1 year.	2	0	0	0	0	4	1	1	1	1	19	19	Arizona.	
Connecticut.....	1916, 1917-2 years.	162	177	166	183	197	170	165	168	152	161	139	2,016	Connecticut.	
Hawaii.....	1916, 1917-2 years.	86	98	78	65	71	73	61	63	57	74	879	879	Hawaii.	
Illinois.....	1916, 1917-1 year.	59	1,235	1,204	784	1,089	1,306	1,237	1,239	1,239	1,552	1,538	12,710	Illinois.	
Indiana.....	1916, 1917-2 years.	247	233	301	276	251	227	225	213	195	190	179	2,861	Indiana.	
Iowa.....	1916, 1917-2 years.	150	167	176	164	172	205	183	196	147	164	136	1,939	Iowa.	
Louisiana.....	1916, 1917-1 year.	266	259	288	272	346	202	322	289	262	224	239	235	Louisiana.	
Maryland.....	1916, 1917-2 years.	689	688	700	686	711	713	776	658	621	608	663	8,121	Maryland.	
Massachusetts.....	1916, 1917-1 year.	373	356	400	413	442	412	346	369	372	388	292	336	Massachusetts.	
Minnesota.....	1916, 1917-2 years.	24	12	29	27	19	14	21	21	23	21	12	267	Minnesota.	
Montana.....	1916, 1917-1 year.	4	2	9	8	6	5	9	4	3	5	4	64	Montana.	
Nevada.....	1916, 1917-2 years.	2,641	2,413	2,956	2,464	2,506	2,637	2,353	2,561	2,087	2,261	2,171	1,986	New York.	
New York.....	1916, 1917-1 year.	5	13	8	7	7	20	21	26	17	11	16	174	New York.	
North Dakota.....	1916-1 year.	49	44	48	48	48	27	26	30	29	31	32	31	North Dakota.	
Oregon.....	1916, 1917-2 years.	65	63	86	90	83	92	78	62	66	58	58	425	Oregon.	
Rhode Island.....	1916, 1917-1 year.	40	28	23	17	24	27	30	12	17	11	19	853	Rhode Island.	
South Carolina.....	1916, 1917-1 year.	13	10	35	35	7	14	20	12	7	9	7	267	South Carolina.	
South Dakota.....	1916, 1917-2 years.	5	6	5	6	7	4	3	4	7	6	5	146	South Dakota.	
Utah.....	1916, 1917-1 year.	18	11	17	17	38	36	28	26	16	14	21	56	Utah.	
Vermont.....	1916, 1917-2 years.	112	91	113	126	83	102	92	73	95	74	94	1,127	Vermont.	
Washington.....	1916, 1917-1 year.	34	24	40	27	34	35	10	32	25	28	36	368	Washington.	
West Virginia.....	1917-1 year.	34	24	40	27	34	35	10	32	25	28	36	368	West Virginia.	

Cases of tuberculosis (pulmonary) reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	227	238	283	283	268	245	218	212	192	188	160	195	2,688	Alabama.
District of Columbia.....	110	87	81	114	122	91	110	90	83	80	88	88	1,152	District of Columbia.
Indiana.....	201	140	425	326	300	364	316	271	237	147	141	208	3,976	Indiana.
Maryland.....	239	276	419	400	337	289	333	381	302	198	173	3,835	Maryland.	
Massachusetts.....	637	677	814	779	886	764	614	606	570	441	625	7,641	Massachusetts.	
Minnesota.....	330	278	343	417	291	338	323	300	138	266	268	3,611	Minnesota.	
Nebraska.....	33	31	13	27	21	10	44	12	10	11	14	239	Nebraska.	
New York.....	2,326	2,015	2,216	2,746	2,312	2,693	1,784	2,250	1,866	1,330	1,259	24,001	New York.	
North Dakota.....	17	8	37	15	19	7	18	18	5	4	4	168	North Dakota.	
Porto Rico.....	62	76	108	128	156	135	108	155	177	105	95	1,521	Porto Rico.	
Rhode Island.....	65	71	60	66	73	82	66	65	117	34	38	745	Rhode Island.	
South Dakota.....	13	12	11	30	22	13	22	25	26	14	34	39	261	South Dakota.
Utah.....	4	3	4	6	5	3	2	7	2	3	4	47	47	Utah.
Vermont.....	11	12	13	20	23	17	8	42	12	8	12	14	192	Vermont.
West Virginia.....	11	12	31	47	16	12	20	5	15	10	8	8	195	West Virginia.
Wisconsin.....	174	159	159	222	219	171	155	126	109	106	106	112	1,832	Wisconsin.

Deaths registered as due to tuberculosis (pulmonary), 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	State.	
													1887.	1888.
Alabama.....	206	223	233	251	238	215	194	188	156	192	168	172	2,436	Alabama.
Arizona.....	97	118	83	87	52	65	60	113	72	143	143	1,048	Arizona.	
Arkansas.....	91	115	108	120	114	93	62	64	111	68	58	1,240	Arkansas.	
California.....	403	419	498	506	434	386	380	376	358	479	464	507	5,210	California.
Connecticut.....	173	138	184	171	159	133	111	108	110	192	139	161	1,766	Connecticut.
District of Columbia.....	59	45	77	96	63	61	50	60	43	68	41	44	647	District of Columbia.
Hawaii.....	113	78	85	74	90	78	83	68	73	91	81	81	1,005	Hawaii.
Illinois.....	44	27	30	28	29	31	46	39	30	32	33	35	404	Illinois.
Indiana.....	617	587	904	847	700	696	522	491	505	659	635	646	7,619	Indiana.
Iowa.....	282	297	395	384	301	291	240	229	200	242	255	255	3,385	Iowa.
Kansas.....	132	108	116	129	116	116	95	81	90	97	122	123	1,335	Kansas.
Kentucky.....	93	85	85	72	61	61	67	54	83	74	101	911	911	Kentucky.
Louisiana.....	3.3	384	463	427	382	350	281	293	234	362	314	267	4,080	Louisiana.
Maine.....	277	240	280	319	278	288	240	211	204	281	219	223	3,040	Maine.
Maryland.....	48	75	63	79	67	58	53	37	47	80	48	64	719	Maryland.
Massachusetts.....	241	215	319	219	194	210	187	158	187	339	208	196	2,733	Massachusetts.
Michigan.....	397	402	443	527	423	388	389	393	480	398	446	5,010	Michigan.	
Minnesota.....	215	262	300	333	290	234	222	204	174	270	194	252	2,920	Minnesota.
Nebraska.....	167	172	215	194	158	130	118	118	131	144	205	191	1,976	Nebraska.
New York.....	40	50	59	59	38	31	59	31	30	37	31	31	47	New York.
North Dakota.....	1,382	1,287	1,658	1,442	1,284	1,145	1,016	1,001	929	1,522	1,235	1,139	15,040	North Dakota.
Ohio.....	519	526	24	32	24	23	18	18	16	21	21	34	270	Ohio.
Oklahoma.....	570	544	685	681	598	497	451	419	391	517	580	546	6,445	Oklahoma.
Oregon.....	43	50	65	54	46	47	42	26	20	42	50	32	918	Oregon.
Pennsylvania.....	859	910	1,094	1,097	883	797	725	758	724	1,292	939	915	10,923	Pennsylvania.
Porto Rico.....	199	201	201	199	107	235	242	197	198	242	256	2,098	2,098	Porto Rico.
Rhode Island.....	75	68	83	83	72	58	60	65	67	98	73	73	879	Rhode Island.
South Carolina.....	101	131	162	156	177	136	110	118	148	118	140	119	1,616	South Carolina.
South Dakota.....	13	12	30	22	13	22	25	26	14	34	33	39	261	South Dakota.
Texas.....	419	433	484	416	376	345	78	267	298	314	314	314	4,134	Texas.
Utah.....	14	13	18	23	10	11	12	12	22	18	15	180	180	Utah.
Vermont.....	26	20	23	34	38	27	22	19	26	32	23	22	312	Vermont.
Washington.....	82	91	103	111	83	83	58	92	94	98	98	98	1,087	Washington.
Wisconsin.....	140	163	194	233	203	154	182	166	159	150	150	150	2,065	Wisconsin.

Tuberculosis (pulmonary)—Average cases reported per annum and average indicated morbidity rates, 1916–1917; cases reported, deaths registered, indicated morbidity and mortality rates, and number of cases reported for each death registered, 1918.

State.	Estimated population July 1, 1918.	Average.		1918		Number of cases reported for each death registered.			
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.				
Alabama.	2,395,270	1916, 1917—2 years.	2,987	1.3	2,688	1.1	2,436	1.0	1.1
Arizona.	272,034	1916—1 year.	.19	.1			1,048	3.9	
Arkansas.	1,782,905						1,240		
California.	3,119,412						5,210		
Connecticut.	1,280,268	1916—1 year.	2,016	1.6	1,132	2.8	1,766	1.4	1.7
District of Columbia.	401,681						1,005		
Florida.	938,877						1,404		
Hawaii.	225,419	1916, 1917—2 years.		879	4.0		7,619	1.8	
Illinois.	6,916,734	1917—1 year.		12,710	2.0		3,885	1.2	
Indiana.	2,854,167	1916, 1917—2 years.		2,861	1.0	3,076	1,335	.6	
Iowa.	2,224,771								
Kansas.	1,874,105						911		
Kentucky.	2,408,547						4,080	1.7	
Louisiana.	1,884,778	1916—1 year.		1,899	1.1		3,040	1.6	
Maine.	782,191						719		
Maryland.	1,384,538	1916, 1917—2 years.		3,246	2.4	3,658	2.8		
Massachusetts.	3,532,790	1916, 1917—2 years.		8,121	2.2	7,841	2.0		
Michigan.	3,133,678						2,733		
Minnesota.	2,945,287	1916, 1917—2 years.		4,479	2.0	3,611	1.5		
Mississippi.	2,001,456						2,976		
Missouri.	3,448,498						2,652		
Montana.	486,376	1916, 1917—2 years.		267	.6		4,145		
Nebraska.	1,296,877								
Nevada.	11,472	1917—1 year.					480		
New York.	10,646,980	1916, 1917—2 years.		.94	6				
North Dakota.	701,437	1917—1 year.		28,976	2.8	24,001	2.3	15,040	1.4
Ohio.	5,273,814			174	2	168		6,210	1.3
Oklahoma.	2,377,620						6,445		1.2
Oregon.	888,243	1916—1 year.					918		
Pennsylvania.	8,798,067						517		
Puerto Rico.	1,207,677						10,923		
Rhode Island.	657,415	1916, 1917—2 years.					2,508		
South Carolina.	1,600,834	1917—1 year.		853	1.4	1,521		819	1.4
South Dakota.	735,434	1917—1 year.		267	.2	745		1,616	1.0
							261		.4

Tuberculosis (all forms)—Monthly averages of cases reported during the years for which data are available, 1916-17.

State.	Years.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	Average per year.	State.	
Texas.....	4,601,279	1617—1 year.	50	1	47	—	—	—	—	—	4,134	.9	.9	Texas.	
Utah.....	453,648	1916, 1917—2 years.	66	43	565	64	54	94	47	43	24	15	.86	Arkansas.	
Vermont.....	386,192	1916, 1917—2 years.	137	156	158	173	149	152	153	140	152	161	134	California.	
Washington.....	1,680,578	1916, 1917—2 years.	83	80	95	94	92	84	78	84	84	69	1,087	Connecticut.	
West Virginia.....	1,459,165	1917—1 year.	93	101	86	68	77	74	68	62	75	80	94	Vermont.	
Wisconsin.....	2,553,983	1917—1 year.	113	1250	1,211	1,003	866	1,160	1,313	1,245	1,448	1,559	1,357	1,087	Washington.
		330	362	408	377	380	338	364	304	273	264	261	221	1,087	West Virginia.
Kansas.....	128	126	147	112	145	127	119	131	107	95	108	100	1,445	Panhandle.	
Louisiana.....	197	221	183	164	174	201	127	122	139	130	188	187	1,445	Arkansas.	
Maine.....	52	22	24	36	79	43	34	43	37	32	33	33	503	Louisiana.	
Massachusetts.....	744	726	748	775	733	812	723	883	639	690	714	839	5,988	Maine.	
Michigan.....	402	537	537	566	538	593	564	387	419	352	660	660	5,986	Massachusetts.	
Minnesota.....	410	393	442	454	453	438	381	408	395	398	320	368	4,907	Michigan.	
Mississippi.....	513	519	588	620	594	637	592	514	438	432	320	320	6,852	Minnesota.	
New Jersey.....	809	696	845	855	752	718	687	542	681	620	541	541	8,597	Mississippi.	
New York.....	2,697	2,523	2,510	2,792	2,121	2,415	2,688	2,121	2,237	2,190	2,016	2,016	20,613	Panhandle.	
North Dakota.....	6	14	10	14	21	21	22	19	16	18	12	21	1,199	New York.	
Ohio.....	527	530	689	623	582	568	432	46	424	434	394	6,225	North Dakota.		
Oklahoma.....	54	54	72	42	40	46	34	32	33	22	52	20	501	Ohio.	
Oregon.....	43	37	71	56	46	49	51	48	36	41	43	44	565	Oklahoma.	
Pennsylvania.....	1,086	835	1,117	1,136	1,068	1,034	1,009	1,009	795	801	778	769	11,457	Oregon.	
Porto Rico.....	1,59	63	66	73	71	82	68	99	99	116	85	103	951	Pennsylvania.	
Rhode Island.....	68	65	62	50	50	50	67	73	61	56	62	62	903	Porto Rico.	
South Carolina.....	17	18	23	10	18	22	39	14	49	14	21	21	251	Rhode Island.	
South Dakota.....	13	9	12	36	12	14	20	14	7	13	8	8	157	South Carolina.	
Texas.....	147	88	106	84	69	78	55	103	19	147	149	149	1,239	South Dakota.	
Virginia.....	321	288	317	285	220	288	283	227	227	212	232	232	3,370	Texas.	
West Virginia.....	35	24	41	28	35	33	25	28	28	36	44	44	374	Virginia.	
Wisconsin.....	140	117	182	136	177	186	213	107	115	122	116	116	1,800	West Virginia.	
Wyoming.....	6	4	2	4	2	2	2	2	2	2	6	6	56	Wyoming.	

August 15, 1919.

1890

Cases of tuberculosis (all forms) reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	238	283	261	297	284	265	235	230	200	190	173	205	2,870	Alabama.
Arkansas.....	42	74	123	130	127	61	59	43	43	56	52	63	804	Arkansas.
California.....	658	628	585	567	810	623	713	703	582	465	444	35	7,534	California.
Colorado.....	85	110	76	4	93	47	59	50	40	30	15	630	Colorado.
Connecticut.....	138	148	178	205	225	236	240	208	174	114	155	146	2,167	Connecticut.
Florida.....	37	63	42	39	43	40	35	43	75	24	40	38	522	Florida.
Illinois.....	1,840	1,142	1,722	1,446	1,602	1,353	1,563	1,182	973	801	1,268	637	15,550	Illinois.
Louisiana.....	150	160	207	197	184	196	163	182	180	148	124	134	1,892	Louisiana.
Maine.....	22	30	9	21	34	41	19	23	23	25	19	19	296	Maine.
Massachusetts.....	706	753	898	865	966	818	696	664	619	465	570	589	8,590	Massachusetts.
Michigan.....	130	442	330	230	148	667	360	256	302	259	259	204	3,655	Michigan.
Minnesota.....	386	282	353	428	290	346	332	309	138	268	271	370	2,700	Minnesota.
Mississippi.....	435	540	678	637	614	610	608	468	369	344	310	247	5,860	Mississippi.
Montana.....	19	16	23	37	18	70	24	37	64	10	10	10	190	Montana.
New Jersey.....	518	602	819	826	731	624	724	666	594	509	587	766	7,946	New Jersey.
North Carolina.....	286	361	350	227	307	589	283	441	181	181	183	131	3,521	North Carolina.
Ohio.....	569	643	647	628	656	655	565	579	456	320	274	372	6,408	Ohio.
Oregon.....	38	57	79	70	78	91	78	102	81	46	42	42	793	Oregon.
Pennsylvania.....	852	98	1,077	1,244	1,005	1,186	859	903	780	361	251	629	10,114	Pennsylvania.
Porto Rico.....	70	85	118	138	174	150	217	165	186	119	114	140	1,698	Porto Rico.
Rhode Island.....	66	74	63	106	63	77	86	72	68	19	19	38	792	Rhode Island.
South Carolina.....	11	35	23	32	40	28	24	36	20	10	10	17	298	South Carolina.
South Dakota.....	17	16	17	35	27	15	23	28	28	20	10	37	443	South Dakota.
Texas.....	127	148	198	215	134	162	106	113	214	76	49	48	1,590	Texas.
Virginia.....	221	267	336	337	252	214	212	207	158	174	181	129	2,638	Virginia.
Washington.....	62	95	106	137	160	103	102	102	102	1	1	79	1,172	Washington.
Wyoming.....	1	4	Wyoming.

Deaths registered as due to tuberculosis (all forms), 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	217	248	261	265	252	231	211	206	164	203	172	182	2,612	Alabama.
Arizona.....	103	90	122	97	59	50	69	62	116	76	146	146	1,116	Arizona.
Arkansas.....	98	65	117	114	122	117	97	68	66	112	94	191	1,289	Arkansas.
California.....	435	568	472	568	441	424	426	419	533	500	563	561	5,981	California.
Connecticut.....	201	158	188	199	192	145	131	126	139	217	156	184	2,036	Connecticut.
Delaware.....	47	42	45	34	42	34	22	38	37	46	20	33	440	Delaware.
District of Columbia.....	70	54	83	66	66	66	62	70	47	69	47	50	739	District of Columbia.
Florida.....	119	82	94	79	98	85	87	73	76	97	100	91	1,081	Florida.
Hawaii.....	51	30	35	36	41	53	45	35	36	34	42	473	473	Hawaii.
Illinois.....	686	656	890	939	798	682	597	567	566	740	703	696	8,520	Illinois.
Indiana.....	323	420	461	361	336	296	276	276	247	280	289	316	3,944	Indiana.
Iowa.....	150	132	132	167	139	140	121	98	112	108	138	135	1,563	Iowa.
Kansas.....	99	73	108	99	88	74	73	71	65	92	83	108	1,038	Kansas.
Kentucky.....	401	425	513	479	418	409	297	333	280	397	341	286	4,591	Kentucky.
Louisiana.....	295	257	302	342	298	285	266	227	218	308	241	238	3,278	Louisiana.
Maine.....	72	91	87	92	86	81	81	68	64	97	57	73	921	Maine.
Massachusetts.....	460	470	510	605	490	461	427	393	451	528	462	501	5,738	Massachusetts.
Michigan.....	278	313	351	380	330	279	260	252	210	288	235	286	3,463	Michigan.
Minnesota.....	199	188	219	260	254	183	167	151	166	173	229	219	2,408	Minnesota.
Nebraska.....	50	61	52	52	37	67	38	38	35	49	43	52	594	Nebraska.
New Jersey.....	392	388	522	475	426	347	347	342	316	485	351	479	4,738	New Jersey.
New York.....	1,552	1,448	1,853	1,631	1,468	1,335	1,188	1,158	1,058	1,698	1,332	1,259	16,693	New York.
North Dakota.....	19	26	32	32	31	21	22	16	20	21	35	35	239	North Dakota.
Ohio.....	685	632	837	809	728	607	549	511	477	610	698	636	7,649	Ohio.
Oklahoma.....	53	54	95	99	130	80	93	90	64	92	81	75	1,001	Oklahoma.
Oregon.....	99	1,061	1,287	1,316	1,089	947	893	905	25	56	56	39	638	Oregon.
Pennsylvania.....	214	221	222	210	211	115	115	267	269	210	262	270	12,875	Pennsylvania.
Porto Rico.....	95	87	106	112	86	77	73	87	80	117	88	85	2,683	Porto Rico.
Rhode Island.....	112	144	176	172	191	151	121	131	160	131	149	149	1,083	Rhode Island.
South Carolina.....	17	16	17	17	35	27	15	23	28	20	37	44	1,757	South Carolina.
South Dakota.....	439	417	464	502	473	412	380	103	278	315	345	323	4,456	South Dakota.
Texas.....	15	15	24	24	10	15	16	17	12	23	19	17	207	Texas.
Utah.....	33	24	31	41	36	25	22	31	34	26	25	25	371	Utah.
Vermont.....	106	122	140	114	152	107	117	80	84	108	114	114	1,364	Vermont.
Washington.....	150	182	224	268	238	180	203	186	188	173	147	147	2,337	Washington.
Wisconsin.....	6	4	9	4	3	6	5	6	7	6	6	6	61	Wyoming.

Tuberculosis (all forms)—Average number of cases reported per annum and average indicated morbidity rates, 1916-17; cases reported, deaths registered, indicated morbidity and mortality rates, and number of cases reported for each death registered, 1918.

State.	Estimated population, July 1, 1918.	Average.			1918			State.
		Years.	Cases reported per annum.	Cases per 1,000 inhabitants.	Total cases reported.	Cases per 1,000 inhabitants.	Total deaths registered.	
Alabama.....	2,385,270	1917-1 year.....	597	1.2	2,870	1.2	2,612	1.1
Arizona.....	272,084	1917-1 year.....	6,988	2.3	864	1.115	4,115	4.1
Arkansas.....	1,792,965	1916, 1917-2 years.....	6,988	2.3	7,534	2.4	1,289	.7
California.....	3,116,412	1916, 1917-2 years.....	6,988	2.3	6,629	2.4	5,861	1.9
Colorado.....	1,014,588	1917-1 year.....	1,785	1.4	2,167	1.7	2,036	1.1
Connecticut.....	1,286,268	1916, 1917-2 years.....	1,082	2.8	—	—	440	1.6
Delaware.....	216,941	1916, 1917-2 years.....	—	—	—	—	739	1.8
District of Columbia.....	401,681	1916, 1917-2 years.....	—	—	—	—	1,081	1.2
Florida.....	938,877	1916, 1917-2 years.....	954	3.6	522	—	473	2.1
Hawaii.....	223,419	1916, 1917-2 years.....	—	—	—	—	—	—
Illinois.....	6,317,734	1917-1 year.....	13,999	2.2	15,559	2.5	8,950	1.8
Indiana.....	2,854,167	1917-1 year.....	3,875	1.4	—	—	3,944	1.4
Iowa.....	2,224,771	1916, 1917-2 years.....	—	—	—	—	1,583	1.7
Kansas.....	1,874,196	1916, 1917-2 years.....	1,445	.8	—	—	1,088	.6
Kentucky.....	2,404,547	1917-1 year.....	—	—	—	—	4,501	1.9
Louisiana.....	1,864,773	1917-1 year.....	1,870	1.0	1,902	—	3,278	1.7
Maine.....	752,191	1916, 1917-2 years.....	5,533	6	2,906	—	921	1.2
Massachusetts.....	3,832,190	1916, 1917-2 years.....	8,839	2.4	8,589	2.2	5,738	1.5
Michigan.....	3,133,678	1917-1 year.....	5,968	1.9	3,655	1.2	3,463	1.1
Minnesota.....	2,345,287	1917-1 year.....	4,907	2.1	3,700	1.6	2,408	1.5
Mississippi.....	2,001,486	1917-1 year.....	6,852	3.5	5,860	2.9	2,798	1.4
Missouri.....	3,448,496	1917-1 year.....	—	—	—	—	4,587	1.3
Montana.....	456,376	1916, 1917-2 years.....	—	—	599	1.0	—	—
Nebraska.....	1,266,877	1916, 1917-2 years.....	8,597	2.9	7,948	2.6	4,730	1.5
New Jersey.....	3,030,371	1916, 1917-2 years.....	29,613	2.9	—	—	16,933	1.6
New York.....	10,646,980	1916, 1917-2 years.....	—	—	3,521	1.4	—	—
North Carolina.....	2,488,025	1917-1 year.....	—	—	—	—	—	—
North Dakota.....	701,437	1916, 1917-2 years.....	168	.3	—	—	269	4
Ohio.....	5,223,914	1916, 1917-2 years.....	6,225	1.2	6,403	—	7,649	1.5
Oklahoma.....	2,377,029	1917-1 year.....	561	.2	—	—	1,001	.4
Oregon.....	388,243	1916, 1917-2 years.....	565	.7	793	.9	638	.7
Pennsylvania.....	8,768,067	1917-1 year.....	11,457	1.3	10,115	1.8	12,875	1.2
Puerto Rico.....	1,247,677	1917-1 year.....	951	.8	1,986	—	2,693	2.2
Rhode Island.....	637,415	1916, 1917-2 years.....	903	1.5	792	1.0	1,093	1.7

August 15, 1919.

South Carolina.....	1,860,934	1916—1 year.....	251	.2	286	1,757	1.1
South Dakota.....	735,434	1917—1 year.....	157	.2	307	307	.4
Texas.....	4,801,279	1917—1 year.....	1,239	.3	1,590	4,456	1.0
Utah.....	453,648	do.....				207	.5
Vermont.....	386,182					371	1.0
Virginia.....	2,234,030	1916, 1917—2 years.....	3,370	1.5	2,638	1.2	
Washington.....	1,860,578	1917—1 year.....	374	.3	1,172	1,364	.8
West Virginia.....	1,439,165	1917—1 year.....	1,800	.7			
Wisconsin.....	2,553,983	1916, 1917—2 years.....	56	.3	4	2,337	.9
Wyoming.....	190,380	1916—1 year.....				.61	.3

Typhoid fever—Monthly averages of cases reported during the years for which data are available, 1913-1917.

State.	Years.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Average per year.	State.	
Alabama.....	1913, 1915-1917—4 years.	81	67	50	65	117	310	575	532	339	265	152	89	2,642	Alabama.	
Arizona.....	1916—1 year.....	0	0	3	2	0	9	6	7	5	14	5	0	51	Arizona.	
Arkansas.....	1915—1 year.....	22	49	23	25	56	49	133	128	122	87	48	41	774	Arkansas.	
California.....	1913-1917—5 years.	100	70	76	89	95	105	165	185	150	169	101	104	1,498	California.	
Colorado.....	1914, 1916, 1917—5 years.	24	8	10	20	16	23	54	80	105	114	57	23	634	Colorado.	
Connecticut.....	1913-1917—5 years.	22	22	22	27	29	41	56	144	166	95	94	35	443	Connecticut.	
District of Columbia.....	do.....	15	12	12	13	14	24	37	72	68	47	32	15	361	District of Columbia.	
Hawaii.....	1913—1 year.....	11	17	17	16	15	16	13	11	11	33	23	20	213	Hawaii.	
Idaho.....	1913—1 year.....	0	0	1	0	0	10	12	26	56	23	30	18	176	Idaho.	
Illinois.....	1913-1917—3 years.....	239	171	140	184	136	145	324	512	686	508	427	209	3,741	Illinois.	
Indiana.....	1913-1917—5 years.	113	99	107	101	82	97	219	532	472	345	221	135	2,523	Indiana.	
Kansas.....	1914-1917—4 years.	37	34	41	38	39	51	212	351	335	322	214	125	1,798	Kansas.	
Louisiana.....	1916, 1917—2 years.	46	32	21	41	69	125	160	124	84	107	80	120	1,039	Louisiana.	
Maine.....	1915-1917—3 years.	19	11	6	10	11	7	19	32	47	27	27	22	242	Maine.	
Maryland.....	1913, 1916, 1917—3 years.....	107	71	71	75	79	97	122	122	645	903	407	322	156	Maryland.	
Massachusetts.....	1912-1917—5 years.	102	73	69	92	76	92	138	138	298	364	283	153	98	Massachusetts.	
Michigan.....	1912-1917—5 years.	124	124	113	114	91	119	132	174	322	258	154	104	1,956	Michigan.	
Minnesota.....	1913	73	63	54	45	58	84	165	188	177	106	106	106	1,160	Minnesota.	
Mississippi.....	1914-1917—4 years.	277	193	164	157	299	616	955	1,042	816	642	316	205	5,928	Mississippi.	
Montana.....	1913, 1914, 1916, 1917—4 years.....	51	34	26	36	35	52	99	104	36	35	49	25	13	492	Montana.
Nevada.....	1913, 1915, 1917—3 years.....	7	2	4	4	5	9	11	21	26	22	11	8	130	Nevada.	
New Jersey.....	1912-1917—5 years.	67	47	56	57	75	72	130	237	208	221	124	71	1,455	New Jersey.	
New York.....	do.....	262	244	280	284	277	272	393	601	971	771	438	344	5,115	New York.	
Ohio.....	1913-1917—4 years.....	8	5	7	26	24	7	20	34	39	23	14	22	231	Ohio.	
Oklahoma.....	1913, 1914, 1915, 1917—3 years.....	75	55	29	64	48	133	283	312	264	232	154	98	1,174	Oklahoma.	
Pennsylvania.....	1913-1917—5 years.	13	15	19	10	11	12	19	36	35	49	49	25	1,257	Pennsylvania.	
Porto Rico.....	1916, 1917—2 years.	378	317	317	287	292	382	506	920	1,536	1,278	427	221	6,880	Porto Rico.	
Rhode Island.....	1914, 1915, 1917—3 years.....	30	16	38	34	32	25	30	30	39	39	26	18	23	Rhode Island.	
South Carolina.....	1913-1917—5 years.	7	8	11	11	15	19	38	37	44	44	22	17	244	South Carolina.	
South Dakota.....	1913, 1914, 1917—3 years.....	24	14	23	26	62	194	203	149	147	73	42	19	976	South Dakota.	
Texas.....	1915, 1917—2 years.	37	32	24	41	38	90	158	158	138	145	145	129	1,147	Texas.	
Utah.....	1915-1916, 1917—4 years.....	13	11	20	22	21	39	65	82	131	104	78	26	621	Utah.	
Vermont.....	1914-1917—3 years.....	14	14	16	20	9	8	10	28	39	38	18	12	227	Vermont.	
Virginia.....	1912-1917—5 years.	145	133	95	110	160	345	667	1,024	728	486	321	167	4,361	Virginia.	
Washington.....	do.....	48	30	34	30	35	42	64	64	120	137	117	76	46	776	Washington.
West Virginia.....	1913-1917—5 years.....	71	74	17	63	81	106	160	339	286	165	120	45	1,629	West Virginia.	
Wisconsin.....	1913-1917—5 years.....	42	55	62	58	38	88	146	54	74	52	45	6	67	Wisconsin.	
Wyoming.....	do.....	5	4	2	5	3	4	8	11	9	6	6	6	6	Wyoming.	

1895

August 15, 1919.

Cases of typhoid fever reported, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	44	38	53	100	262	577	432	333	95	50	36		2,094	Alabama.
Arizona.....	7	2	1	5	9	24	5	10	1	0	6		71	Arizona.
Arkansas.....	10	37	20	15	17	42	58	84	76	25	27		473	Arkansas.
California.....	81	60	66	118	106	170	122	109	105	25	25		1,049	California.
Colorado.....	5	3	11	23	8	11	52	110	129	41	7		404	Colorado.
Connecticut.....	18	9	11	16	15	24	38	104	147	30	12		443	Connecticut.
District of Columbia.....	10	6	5	6	7	21	46	39	43	19	5		217	District of Columbia.
Florida.....	17	14	27	23	73	38	98	68	56	31	24		455	Florida.
Illinois.....	89	82	55	52	51	64	212	241	286	102	24		344	Illinois.
Indiana.....	48	42	87	45	64	74	94	187	207	150	82		1,127	Indiana.
Kansas.....	35	30	88	63	68	92	322	392	444	201	88		1,811	Kansas.
Louisiana.....	26	42	66	86	189	321	382	342	198	108	63		1,922	Louisiana.
Maine.....	10	5	6	4	7	13	10	17	17	12	4		112	Maine.
Maryland.....	39	30	75	70	63	154	351	435	222	86	101		1,701	Maryland.
Massachusetts.....	53	39	41	52	69	64	112	184	97	49	44		1,065	Massachusetts.
Michigan.....	44	47	71	47	35	69	66	115	226	128	33		915	Michigan.
Minnesota.....	44	59	82	61	34	22	83	77	78	18	34		614	Minnesota.
Mississippi.....	146	141	165	154	242	601	1,114	1,74	709	305	265		1,165	Mississippi.
Montana.....	2	12	23	3	7	28	31	37	35	11	10		210	Montana.
Nebraska.....	4	3	3	6	9	16	69	30	11	5		156	Nebraska.
New Jersey.....	52	34	46	36	38	65	132	166	64	25	45		739	New Jersey.
New York.....	122	123	168	197	186	250	548	586	461	202	180		3,390	New York.
North Carolina.....	20	17	30	31	46	280	879	949	769	233	75		3,388	North Carolina.
North Dakota.....	15	5	8	4	6	2	21	23	21	4	4		124	North Dakota.
Ohio.....	142	130	138	126	122	192	253	443	841	324	91		2,918	Ohio.
Oregon.....	14	4	13	10	7	12	15	26	24	35	16		183	Oregon.
Pennsylvania.....	146	201	328	288	220	291	316	643	976	660	185		4,559	Pennsylvania.
Puerto Rico.....	13	25	14	39	49	24	37	28	16	14	8		176	Puerto Rico.
Rhode Island.....	4	8	2	23	8	6	14	16	36	4	8		125	Rhode Island.
South Carolina.....	5	8	10	11	29	75	138	289	97	52	23		763	South Carolina.
South Dakota.....	1	1	9	3	5	2	3	16	23	13	3		92	South Dakota.
Texas.....	61	45	29	34	114	302	356	265	264	106	38		14	Texas.
Utah.....	10	8	11	10	21	19	27	37	37	34	9		18	Utah.
Vermont.....	14	8	2	1	14	6	5	31	32	21	6		143	Vermont.
Virginia.....	56	57	65	114	123	186	315	517	355	201	96		2,141	Virginia.
Washington.....	38	35	46	25	41	90	165	88	88	23	7		694	Washington.
West Virginia.....	48	42	96	70	104	177	204	192	69	61	18		1,127	West Virginia.
Wisconsin.....	22	31	56	42	36	35	29	43	43	19	4		368	Wisconsin.
Wyoming.....	1	2	2	3	23	22	9	3	3	3	1		66	Wyoming.

August 15, 1919.

1896

Deaths registered as due to typhoid fever, 1918.

State.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.	State.
Alabama.....	26	8	20	19	33	58	118	114	89	57	33	23	598	Alabama.
Arizona.....	4	3	5	5	3	10	9	4	2	8	5	2	56	Arizona.
Arkansas.....	11	16	5	7	15	19	26	27	36	13	49	49	243	Arkansas.
California.....	9	18	13	15	16	17	19	24	17	28	8	10	194	California.
Connecticut.....	5	3	4	4	2	4	4	9	17	17	2	4	75	Connecticut.
Delaware.....	0	0	3	3	1	1	1	12	13	18	2	5	59	Delaware.
District of Columbia.....	4	2	1	1	1	1	1	12	5	7	3	5	48	District of Columbia.
Florida.....	9	5	12	26	33	38	26	26	27	19	13	13	250	Florida.
Hawaii.....	3	6	6	2	6	1	8	4	1	3	0	5	45	Hawaii.
Illinois.....	26	37	24	21	33	27	37	84	75	86	44	39	533	Illinois.
Indiana.....	19	16	24	26	20	23	23	46	48	45	29	39	396	Indiana.
Iowa.....	9	9	12	9	7	13	16	18	17	31	4	9	154	Iowa.
Kansas.....	20	11	11	8	12	13	32	46	37	51	29	14	284	Kansas.
Kentucky.....	29	16	23	33	29	43	56	104	112	88	63	27	618	Kentucky.
Louisiana.....	21	26	28	31	59	104	110	118	70	72	34	18	691	Louisiana.
Maine.....	4	5	4	4	2	7	2	6	8	11	3	2	58	Maine.
Maryland.....	10	9	7	13	8	10	14	32	42	65	10	17	237	Maryland.
Massachusetts.....	9	8	5	8	9	13	11	17	26	36	9	7	158	Massachusetts.
Michigan.....	22	15	20	35	21	16	18	34	30	64	29	17	321	Michigan.
Minnesota.....	5	4	23	10	5	1	5	9	11	7	5	3	88	Minnesota.
Nebraska.....	3	3	4	4	4	5	8	5	9	9	1	1	58	Nebraska.
New Jersey.....	13	4	14	16	4	7	8	23	18	41	4	10	162	New Jersey.
New York.....	27	21	34	28	44	30	50	55	78	145	31	33	576	New York.
North Dakota.....	47	2	2	1	1	1	1	3	3	9	2	1	26	North Dakota.
Ohio.....	38	42	52	41	34	38	76	113	181	68	41	771	Ohio.	
Oklahoma.....	14	16	11	14	11	24	82	91	67	61	30	30	468	Oklahoma.
Oregon.....	2	2	6	3	5	4	9	9	5	5	16	6	6	Oregon.
Pennsylvania.....	45	52	68	68	62	43	76	87	129	108	62	44	924	Pennsylvania.
Porto Rico.....	17	21	15	20	9	8	15	13	14	15	3	7	157	Porto Rico.
Rhode Island.....	1	3	3	1	1	2	1	4	6	6	7	4	35	Rhode Island.
South Carolina.....	2	5	11	14	18	57	70	98	90	66	28	33	482	South Carolina.
South Dakota.....	47	46	31	67	136	176	148	113	117	71	61	3	37	South Dakota.
Texas.....	65	53	1	2	2	3	4	2	3	3	2	2	2	Texas.
Utah.....	3	1	2	2	2	1	2	1	4	7	4	1	31	Utah.
Vermont.....	2	6	2	2	2	2	2	1	1	18	17	10	98	Vermont.
Washington.....	4	10	18	12	5	5	4	11	4	10	7	3	94	Washington.
Wisconsin.....	1	2	2	2	2	2	2	6	6	10	7	1	18	Wyoming.

Typhoid fever—Average number of cases reported per annum and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated mortality, morbidity, and fatality rates, 1918.

State.	Estimated population July 1, 1918.	Average.		1918.		State.	
		Years.	Cases reported per annum.	Total cases reported.	Cases per 1,000 inhabitants.		
Alabama.....	2,385,270	1913,1915-1917—4 years.....	2,642	1.1	2,094	0.9	Alabama.
Arizona.....	272,034	1916—1 year.	51	.2	71	.3	Arizona.
Arkansas.....	1,792,965	1917—1 year.	774	.4	473	.3	Arkansas.
California.....	3,119,412	1913-1917—5 years.....	1,409	.5	1,049	.3	California.
Colorado.....	1,014,581	1914,1916-1917—3 years.....	534	.6	404	.4	Colorado.
Connecticut.....	1,398,288	1913-1917—5 years.....	733	.6	443	.3	Connecticut.
Delaware.....	216,941	1913-1917—5 years.....	361	1.0	217	.5	Delaware.
District of Columbia.....	401,681	1913-1917—5 years.....	485	.5	250	.5	District of Columbia.
Florida.....	938,877	1913-1917—5 years.....	213	1.0	45	.12	Florida.
Hawaii.....	223,410	1913-1917—1 year.	176	.5	250	.27	Hawaii.
Illinois.....	6,317,734	1913,1914,1916-1917—3 years.....	3,741	.6	1,344	.2	Illinois.
Indiana.....	2,884,167	1913-1917—5 years.....	2,523	.9	1,127	.4	Indiana.
Iowa.....	461,786	1913-1917—5 years.....	170	.5	127	.07	Iowa.
Kansas.....	1,874,195	1913-1917—5 years.....	1,790	1.0	811	1.0	Kansas.
Kentucky.....	2,408,547	1914-1917—4 years.....	1,009	.6	922	1.0	Kentucky.
Louisiana.....	1,884,775	1916,1917—2 years.....	2,242	.3	1,701	1.2	Louisiana.
Maine.....	1,782,191	1916,1917—3 years.....	2,788	2.0	1,701	1.2	Maine.
Maryland.....	1,384,539	1915-1917—3 years.....	1,919	.6	1,065	.3	Maryland.
Massachusetts.....	3,822,790	1913,1915-1917—5 years.....	1,936	.6	915	.3	Massachusetts.
Michigan.....	3,133,678	1913-1917—5 years.....	1,150	.5	614	.3	Michigan.
Minnesota.....	2,345,287	do.....	5,926	3.1	5,141	2.6	Minnesota.
Mississippi.....	2,001,466	1914-1917—4 years.....	492	1.1	210	.4	Mississippi.
Missouri.....	3,448,498	1913,1914,1916,1917—4 years.....	1,266,877	1.1	156	.1	Missouri.
Montana.....	466,376	1913-1917—3 years.....	114,742	1.3	130	.1	Montana.
Nebraska.....	3,080,371	1913-1917—5 years.....	1,455	.5	739	.2	Nebraska.
New Jersey.....	10,666,989	do.....	5,175	5	3,330	3	New Jersey.
New York.....	2,468,025	1917—1 year	2,701,437	3	3,388	1.4	New York.
North Carolina.....	5,233,814	1914-1917—4 years.....	231	.3	124	.2	North Carolina.
Ohio.....	5,233,814	1913,1915-1917—3 years.....	4,174	.8	2,918	.6	Ohio.
Oklahoma.....	2,377,629	1913-1917—5 years.....	1,757	.8	183	.2	Oklahoma.
Oregon.....	888,243	1913-1917—2 years.....	6,880	.8	4,559	.5	Oregon.
Pennsylvania.....	8,798,067	1916,1917—2 years.....	924	.11	70	.08	Pennsylvania.

Typhoid fever—Average number of cases reported per annum, and average indicated morbidity rates, 1913-1917; cases reported, deaths registered, and indicated mortality, morbidity, and fatality rates, 1918—Continued.

State.	Estimated population July 1, 1918.	Average.			1918			State.
		Years.	Cases reported per annum.	Cases per 1,000 inhabitants.	Total cases reported.	Deaths per 1,000 inhabitants.	Fatalities per 100 cases.	
Porto Rico.....	1,247,677	1914, 1915, 1917-3 years.....	345	0.3	275	0.2	157	Porto Rico.
Rhode Island.....	637,415	1914, 1916, 1917-3 years.....	244	.4	132	.2	35	Rhode Island.
South Carolina.....	1,689,934	1912, 1913, 1914, 1917-5 years.....	978	.6	783	.5	482	South Carolina.
South Dakota.....	735,434	1913, 1914, 1917-3 years.....	129	.2	92	.1	37	South Dakota.
Texas.....	4,601,279	1915, 1917-2 years.....	1,	1.3	1,621	1.4	1,057	Texas.
Utah.....	453,618	1913-1915, 1917-4 years.....	621	1.5	231	.5	23	Utah.
Vermont.....	386,192	1914-1917-5 years.....	227	.6	143	.4	31	Vermont.
Virginia.....	2,234,030	1913-1917-5 years.....	4,391	2.0	2,141	1.0	141	Virginia.
Washington.....	1,680,578	1917-1 do.....	778	.5	694	.4	98	Washington.
West Virginia.....	1,459,165	1913-1917-5 years.....	1,529	1.1	1,127	.8	94	West Virginia.
Wisconsin.....	2,533,983	1913-1917-5 years.....	623	.3	388	.1	18	Wisconsin.
Wyoming.....	190,380	do.....	67	.4	66	.3	18	Wyoming.

Cases of typhus fever reported, 1918.

1899

August 15, 1912.

Deaths registered as due to typhus fever, 1918.

Typhus fever—Cases reported, deaths registered, and indicated morbidity, mortality, and fatality rates, 1918.

State.	Estimated population July 1, 1918.	Total cases reported.	Cases per 1,000 inhabitants.	Deaths per 1,000 inhabitants.	Fatalities per 100 cases.	State.	Estimated population July 1, 1918.	Total cases reported.	Cases per 1,000 inhabitants.	Deaths per 1,000 inhabitants.	Fatalities per 100 cases.	
Connecticut.....	1,286,268	1	0.0098	1	0.009	Massachusetts.....	3,832,790	3,869,371	2	0.0005	1	0.0003
Colorado.....	1,014,591	1	.0011	1	.0011	New Jersey.....	3,869,371	3,869,371	1	.0001	1	.0003
Florida.....	938,877	1	.0029	3	.0022	Pennsylvania.....	8,798,067	4,601,279	11	.0023	1	.0002
Maryland.....	1,384,530	4				Texas.....						9.1

August 15, 1919.

1900

DEATHS DURING WEEK ENDED AUG. 2, 1919, IN CITIES.

From the "Weekly Health Index," Aug. 5, 1919, issued by the Bureau of the Census, Department of Commerce.

Deaths from all causes in certain large cities of the United States during the week ended Aug. 2, 1919, infant mortality (per cent), annual death rates, and comparison with corresponding week of preceding years.

City.	Population July 1, 1918, esti- mated.	Week ended Aug. 2, 1919.		Average annual death rate per 1,000. ¹	Per cent of deaths under 1 year.	
		Total deaths.	Death rate. ¹		Week ended Aug. 2, 1919.	Previous year or years. ²
Albany, N. Y.	112,565	22	10.2	C 17.6	4.5	C 5.3
Atlanta, Ga.	201,732	54	14.0	C 12.7	5.6	C 14.3
Baltimore, Md.	3,660,981	211	16.4	A 18.9	26.1	A 28.0
Boston, Mass.	785,245	173	11.5	A 14.4	17.9	A 17.5
Buffalo, N. Y.	473,229	119	13.1	C 15.6	26.9	C 33.8
Cambridge, Mass.	111,432	20	9.4	A 10.9	5.0	A 26.3
Chicago, Ill.	2,596,681	553	11.1	A 13.7	23.0	A 26.7
Cincinnati, Ohio	418,022	111	13.8	C 10.9	16.2	C 17.2
Cleveland, Ohio	810,306	170	10.9	C 11.3	14.7	C 31.8
Columbus, Ohio	225,296	57	13.2	C 12.7	12.3	C 14.5
Dayton, Ohio	130,655	34	13.6	C 12.4	17.6	C 6.5
Denver, Colo.		55			16.4	
Fall River, Mass.	128,392	33	13.4	C 19.9	39.4	C 51.0
Grand Rapids, Mich.	135,450	14	5.4	C 10.8	14.3	C 10.7
Indianapolis, Ind.	290,389	85	15.3	C 13.7	9.4	C 18.4
Jersey City, N. J.	318,770	73	11.9	C 9.8	12.3	C 30.0
Kansas City, Mo.	313,785	102	16.9	C 16.5	16.7	C 26.3
Los Angeles, Calif.	568,495	104	9.5	A 12.3	13.5	A 10.0
Louisville, Ky.	242,707	75	16.1	C 14.4	10.7	C 4.5
Lowell, Mass.	109,081	27	12.9	A 18.2	59.3	A 35.1
Memphis, Tenn.	154,759	58	19.5	C 17.9	25.9	C 11.3
Milwaukee, Wis.	453,481	95	10.9	A 12.1	31.6	A 16.5
Minneapolis, Minn.	383,442	72	9.8	C 8.7	11.1	C 6.3
Nashville, Tenn.	119,215	50	21.9	C 19.2	16.0	C 13.6
Newark, N. J.	428,684	99	12.0	C 13.0	31.3	C 23.4
New Haven, Conn.	154,865	33	11.1	C 12.5	21.2	C 16.2
New Orleans, La.	382,273	126	17.2	A 18.1	14.3	A 14.2
New York, N. Y.	5,215,879	1,198	12.0	C 11.2	22.6	C 21.2
Oakland, Calif.	214,206	44	10.7	A 10.1	2.3	A 9.8
Omaha, Nebr.	180,264	34	9.8	C 11.3	23.5	C 15.4
Philadelphia, Pa.	1,761,371	429	12.7	A 17.1	22.1	A 23.9
Pittsburgh, Pa.	593,303	128	11.2	C 14.9	23.4	C 27.8
Portland, Oreg.		45			11.1	C 11.8
Providence, R. I.	263,613	49	9.7	C 12.7	24.5	C 32.8
Richmond, Va.	160,719	42	13.6	C 13.3	26.2	C 12.2
Rochester, N. Y.	264,856	50	9.8	C 15.6	2.0	C 20.3
St. Louis, Mo.	779,251	158	10.6	C 13.6	12.7	C 13.2
St. Paul, Minn.	257,699	38	7.7	C 9.5	13.2	C 12.8
Spokane, Wash.		24			4.2	C 11.1
Syracuse, N. Y.	161,404	28	9.0	C 17.8	14.3	C 29.1
Toledo, Ohio	262,234	55	10.9	A 14.4	5.5	A 20.8
Washington, D. C.	401,681	130	16.9	A 18.0	19.2	A 17.1
Worcester, Mass.	173,650	50	15.0	C 13.2	20.6	C 15.9

¹ Annual rates per 1,000 estimated population.² "A" indicates data for the corresponding week of the years 1913 to 1917, inclusive. "C" indicates data for the corresponding week of the year 1918.³ Population estimated as of July 1, 1919.⁴ Data are based on statistics of 1915, 1916, and 1917.

Summary of information received by telegraph from industrial insurance companies for week ended Aug. 2, 1919.

Policies in force	40,774,651
Number of death claims	6,846
Death claims per 1,000 policies in force, annual rate	8.8

PREVALENCE OF DISEASE.

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring.

UNITED STATES.

CURRENT STATE SUMMARIES.

Telegraphic Reports for Week Ended Aug. 9, 1919.

These reports are preliminary, and the figures are subject to change when later returns are received by the State health officers.

ALABAMA.		CONNECTICUT.	
	Cases.		Cases.
Diphtheria.....	14	Cerebrospinal meningitis:	
Malaria.....	32	Torrington.....	1
Poliomyelitis.....	1	Seymour.....	1
Scarlet fever.....	7	Poliomyelitis:	
Smallpox.....	2	Torrington.....	2
Tuberculosis.....	41		
Typhoid fever.....	38		
ARKANSAS.			
Chicken pox.....	9	Chancroid.....	2
Diphtheria.....	1	Chicken pox.....	1
Influenza.....	2	Diphtheria:	
Lethargic encephalitis.....	1	Wilmington.....	1
Malaria.....	250	Gonorrhea.....	36
Pellagra.....	12	Influenza:	
Poliomyelitis.....	1	Milton.....	1
Scarlet fever.....	2	Malaria:	
Trachoma.....	2	Dover.....	1
Tuberculosis.....	8	Measles.....	1
Typhoid fever.....	16	Mumps.....	3
Whooping cough.....	6	Scarlet fever:	
CALIFORNIA.			
Influenza.....	7	Townsend.....	1
Poliomyelitis:		Syphilis.....	3
Redwood City.....	1	Tuberculosis:	
Claremont.....	1	Wilmington.....	3
Smallpox:		New Castle.....	1
Oakland.....	2	Typhoid fever:	
Monterey County.....	1	Laurel.....	1
San Francisco.....	7	Harrington.....	1
Santa Clara County.....	2	McDonough.....	1
Los Angeles County.....	4	Hartly.....	1
San Joaquin County.....	1	Townsend.....	1
Stockton.....	1	Wyoming.....	3
Long Beach.....	1		
Santa Rosa.....	1		
Typhoid fever:			
Sacramento.....	2		
Fresno County.....	1		
Modoc County.....	1		
Woodland.....	1		
Los Angeles County.....	1		
Perris.....	1		
Riverside.....	1		
FLORIDA.			
Cerebrospinal meningitis:			
Orange County.....			1
Walton County.....			2
Diphtheria.....			1
Influenza.....			5
Malaria.....			61
Scarlet fever.....			3
Typhoid fever.....			5
GEORGIA.			
Actinomycosis.....			2
Acute infectious conjunctivitis.....			3
Cerebrospinal meningitis.....			1

August 15, 1919.

1902

GEORGIA—continued.	Cases.	ILLINOIS—continued.	Cases.
Chicken pox.....	4	Typhoid fever—Continued.	
Diphtheria.....	18	Elgin.....	4
Dysentery (amebic).....	3	Galesburg.....	2
Dysentery (bacillary).....	6	Bluff Precinct.....	10
Gonorrhea.....	258	Jacksonville.....	2
Hookworm.....	29	Moline.....	2
Influenza.....	2	Springfield.....	2
Malaria.....	128	Joliet.....	3
Mumps.....	4	Scattering.....	9
Pneumonia (acute lobar).....	13		
Scarlet fever.....	3	INDIANA.	
Septic sore throat.....	2	Diphtheria:	
Smallpox.....	9	Richmond.....	66
Syphilis.....	138	Johnson County.....	1
Tetanus.....	1	Montgomery County.....	1
Tuberculosis (pulmonary).....	14	Tipton County.....	1
Typhid fever.....	67	Ripley County.....	1
Whooping cough.....	8	Randolph County.....	2
		Harrison County.....	4
ILLINOIS.		Hendricks County.....	2
Cerebrospinal meningitis:		Grant County.....	7
Springfield.....	1	Decatur County.....	1
Standard.....	1	Blackford County.....	7
Diphtheria:		Lake County.....	1
Chicago.....	62	Washington County.....	2
Harvey.....	2	Kosciusko County.....	3
Lemont.....	2	Sullivan County.....	2
Dekalb.....	2	Boone County.....	2
Crescent City.....	3	Gonorrhea.....	119
Mooseheart.....	2	Measles:	
Livingston County—		Prevalent in—	
Owego Township.....	2	Clay County.	
Peoria.....	2	Owen County.	
Scattering.....	11	Lake County.	
Gonorrhea.....	282	Delaware County.	
Lethargic encephalitis:		Noble County.	
La Salle.....	1	Rabies in animals:	
Poliomyelitis:		Orange County.....	1
Chicago.....	14	Scarlet fever:	
La Grange.....	1	Prevalent in—	
Cumberland County—		Madison County.	
Spring Point Township.....	1	Clinton County.	
Kewanee.....	1	Elkhart County.	
Galesburg.....	1	Hendricks County.	
Knox County—		Jay County.	
Lynn Township.....	1	Delaware County.	
Wenona.....	3	Smallpox:	
Sigel.....	1	Prevalent in—	
Bradford.....	1	Wayne County.	
Freeport.....	1	Blackford County.	
Monmouth.....	2	Ripley County.	
Scarlet fever:		Elkhart County.	
Chicago.....	16	Syphilis.....	102
Springfield.....	6	Typhoid fever:	
Marion.....	2	Prevalent in—	
Scattering.....	4	Jennings County.	
Smallpox:		Harrison County.	
Hillview.....	2	Owen County.	
Momence.....	2	Delaware County.	
Scattering.....	4	Shelby County.	
Syphilis.....	190	Parke County.	
Typhoid fever:		Sullivan County.	
Chicago.....	7	IOWA.	
Edgewood.....	2	Chancroid.....	2
St. Elmo.....	2	Diphtheria:	
Sesser.....	2	Emmetsburg.....	1
		Cerro Gordo County.....	1

1903

August 15, 1919.

IOWA—continued.	Cases.	MASSACHUSETTS—continued.	Cases.
Gonorrhœa.....	32	Mumps.....	17
Scarlet fever:		Ophthalmia neonatorum.....	17
Burlington.....	1	Pellagra.....	1
Des Moines.....	2	Pneumonia (lobar).....	16
Dubuque.....	1	Scarlet fever.....	51
Butler County.....	1	Septic sore throat.....	10
Humboldt County.....	1	Syphilis.....	73
Smallpox:		Tetanus.....	1
Bennett.....	1	Typhoid fever.....	24
Boone.....	2	Tuberculosis.....	154
Syphilis.....	8	Whooping cough.....	112
KANSAS.			
Cerebrospinal meningitis:		MINNESOTA.	
Carlton.....	1	Cerebrospinal meningitis.....	4
Diphtheria.....	12	Chancroid.....	5
Scarlet fever.....	9	Gonorrhœa.....	107
Smallpox.....	18	Poliomyelitis.....	5
LOUISIANA.			
Chancroid.....	45	Smallpox:	
Diphtheria.....	9	Grant County—Lawrence Township.....	5
Gonorrhœa.....	212	Otter Tail County—	
Influenza.....	2	Astad Township.....	1
Pellagra.....	9	Buse Township.....	1
Smallpox.....	2	Traverse County—Croke Township.....	3
Syphilis.....	117	Wright County—Silver Creek Township....	1
Typhoid fever.....	31	Syphilis.....	108
MAINE.			
Chicken pox.....	1	MONTANA.	
Diphtheria:		Cerebrospinal meningitis:	
Portland.....	3	Absarokee.....	1
Friendship.....	1	Diphtheria.....	10
Winslow.....	1	Influenza.....	3
Gonorrhœa.....	30	Scarlet fever.....	3
Measles.....	1	Typhoid fever.....	14
Scarlet fever:		NEW JERSEY.	
Portland.....	3	Influenza.....	2
Westbrook.....	1	Pneumonia.....	15
Farmington.....	1	NEW YORK.	
Smallpox:		(Exclusive of New York City.)	
Boothbay.....	5	Cerebrospinal meningitis:	
Lewiston.....	5	Buffalo.....	1
Syphilis.....	20	Tonawanda.....	1
Tuberculosis.....	17	Diphtheria.....	135
Typhoid fever:		Gonorrhœa (voluntary reports).....	47
Harrison.....	1	Influenza.....	3
Portland.....	1	Measles.....	90
Calais.....	1	Poliomyelitis:	
Rumford.....	1	Rotterdam.....	1
Whooping cough:		Ithaca.....	1
Portland.....	2	Hague.....	1
Stonington.....	1	Newark.....	1
South Berwick.....	3	Yonkers.....	1
Islesboro.....	3	Pneumonia.....	33
Farmington.....	1	Scarlet fever.....	50
MASSACHUSETTS.		Syphilis (voluntary reports).....	170
Cerebrospinal meningitis.....	6	Typhoid fever:	
Chicken pox.....	29	Tonawanda.....	11
Diphtheria.....	76	Scattering.....	35
German measles.....	3	Whooping cough.....	175
Gonorrhœa.....	214	NORTH CAROLINA.	
Influenza.....	5	Chancroid.....	19
Malaria.....	1	Chicken pox.....	13
Measles.....	42	Cholera infantum.....	5
		Diphtheria.....	31
		Dysentery (amebic).....	1

August 15, 1919.

1904

NORTH CAROLINA—continued.		WEST VIRGINIA—continued.	
	Cases.		Cases.
Dysentery (bacillary).....	1	Measles: Charleston.....	2
Gonorrhea.....	121	Fairmont.....	1
Measles.....	26	Keyser.....	1
Ophthalmia neonatorum.....	1	Poliomyelitis: Parkersburg.....	1
Paratyphoid fever.....	2	Scarlet fever: Charleston.....	1
Pneumonia (broncho).....	3	Clarksburg.....	3
Pneumonia (lobar).....	3	Fairmont.....	1
Scarlet fever.....	24	Huntington.....	2
Septic sore throat.....	7	Smallpox: Morgantown.....	1
Smallpox.....	14	Wheeling.....	2
Syphilis.....	42	Typhoid fever: Beckley.....	1
Trachoma.....	1	Clarksburg.....	1
Typhoid fever.....	144	Huntington.....	14
Whooping cough.....	85	Keyser.....	1
OHIO.		Martinsburg.....	3
Smallpox:		Wellsburg.....	1
Uhrichsville.....	7	Weston.....	1
Youngstown.....	9	WISCONSIN.	
Typhoid fever:		Cerebrospinal meningitis:	
Struthers.....	3	Milwaukee.....	1
Warren.....	5	State.....	1
Cambridge.....	5	Chicken pox:	
VERMONT.		Milwaukee.....	2
No outbreak or unusual prevalence.		State.....	4
VIRGINIA.		Diphtheria:	
Poliomyelitis:		Milwaukee.....	18
Richmond.....	1	State.....	19
Henrico County.....	1	Measles:	
WASHINGTON.		Milwaukee.....	3
Chicken pox.....	23	State.....	9
Diphtheria.....	16	Poliomyelitis:	
Gonorrhea.....	29	Milwaukee.....	0
Measles.....	7	State.....	18
Mumps.....	14	Scarlet fever:	
Scarlet fever.....	34	Milwaukee.....	9
Smallpox.....	37	State.....	16
Syphilis.....	2	Smallpox:	
Tuberculosis.....	14	Milwaukee.....	3
Typhoid fever.....	12	State.....	20
Whooping cough.....	31	Tuberculosis.....	17
WEST VIRGINIA.		Typhoid fever.....	12
Diphtheria:		Whooping cough:	
Huntington.....	2	Milwaukee.....	45
Moundsville.....	1	State.....	8
Weston.....	2		

SUMMARY OF CASES REPORTED MONTHLY BY STATES.

Tables showing by counties the reported cases of cerebrospinal meningitis, malaria, pellagra, poliomyelitis, smallpox, and typhoid fever are published under the names of these diseases. (See names of these and other diseases in the table of contents.)

The following monthly State reports include only those which were received during the current week. These reports appear each week as received.

State.	Cer- ebro- spinal menin- gitis.	Diph- theria.	Malaria.	Measles.	Pel- lagra.	Poli- omy- elitis.	Scarlet fever.	Small- pox.	Ty- phoid fever.
June, 1919:									
Alabama.....	6	16	213	41	59	12	16	33	112
Indiana.....	3	94	-----	584	-----	1	160	277	42
Nebraska.....	1	27	-----	81	-----	-----	38	382	3
Virginia.....	5	93	430	458	47	20	62	128	267

1905

August 15, 1919.

ANTHRAX.

Wilmington, Del., and Woburn, Mass.

During the week ended July 26, 1919, one case of anthrax was reported at Wilmington, Del. On July 22, 1919, a case was reported at Woburn, Mass.

CEREBROSPINAL MENINGITIS.

State Reports for June, 1919.

Place.	New cases reported.	Place.	New cases reported.
Alabama:		Nebraska:	
Crenshaw County.....	1	Lancaster County.....	1
Houston County.....	1		
Lee County.....	1		
Mobile County.....	1		
Talladega County.....	1		
Wilcox County.....	1		
Total.....	6		
Indiana:		Virginia:	
Clark County.....	1	Buchanan County.....	1
Jay County.....	1	James City County—	1
Lake County.....	1	Williamsburg.....	1
Total.....	3	Mathews County.....	1
		Middlesex County.....	1
		Norfolk County.....	1
		Total.....	5

City Reports for Week Ended July 26, 1919.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Appleton, Wis.....	1	New York, N. Y.....	1	1
Benton Harbor, Mich.....	1	Norfolk, Va.....	1
Boston, Mass.....	1	Pawtucket, R. I.....	1
Bridgeport, Conn.....	1	Philadelphia, Pa.....	1
Chicago, Ill.....	1	Pittsfield, Mass.....	1
Cleveland, Ohio.....	1	Providence, R. I.....	1	1
Colorado Springs, Colo.....	1	1	Rahway, N. J.....	1
Louisville, Ky.....	1	Richmond, Va.....	1	1
Memphis, Tenn.....	1	1	San Francisco, Calif.....	1
Milwaukee, Wis.....	4	3			

DIPHTHERIA.

See Telegraphic weekly reports from States, p. 1901; Monthly summaries by States, p. 1904; and Weekly reports from cities, p. 1913.

LEPROSY.

San Francisco, Calif.

During the week ended July 26, 1919, one case of leprosy was reported at San Francisco, Calif.

August 15, 1919.

1906

MALARIA.**State Reports for June, 1919.**

Place.	New cases reported.	Place.	New cases reported.
Alabama:		Virginia—Continued.	
Autauga County.....	15	Halifax County.....	17
Bibb County.....	1	South Boston.....	11
Colbert County.....	2	Hanover County.....	21
Geneva County.....	1	Henrico County—	
Greene County.....	6	Richmond.....	5
Houston County.....	108	Isle of Wight County.....	26
Jackson County.....	1	King and Queen County.....	10
Jefferson County.....	8	Lancaster County.....	3
Lamar County.....	25	Loudoun County.....	6
Limestone County.....	1	Lunenburg County.....	9
Marengo County.....	1	Victoria.....	2
Marshall County.....	1	Mecklenburg County.....	32
Mobile County.....	2	Middlesex.....	9
Montgomery County.....	6	Nansemond County.....	13
Morgan County.....	1	Whaleyville.....	3
Shelby County.....	2	Suffolk.....	6
Sumter County.....	1	Northampton County.....	3
Talladega County.....	1	Cape Charles.....	11
Tuscaloosa County.....	24	Pittsylvania County.....	11
Walker County.....	6	Powhatan County.....	5
Total.....	213	Princess Anne County.....	9
		Oceania.....	11
Virginia:		Virginia Beach.....	2
Accomac County.....	1	Prince Edward County—	18
Alleghany County—		Farmville.....	
Clifton Forge.....	2	Prince George County—	
Amherst County—		Hopewell.....	
Woodstock.....	2	Prince William County—	
Appomattox County.....	13	Quantico.....	
Botetourt County.....	17	Roanoke County—	
Brunswick County—		Salem.....	
Lawrenceville.....	17	Shenandoah County.....	
Buckingham County.....	2	Southampton County.....	
Campbell County.....	3	Franklin.....	
Caroline County.....	13	Sebrell.....	
Charles City County.....	7	Stafford County—	
Chesterfield County—		Falmouth.....	
Winterpock.....	2	Surry County.....	
Dickenson County—		Sussex County.....	
Clinwood.....	1	Stony Creek.....	
Elizabeth City County—		Warwick County—	
Hampton.....	2	Camp Custis.....	
Essex County.....	5	Denigh.....	
Fluvanna County.....	2	Washington County.....	
Franklin County.....	3	Wise County.....	
Giles County.....	1	York County.....	
Greenville County—		Total.....	430
North Emporia.....	23		

City Reports for Week Ended July 26, 1919.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Baton Rouge, La.....	2	Newark, N. J.....	3
Belleview, N. J.....	1	New Orleans, La.....	3	1
Birmingham, Ala.....	1	1	North Little Rock, Ark.....	1
Charleston, S. C.....	1	Paterson, N. J.....	1
Chicago, Ill.....	1	Portsmouth, Va.....	1
Dallas, Tex.....	10	1	Richmond, Va.....	1
East Orange, N. J.....	1	Rome, Ga.....	6
Elgin, Ill.....	2	San Francisco, Calif.....	2
Fort Worth, Tex.....	1	Savannah, Ga.....	3
Little Rock, Ark.....	4	Toledo, Ohio.....	1
Memphis, Tenn.....	3	1	Tuscaloosa, Ala.....	3

MEASLES.

See Telegraphic weekly reports from States, p. 1901; Monthly summaries by States, p. 1904; and Weekly reports from cities, p. 1913.

1907.

August 15, 1919.

PELLAGRA.

State Reports for June, 1919.

Place.	New cases reported.	Place.	New cases reported.
Alabama:		Virginia—Continued.	
Colbert County.....	1	Hanover County.....	1
Conecuh County.....	1	Henrico County—	
Geneva County.....	2	Glen Allen.....	1
Jefferson County.....	7	Richmond.....	3
Mobile County.....	21	Henry County—	
Monroe County.....	2	Martinsville.....	1
Montgomery County.....	8	Isle of Wight County.....	1
Randolph County.....	1	Loudoun County—	
Talladega County.....	3	Leesburg.....	1
Tuscaloosa County.....	11	Mathews County.....	1
Walker County.....	2	Northampton County.....	4
Total.....	59	Patrick County.....	1
		Pittsylvania County—	
		Danville.....	1
		Powhatan County.....	1
Virginia:		Scott County.....	1
Alexandria County.....	1	Southampton County.....	1
Alleghany County—		Sebrell.....	1
Clifton Forge.....	1	Spotsylvania County—	
Augusta County.....	1	Fredericksburg.....	1
Buckingham County.....	1	Surry County.....	6
Chesterfield County—		Sussex County.....	1
Winterpock.....	1	Wise County.....	2
Dinwiddie County.....	1	Total.....	47
Elizabeth City County—			
Hampton.....	2		
Essex County.....	2		
Greenville County.....	6		
North Emporia.....	2		

City Reports for Week Ended July 26, 1919.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Austin, Tex.....		1	Los Angeles, Calif.....	1	
Birmingham, Ala.....		1	Macon, Ga.....		1
Charleston, S. C.....		1	Memphis, Tenn.....		1
Charlotte, N. C.....		1	Montgomery, Ala.....	1	
Dallas, Tex.....	8		Nashville, Tenn.....	1	1
Fort Worth, Tex.....		1	New Orleans, La.....	1	
Lexington, Ky.....		1	Tuscaloosa, Ala.....	2	
Little Rock, Ark.....	1				

PLAQUE-INFECTED GROUND SQUIRRELS.

Alameda County, Calif.

On July 17, 1919, a plague-infected ground squirrel was reported found in Alameda County, Calif. Another one was reported found on July 18, 1919. Diagnoses were based on animal inoculation and cultures. Intensive hunting and poisoning operations are being carried on.

August 15, 1919.

1908

PNEUMONIA.

City Reports for Week Ended July 26, 1919.

Place.	Lobar.		All forms.		Place.	Lobar.		All forms.	
	Cases.	Deaths.	Cases.	Deaths.		Cases.	Deaths.	Cases.	Deaths.
Anderson, Ind.		1			Martins Ferry, Ohio			1	
Arlington, Mass.	1	1			Memphis, Tenn.			1	
Atlanta, Ga.		3			Minneapolis, Minn.			2	
Baltimore, Md.	1	1			Newark, N. J.	3	4		
Birmingham, Ala.		2			New Bedford, Mass.	1	1		
Boston, Mass.	5	1			New Orleans, La.			1	
Bridgeport, Conn.		1			New York, N. Y.			9	62
Buffalo, N. Y.		1			Oakland, Calif.			2	
Cairo, Ill.		1			Oak Park, Ill.			1	
Charlotte, N. C.		1			Omaha, Nebr.				2
Chicago, Ill.		1	55	27	Orange, N. J.			1	
Cincinnati, Ohio		1			Oshkosh, Wis.			1	
Cleveland, Ohio	5	5			Paterson, N. J.	18		1	
Columbus, Ga.	2				Pawtucket, R. I.				
Dayton, Ohio	1	1			Philadelphia, Pa.	17	4		
Denver, Colo.		1		3	Portland, Oreg.				1
Detroit, Mich.	1	6	3	11	Portsmouth, Va.			1	
East Chicago, Ind.		6		1	Providence, R. I.			2	
Elizabeth, N. J.		1			Racine, Wis.			1	
Elmira, N. Y.				1	Richmond, Va.			3	
Everett, Mass.			1		Rochester, N. Y.	1	1		
Fort Worth, Tex.		4			Rockford, Ill.			1	
Fresno, Calif.		1			Rome, N. Y.	3			
Grand Rapids, Mich.	1				Salt Lake City, Utah			1	
Holyoke, Mass.	1	1			San Francisco, Calif.	5	7		
Jersey City, N. J.		2			Santa Cruz, Calif.	1			
Johnstown, N. Y.	1	1			Savannah, Ga.			1	
La Crosse, Wis.	1	1			Schenectady, N. Y.	2			
Lawrence, Kans.	1				Springfield, Ill.			1	
Lawrence, Mass.	1				Terre Haute, Ind.			1	
Lexington, Ky.		1			Topeka, Kans.	1			
Los Angeles, Calif.	1				Trenton, N. J.	1			
Lowell, Mass.	1				Troy, N. Y.			1	
Lynn, Mass.	1	1			Washington, D. C.			2	
Madison, Wis.		2			Wheeling, W. Va.				1
Malden, Mass.	1				Yonkers, N. Y.	1	1		
Mankato, Minn.			1						

POLIOMYELITIS (INFANTILE PARALYSIS).

State Reports for June, 1919.

Place.	New cases reported.	Place.	New cases reported.
Alabama:		Virginia—Continued.	
Jefferson County	8	Elizabeth City County—	
Pike County	1	Hampton	2
Tuscaloosa County	1	Phoebeus	
Walker County	2	Hanover County	2
Total.....	12	Henrico County—	
Indiana:		Richmond	1
Switzerland County.....	1	Lee County	1
Virginia:		Washington County.....	1
Culpeper County	2	Total.....	20
Dickenson County	4		
Clintwood.....	2		

1909

August 15, 1919.

POLIOMYELITIS (INFANTILE PARALYSIS)—Continued.**City Reports for Week Ended July 26, 1919.**

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Baltimore, Md.....	4	2	Marinette, Wis.....	1
Charleston, W. Va.....	2	2	Milwaukee, Wis.....	22	2
Chicago, Ill.....	8	1	Natick, Mass.....	1
Cleveland, Ohio.....	1	Omaha, Nebr.....	5	2
Flint, Mich.....	1	Pontiac, Mich.....	3	2
Fond du Lac, Wis.....	1	Richmond, Va.....	1
Janesville, Wis.....	1	Topeka, Kans.....	1
Kenosha, Wis.....	1			

RABIES IN ANIMALS.**City Reports for Week Ended July 26, 1919.**

During the week ended July 26, 1919, three cases of rabies in animals were reported at Akron, Ohio, and one at Chillicothe, Ohio.

SCARLET FEVER.

See Telegraphic weekly reports from States, p. 1901; Monthly summaries by States, p. 1904; and Weekly reports from cities, p. 1913.

SMALPOX.**State Reports for June, 1919.**

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Alabama:			Indiana—Continued.		
Chilton County.....	1	Sullivan County.....	1
Clarke County.....	2	Switzerland County.....	3
Colbert County.....	1	1	Tipppecanoe County.....	10
DeKalb County.....	1	Tipton County.....	3
Jefferson County.....	6	Vermillion County.....	4
Mobile County.....	18	1	Vigo County.....	1
Morgan County.....	2	Wells County.....	5
Talladega County.....	1	White County.....	5
Wilcox County.....	1	Total.....	277
Total.....	33	2			
Indiana:			Nebraska:		
Allen County.....	11	Adams County.....	2
Boone County.....	1	Antelope County.....	1
Cass County.....	53	Box Butte County.....	5
Dearborn County.....	14	Boyd County.....	1
Delaware County.....	5	Buffalo County.....	24
Elkhart County.....	4	Burt County.....	1
Fayette County.....	3	Cass County.....	3
Floyd County.....	1	Cherry County.....	2
Fountain County.....	1	Cheyenne County.....	1
Fulton County.....	2	Clay County.....	19
Grant County.....	6	Custer County.....	54
Hamilton County.....	7	Dixon County.....	1
Hendricks County.....	1	Douglas County.....	81
Henry County.....	9	Franklin County.....	2
Howard County.....	8	Furnas County.....	1
Huntington County.....	2	Gage County.....	15
Jennings County.....	3	Jefferson County.....	4
Johnson County.....	1	Keith County.....	1
Knox County.....	4	Lancaster County.....	117
Kosciusko County.....	1	Otoe County.....	1
Lake County.....	3	Phelps County.....	4
LaPorte County.....	14	Richardson County.....	23
Lawrence County.....	6	Saline County.....	1
Madison County.....	50	Saunders County.....	19
Marion County.....	9	Sheridan County.....	1
Ohio County.....	3	Stanton County.....	1
Parke County.....	2	Thomas County.....	1
Randolph County.....	2	Wayne County.....	1
Ripley County.....	10	York County.....	1
Shelby County.....	4	Total.....	382
St. Joseph County.....	5			

August 15, 1919.

1910

SMALLPOX—Continued.**State Reports for June, 1919—Continued.**

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Virginia:			Virginia—Continued.		
Accomac County.....	10		Nottoway County—		
Albemarle County—			Blacksome.....	2	
Charlottesville.....	2		Pittsylvania County—		
Alleghany County—			Danville.....	1	
Clifton Forge.....	2		Princess Anne County.....	1	
Covington.....	8		Pulaski County.....	1	
Bath County.....	10		Roanoke County.....	21	
Bedford County.....	1		Vinton.....	1	
Buchanan County.....	1		Rockbridge County—		
Campbell County.....	4		Glasgow.....	1	
Chesterfield County.....	7		Scott County.....	8	
Franklin County.....	1		Sussex County.....	2	
Henrico County—			Tazewell County—		
Richmond.....	1		Pocahontas.....	2	
Henry County.....	4		Warren County—		
Isle of Wight County.....	10		Front Royal.....	5	
Loudoun County.....	1		Warwick County—		
Louisa County—			Denbigh.....	1	
Louisa.....	3		Washington County.....	10	
Lunenburg County.....	3		Total.....	138	
Mathews County.....	2				
Montgomery County.....	1				
Norfolk County.....	9				
Portsmouth.....	2				

City Reports for Week Ended July 26, 1919.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Aberdeen, Wash.....	2		Lincoln, Nebr.....	2	
Anniston, Ala.....	1		Logansport, Ind.....	6	
Appleton, Wis.....	1		Los Angeles, Calif.....	1	
Atlanta, Ga.....	6		Madison, Wis.....	1	
Battle Creek, Mich.....	1		Marquette, Mich.....	1	
Beatrice, Nebr.....	1		Memphis, Tenn.....	1	
Bellingham, Wash.....	1		Milwaukee, Wis.....	3	
Birmingham, Ala.....	1		Minneapolis, Minn.....	6	
Boise, Idaho.....	1		Morgantown, W. Va.....	3	
Buffalo, N. Y.....	3		Oakland, Calif.....	2	
Chanute, Kans.....	1		Omaha, Nebr.....	3	
Cincinnati, Ohio.....	1		Portland, Oreg.....	27	
Cleveland, Ohio.....	5		Portsmouth, Ohio.....	1	
Dallas, Tex.....	6		Roanoke, Va.....	4	
Danville, Ill.....	1		Rock Island, Ill.....	1	
Davenport, Iowa.....	5		Rome, Ga.....	1	
Denver, Colo.....	3		St. Paul, Minn.....	3	
Detroit, Mich.....	4		Salt Lake City, Utah.....	1	
Duluth, Minn.....	2		San Francisco, Calif.....	5	
El Paso, Tex.....	4		San Jose, Calif.....	1	
Eureka, Calif.....	2		Seattle, Wash.....	21	
Fort Worth, Tex.....	3		Spokane, Wash.....	3	
Grand Rapids, Mich.....	1		Tacoma, Wash.....	1	
Hoquiam, Wash.....	4		Tulsa, Okla.....	1	
Hutchinson, Kans.....	1		Wichita, Kans.....	1	
Independence, Mo.....	3		Winston-Salem, N. C.....	1	
Indianapolis, Ind.....	2		Yakima, Wash.....	4	
Knoxville, Tenn.....	3		Youngstown, Ohio.....	10	
La Fayette, Ind.....	4	1			

TETANUS.**City Reports for Week Ended July 26, 1919.**

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Chicago, Ill.....	2	1	Milwaukee, Wis.....		
Detroit, Mich.....	2	1	Newark, N. J.....	1	1
Kansas City, Kans.....	1		New Orleans, La.....		
Los Angeles, Calif.....	1		Rochester, N. Y.....	1	
Macon, Ga.....			Wilmington, N. C.....		
Memphis, Tenn.....		1			

1911

August 15, 1919.

TUBERCULOSIS.

See Telegraphic weekly reports from States, p. 1901, and weekly reports from cities, p. 1913.

TYPHOID FEVER.**State Reports for June, 1919.**

Place.	New cases reported.	Place.	New cases reported.
Alabama:		Virginia—Continued.	
Barbour County.....	1	Amelia County.....	5
Bullock County.....	2	Amherst County.....	1
Calhoun County.....	2	Woodson.....	1
Chambers County.....	1	Appomattox County.....	5
Chilton County.....	5	Augusta County.....	7
Coffee County.....	3	Staunton.....	2
Colbert County.....	4	Bath County—	
Crenshaw.....	3	Williamsville.....	1
Cullinan County.....	1	Bedford County.....	4
Etowah County.....	1	Botetourt County.....	3
Fayette County.....	1	Brunswick County.....	2
Geneva County.....	3	Buchanan County.....	1
Greene County.....	1	Buckingham County.....	1
Henry County.....	3	Campbell County.....	3
Jackson County.....	1	Caroline County.....	6
Jefferson County.....	29	Carroll County.....	2
Lamar County.....	3	Clarke County—	
Lawrence County.....	2	White Post.....	1
Lee County.....	1	Dickenson County—	
Limestone County.....	1	Clintwood.....	2
Marion County.....	1	Dinwiddie County—	
Marshall County.....	1	Carson.....	3
Mobile County.....	12	Elizabeth City County—	
Monroe County.....	2	Hampton.....	1
Montgomery County.....	8	Hobkiss.....	1
Morgan County.....	0	Essex County.....	3
Pickens County.....	2	Floyd County.....	6
Pike County.....	6	Franklin County.....	1
Randolph County.....	5	Frederick County—	
Sumter County.....	1	Winchester.....	3
Talladega County.....	2	Grayson County—	
Tallapoosa County.....	1	Fife Creek.....	5
Tuscaloosa County.....	3	Greeneville County—	
Total.....	112	North Emporia.....	2
Indiana:		Isle of Wight County—	
Adams County.....	1	Carrsville.....	1
Blackford County.....	1	Dendron.....	1
Cass County.....	1	James City County—	
Clark County.....	5	Williamsburg.....	2
Decatur County.....	1	King and Queen County.....	4
Elkhart County.....	1	Halifax.....	2
Gibson County.....	2	South Boston.....	1
Howard County.....	3	Hanover County.....	4
Huntington County.....	2	Henrico County—	
Kosciusko County.....	1	Richmond.....	6
Lake County.....	9	Henry County.....	4
Marion County.....	4	Lancaster County—	
Perry County.....	1	Irvington.....	1
Scott County.....	1	Lee County.....	5
St. Joseph County.....	5	Loudoun County.....	1
Union County.....	1	Louisa County—	
Vanderburgh County.....	1	Mineral.....	5
Warren County.....	1	Lunenburg County.....	3
White County.....	1	Madison County.....	2
Total.....	42	Mecklenburg County.....	1
Nebraska:		Middlesex County.....	1
Cedar County.....	1	Montgomery County—	
Seward County.....	2	Beloit.....	1
Total.....	3	Radford.....	6
Virginia:		Nancemond County.....	6
Accomac County.....	1	Nelson County.....	1
Albemarle County.....	2	Fairington.....	1
Charlottesville.....	1	Tyrone.....	1
Alexandria County—		Norfolk County.....	10
Alexandria.....	1	Norfolk.....	1
Alleghany County—		Portsmouth.....	6
Clifton Forge.....	1	Northampton County.....	10
Covington.....	1	Cape Charles.....	11
		Franktown.....	2
		Northumberland County.....	1
		Nottoway County—	
		Elsewhere.....	2
		Orange County.....	3
		Gordonsville.....	2

August 15, 1919,

1912

TYPHOID FEVER—Continued.
State Reports for June, 1919—Continued.

Place.	New cases reported.	Place.	New cases reported.
Virginia—Continued.		Virginia—Continued.	
Patrick County.....	7	Southampton County.....	17
Pittsylvania County—		Drewryville.....	2
Danville.....	2	Newsoms.....	1
Powhatan County.....	3	Sebrell.....	3
Princess Anne County.....	9	Sedley.....	1
Prince Edward County—		Tazewell County—	
Prospect.....	1	Pocahontas.....	2
Roanoke County—		Warwick County—	
Roanoke.....	2	Lee Hall.....	1
Salem.....	2	Newport News.....	4
Rockbridge County—		Washington County—	
Buena Vista.....	2	Abingdon.....	1
Rockingham County.....	13	Wise County—	
Harrisonburg.....	1	Coeburn.....	1
Dayton.....	1	Exeter.....	1
Russell County—		Inman.....	1
Belfast Mills.....	1	Roaring Fork.....	1
Dante.....	2	Toms Creek.....	1
Scott County—		Wythe County—	
Nickelsville.....	2	York County.....	1
Shenandoah County.....	3		
Smyth County.....	5	Total.....	297

City Reports for Week Ended July 26, 1919.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Akron, Ohio.....	3		Memphis, Tenn.....	6	3
Alton, Ill.....	1		Milwaukee, Wis.....	1	3
Asbury Park, N. J.....	2		Mobile, Ala.....	1	
Atlanta, Ga.....	2		Montgomery, Ala.....	3	
Baltimore, Md.....	18		Nashville, Tenn.....	8	
Baton Rouge, La.....	1	1	New Britain, Conn.....	1	
Battle Creek, Mich.....	1		New Haven, Conn.....	1	
Bedford, Ind.....	1		New Orleans, La.....	5	
Birmingham, Ala.....	2	1	New York, N. Y.....	10	1
Bloomfield, N. J.....	2		Norfolk, Va.....	1	
Buffalo, N. Y.....	1	1	North Little Rock, Ark.....	1	
Canton, Ohio.....	1		North Tonawanda, N. Y.....	2	1
Charleston, W. Va.....	1		Oklahoma City, Okla.....	4	1
Charlotte, N. C.....	9		Parsons, Kans.....	1	
Chicago, Ill.....	11		Pawtucket, R. I.....		1
Cincinnati, Ohio.....	1		Philadelphia, Pa.....	12	2
Cleveland, Ohio.....	1		Pocatello, Idaho.....	1	
Coffeyville, Kans.....	3	1	Pomona, Calif.....	8	1
Colorado Springs, Colo.....	1		Port Chester, N. Y.....	2	
Columbus, Ga.....	3		Portland, Me.....	4	
Covington, Ky.....	1		Portland, Oreg.....	1	
Cumberland, Md.....	1		Ft. Portsmouth, Va.....	2	2
Dallas, Tex.....	2		Poughkeepsie, N. Y.....	1	1
Dayton, Ohio.....	3		Providence, R. I.....	5	2
Denver, Colo.....	1		Quincy, Mass.....	2	1
Detroit, Mich.....	16	3	Reno, Nev.....	2	
Durham, N. C.....	1		Richmond, Va.....	4	1
East Chicago, Ind.....		2	Roanoke, Va.....	2	
East St. Louis, Ill.....	1		Rochester, N. Y.....	4	
Elgin, Ill.....	1		Rome, Ga.....	1	
Elmira, N. Y.....	2		St. Joseph, Mo.....	3	
Flint, Mich.....	2		St. Louis, Mo.....	7	1
Fort Worth, Tex.....	3	2	St. Paul, Minn.....	1	
Grand Rapids, Mich.....	1		Salt Lake City, Utah.....	1	
Great Falls, Mont.....	1		San Francisco, Calif.....	2	2
Hammond, Ind.....	1		Savannah, Ga.....	3	
Haverhill, Mass.....	1	1	Schenectady, N. Y.....	1	
Houston, Tex.....	3		Spartanburg, S. C.....	1	
Hutchinson, Kans.....	4		Spokane, Wash.....	2	
Independence, Mo.....	3		Springfield, Ohio.....	1	
Indianapolis, Ind.....	2	1	Terre Haute, Ind.....	2	
Kansas City, Kans.....	1		Topeka, Kans.....		1
Knoxville, Tenn.....	2	1	Trenton, N. J.....	1	
Lawrence, Mass.....	1		Tulsa, Okla.....	16	
Leominster, Mass.....	1	1	Tuscaloosa, Ala.....	1	
Little Rock, Ark.....	1		Waco, Tex.....	6	
Los Angeles, Calif.....	7		Washington, D. C.....	4	
Louisville, Ky.....	10	3	Wichita, Kans.....	1	
Dynn, Mass.....	2		Wilmington, N. C.....	1	
Macon, Ga.....	6		Winston-Salem, N. C.....	3	
Martinsburg, W. Va.....	2		Youngstown, Ohio.....	1	

1913

August 15, 1919.

TYPHUS FEVER.

Baltimore, Md.

During the week ended July 26, 1919, one case of typhus fever was reported at Baltimore, Md.

DIPHTHERIA, MEASLES, SCARLET FEVER, AND TUBERCULOSIS.

City Reports for Week Ended July 26, 1919.

City.	Population as of July 1, 1917 (estimated by U. S. Census Bureau).	Total deaths from all causes.	Diphtheria.		Measles.		Scarlet fever.		Tuberculosis.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Aberdeen, Wash.	21,392	—	—	—	—	—	4	—	—	—
Abilene, Tex.	14,954	—	1	—	—	—	—	—	—	—
Adams, Mass.	14,406	3	—	—	—	—	—	—	—	—
Adrian, Mich.	11,570	3	—	—	—	—	—	—	—	—
Akron, Ohio	93,604	17	—	—	3	—	2	—	14	—
Alameda, Calif.	28,433	5	2	—	—	—	—	—	2	2
Albany, N. Y.	106,632	—	—	—	1	—	2	—	2	—
Alton, Ill.	23,783	—	1	—	—	—	—	—	—	—
Anderson, Ind.	24,230	5	—	—	—	—	—	—	—	—
Ansonia, Conn.	16,954	3	—	—	—	—	—	—	1	1
Appleton, Wis.	18,005	—	—	—	—	—	1	—	—	—
Arlington, Mass.	13,073	6	1	—	—	—	—	—	—	—
Asbury Park, N. J.	14,620	8	—	—	—	—	1	—	—	—
Atchison, Kans.	16,785	—	1	—	—	—	—	—	1	—
Atlanta, Ga.	196,144	39	—	—	2	—	1	—	2	2
Atlantic City, N. J.	59,515	12	1	—	4	—	1	—	7	1
Attleboro, Mass.	19,776	2	—	—	—	—	—	—	1	—
Austin, Tex.	35,612	16	—	—	—	—	—	—	1	1
Bakersfield, Calif.	17,543	4	—	—	—	—	—	—	—	—
Baltimore, Md.	594,637	200	15	1	4	—	10	—	27	22
Baton Rouge, La.	17,544	3	—	—	4	—	3	—	—	—
Battle Creek, Mich.	39,159	—	2	—	—	—	1	—	—	—
Bayonne, N. J.	72,204	—	3	—	—	—	1	—	2	—
Beatrice, Nebr.	10,437	3	—	—	—	—	—	—	—	—
Beaumont, Tex.	28,851	9	—	—	—	—	—	—	—	—
Bedford, Ind.	10,613	1	—	—	—	—	1	—	—	—
Beloit, Wis.	18,547	5	—	—	—	—	—	—	1	—
Benton Harbor, Mich.	11,099	3	—	—	—	—	—	—	—	—
Berkeley, Calif.	60,427	2	2	—	—	—	1	—	—	—
Berlin, N. H.	13,892	4	—	—	—	—	—	—	—	1
Beverly, Mass.	22,128	2	—	—	—	—	—	—	—	—
Biddeford, Me.	17,760	7	—	—	—	—	—	—	—	—
Binghamton, N. Y.	54,864	7	—	—	—	—	3	—	3	—
Birmingham, Ala.	189,716	51	2	1	—	—	5	—	8	7
Bloomfield, N. J.	19,013	—	1	—	1	—	—	—	1	—
Boise, Idaho.	35,951	9	—	—	—	—	2	—	—	—
Boston, Mass.	767,813	155	42	—	15	1	9	—	35	15
Brazil, Ind.	10,472	5	—	—	—	—	—	—	—	—
Bridgeport, Conn.	124,724	28	2	—	2	1	1	—	6	6
Bristol, Conn.	16,318	1	—	—	—	—	—	—	—	—
Brockton, Mass.	69,152	5	1	1	6	—	—	—	3	—
Brookline, Mass.	33,526	9	—	—	—	—	—	—	2	1
Brunswick, Ga.	10,984	5	—	—	—	—	—	—	1	—
Buffalo, N. Y.	475,781	122	41	2	10	—	9	—	28	8
Burlington, Iowa.	25,144	2	—	—	—	—	2	—	—	—
Burlington, Vt.	21,802	5	1	—	—	—	—	—	—	1
Butte, Mont.	44,057	—	1	—	—	—	2	—	—	—
Cadillac, Mich.	10,158	4	4	1	—	—	—	—	—	—
Cairo, Ill.	15,995	7	—	—	—	—	—	—	—	—
Cambridge, Mass.	114,293	23	2	—	5	—	3	—	1	—
Camden, N. J.	108,117	—	1	—	—	—	—	—	3	—
Canton, Ohio.	62,566	12	3	—	—	—	—	—	1	2
Chanute, Kans.	12,968	1	—	—	—	—	—	—	—	—
Charleston, S. C.	61,041	20	—	—	—	—	—	—	—	2
Charleston, W. Va.	31,060	10	2	—	—	—	—	—	—	—
Charlotte, N. C.	40,759	15	—	—	2	—	—	—	1	1
Chelsea, Mass.	43,405	9	1	1	—	—	—	—	1	1
Cheyenne, Wyo.	11,320	1	—	—	—	—	—	—	1	1
Chicago, Ill.	2,547,201	501	75	8	114	4	21	1	253	61
Chicopee, Mass.	29,950	5	2	—	—	—	—	—	—	3
Chillicothe, Ohio.	15,325	4	—	—	—	—	—	—	—	—
Cincinnati, Ohio.	414,248	91	2	—	19	—	8	—	28	7

¹Population Apr. 15, 1910.

August 15, 1919.

1914

DIPHTHERIA, MEASLES, SCARLET FEVER, AND TUBERCULOSIS—Contd.

City Reports for Week Ended July 26, 1919—Continued.

City.	Population as of July 1, 1917 (estimated by U. S. Census Bureau).	Total deaths from all causes.	Diphtheria.		Measles.		Scarlet fever.		Tuberculosis.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Cleveland, Ohio.....	692,259	146	21	1	27	—	4	—	19	19
Clinton, Iowa.....	27,678	1	1	—	—	—	—	—	—	1
Clinton, Mass.....	113,075	1	—	—	—	—	—	—	—	1
Coffeyville, Kans.....	18,331	—	—	—	—	—	—	—	—	1
Cohoes, N. Y.....	25,292	6	—	—	—	—	—	—	—	1
Colorado Springs, Colo.....	38,965	15	—	—	1	—	—	—	18	6
Columbia, S. C.....	35,165	—	1	—	—	—	—	—	2	—
Columbus, Ga.....	25,306	11	3	1	3	—	—	—	1	2
Columbus, Ohio.....	220,135	60	3	—	2	—	2	4	5	—
Concord, N. H.....	22,853	8	—	—	—	—	—	1	—	1
Covington, Ky.....	59,623	11	1	—	—	—	—	—	—	—
Cranston, R. I.....	26,773	3	—	—	—	—	2	—	2	1
Cumberland, Md.....	26,685	6	1	—	—	—	—	—	—	—
Dallas, Tex.....	129,733	31	—	—	—	—	—	—	2	1
Danbury, Conn.....	22,931	11	1	—	—	—	—	—	—	—
Danvers, Mass.....	10,037	0	—	—	—	—	—	—	1	—
Danville, Ill.....	32,969	—	—	—	—	—	1	—	—	—
Danville, Va.....	20,183	—	1	—	—	—	2	—	2	—
Davenport, Iowa.....	49,618	—	1	—	—	—	2	—	—	—
Dayton, Ohio.....	128,939	28	—	—	1	1	2	—	6	1
Decatur, Ill.....	41,483	10	—	1	—	—	—	—	—	—
Dedham, Mass.....	10,618	0	1	—	—	—	—	—	—	—
Denver, Colo.....	268,439	68	3	1	5	—	1	—	—	17
Des Moines, Iowa.....	104,052	—	1	—	—	—	—	—	—	—
Detroit, Mich.....	619,648	153	24	4	34	1	11	1	61	22
Dover, N. H.....	13,276	6	1	—	—	—	—	—	—	—
Dubuque, Iowa.....	40,093	—	—	—	—	—	—	—	—	1
Duluth, Minn.....	97,077	17	—	—	6	1	1	—	3	1
Durham, N. C.....	25,160	5	—	—	—	—	—	—	1	—
East Chicago, Ind.....	30,285	9	—	—	—	—	—	—	—	—
East Cleveland, Ohio.....	13,864	—	—	—	—	—	1	—	—	—
Easthampton, Mass.....	10,658	—	—	—	—	—	—	—	1	—
East Orange, N. J.....	43,761	6	1	—	—	—	2	—	3	—
East St. Louis, Ill.....	77,312	16	—	—	—	—	—	—	—	2
Elgin, Ill.....	28,562	3	—	—	1	—	—	—	—	—
Elizabeth, N. J.....	88,830	—	1	—	—	—	—	—	8	2
Elmira, N. Y.....	38,272	11	2	—	1	—	—	3	—	—
El Paso, Tex.....	69,149	27	—	—	2	—	—	—	—	7
Englewood, N. J.....	12,603	2	—	—	—	—	—	1	—	—
Eureka, Cal.....	15,142	2	—	—	1	—	—	—	—	—
Evanston, Ill.....	29,304	—	—	—	1	—	—	—	—	1
Everett, Mass.....	40,160	9	—	—	—	—	—	3	—	3
Everett, Wash.....	37,205	—	—	—	1	—	—	—	—	—
Fargo, N. Dak.....	17,872	3	—	—	—	—	—	1	—	—
Findlay, Ohio.....	14,858	3	—	—	—	—	—	3	—	2
Flint, Mich.....	57,385	9	15	—	1	—	—	—	—	—
Fond du Lac, Wis.....	21,486	2	—	—	—	—	1	—	—	—
Fort Scott, Kans.....	10,564	4	—	—	—	—	—	—	—	1
Fort Wayne, Ind.....	78,014	18	—	—	—	—	—	6	—	1
Fort Worth, Tex.....	109,597	39	2	1	—	—	—	—	—	—
Fostoria, Ohio.....	10,959	4	—	—	—	—	—	—	—	—
Framingham, Mass.....	14,149	5	—	—	—	—	1	—	1	—
Freeport, Ill.....	19,844	5	—	—	—	—	—	—	—	—
Fremont, Nebr.....	10,080	3	—	—	—	—	—	—	—	—
Fremont, Ohio.....	11,034	4	—	—	—	—	—	—	—	—
Fresno, Calif.....	36,314	7	—	—	—	—	—	—	—	1
Galesburg, Ill.....	24,629	5	—	—	—	—	—	—	—	—
Gloucester City, N. J.....	11,375	—	1	—	2	—	—	—	1	—
Grand Rapids, Mich.....	132,831	25	1	—	1	—	—	—	1	2
Great Falls, Mont.....	113,948	8	—	—	—	—	—	—	—	—
Green Bay, Wis.....	30,017	13	—	—	—	—	—	—	—	3
Greenfield, Mass.....	12,251	2	—	—	—	—	—	—	—	—
Greensboro, N. C.....	20,171	6	—	—	—	—	1	—	—	2
Greenwich, Conn.....	19,594	—	—	—	—	—	—	—	1	—
Hackensack, N. J.....	17,412	1	—	—	—	—	—	—	—	—
Hammond, Ind.....	27,016	9	—	—	5	—	—	—	—	—
Harrison, N. J.....	17,345	—	1	—	—	—	—	—	—	—
Hartford, Conn.....	112,851	27	6	—	—	—	1	—	—	—
Haverhill, Mass.....	49,180	10	3	1	—	—	—	—	—	3
Hibbing, Minn.....	17,550	—	2	—	—	—	—	—	—	—
Highland Park, Mich.....	33,859	5	1	—	9	—	1	—	—	—
Hoboken, N. J.....	78,324	8	2	—	—	—	—	4	—	—

¹ Population Apr. 15, 1910.

1915

August 15, 1919.

DIPHTHERIA, MEASLES, SCARLET FEVER, AND TUBERCULOSIS—Contd.

City Reports for Week Ended July 26, 1919—Continued.

City.	Popula-tion as of July 1, 1917 (estimated by U. S. Census Bureau).	Total deaths from all causes.	Diphtheria.		Measles.		Scarlet fever.		Tuber-cu-lo-sis.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Holland, Mich.	12,459	3								
Holyoke, Mass.	66,503	13			1		3		2	
Hoquiam, Wash.	12,230		1				2			
Houston, Tex.	116,878	22	2							
Hudson, N. Y.	12,888	5								
Independence, Mo.	11,984	3								
Indianapolis, Ind.	283,622	72	5	1	4		1		8	6
Ironwood, Mich.	15,095	7					1		2	2
Irvington, N. J.	18,710						1		1	
Ithaca, N. Y.	16,017	3								
Jamestown, N. Y.	37,431	6	1		3				3	1
Janesville, Wis.	14,411	4								
Jersey City, N. J.	312,557		19	1	3		1		14	7
Johnstown, N. Y.	10,678	2								
Joplin, Mo.	33,400	4								
Kalamazoo, Mich.	50,408	13	2		4		2		1	
Kankakee, Ill.	14,270	3	1		1					
Kansas City, Kans.	102,096		2		1				3	
Kearny, N. J.	24,325	3							4	
Kenosha, Wis.	32,833	8	1		1		1		4	4
Knoxville, Tenn.	59,112								3	1
Kokomo, Ind.	21,929	11					1		2	
Lackawanna, N. Y.	16,219	6	1		1					
La Crosse, Wis.	31,835	8								1
La Fayette, Ind.	21,481	6							1	1
Lancaster, Ohio.	16,086	5								
Lawrence, Mass.	102,923	18					2		6	2
Leavenworth, Kans.	19,363								1	
Leominster, Mass.	21,365	3			2					
Lexington, Ky.	41,997	26	1							3
Lima, Ohio.	37,145	9			2				1	
Lincoln, Nebr.	48,957	4			1					
Little Rock, Ark.	58,716	10							1	2
Lockport, N. Y.	20,028	5					1			
Logansport, Ind.	21,338	7			1					
Long Beach, Calif.	29,163	14			2					
Long Branch, N. J.	15,733	2	1		1				1	
Lorain, Ohio.	38,266	8			1				1	1
Los Angeles, Calif.	535,455	127	20		3		3		57	25
Louisville, Ky.	240,808	66	2						9	8
Lowell, Mass.	114,366	18	1	1	1		2		7	
Lynchburg, Va.	33,497	8					1		1	
Lynn, Mass.	104,534	17	3				2		3	2
Macon, Ga.	46,099	25								
Madison, Wis.	31,315	12	1							
Malden, Mass.	52,243	5			1		3		2	1
Manchester, Conn.	15,859	1								
Manchester, N. H.	79,607	22	3							3
Mankato, Minn.	10,365	3								
Marinette, Wis.	14,610	3						1		
Marion, Ind.	19,923	3						1		
Marlboro, Ohio.	24,129								2	
Marquette, Mich.	15,255	3			3				1	1
Martins Ferry, Ohio.	12,555	3			10					
Mason City, Iowa	10,135	2								
Medford, Mass.	12,638	7								
Melrose, Mass.	26,681	8								
Memphis, Tenn.	17,724	5			3					
Methuen, Mass.	151,877	12	1		1		1		4	4
Middletown, N. Y.	14,320	5							1	
Middletown, Ohio.	15,990		1				4			
Millford, Mass.	16,384	4					1		2	1
Millford, Mass.	14,280	2								
Milwaukee, Wis.	445,008	95	8	1	7		14		21	5
Minneapolis, Minn.	373,448	75	10	3	5		6		24	11
Missoula, Mont.	19,075	5							2	
Mobile, Ala.	59,201	10					1		1	
Moline, Ill.	27,976		1						1	
Montgomery, Ala.	44,039	14							2	
Morgantown, W. Va.	14,444	2			2					
Morristown, N. J.	13,410	3								
Moundsville, W. Va.	11,513	2								

¹ Population April 15, 1910.

August 15, 1919.

1916

DIPHTHERIA, MEASLES, SCARLET FEVER, AND TUBERCULOSIS—Contd.

City Reports for Week Ended July 26, 1919—Continued.

City.	Population as of July 1, 1917 (estimated by U. S. Census Bureau).	Total deaths from all causes.	Diphtheria.		Measles.		Scarlet fever.		Tuberculosis.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Mount Vernon, N. Y.	37,991	12	1	—	—	—	—	—	1	1
Muscatine, Iowa	17,713	—	—	—	—	—	—	—	—	—
Nashua, N. H.	27,541	7	—	—	—	—	1	2	—	—
Nashville, Tenn.	118,136	40	—	—	—	—	2	—	8	5
Natick, Mass.	10,140	4	—	—	—	—	—	—	—	—
Newark, N. J.	418,789	97	19	1	3	—	5	—	25	12
New Bedford, Mass.	121,622	30	3	2	9	—	2	—	3	2
New Britain, Conn.	55,385	11	—	—	2	—	2	—	8	1
New Brunswick, N. J.	25,865	—	1	—	—	—	—	—	2	—
Newburyport, Mass.	15,291	5	—	—	—	—	—	—	—	—
New Haven, Conn.	152,275	19	1	—	2	—	1	—	7	1
New London, Conn.	21,199	8	1	—	—	—	—	—	2	—
New Orleans, La.	377,01	112	4	—	—	1	—	—	16	14
Newport, Ky.	32,133	9	—	—	—	—	—	—	1	1
Newport, R. I.	30,585	7	—	—	—	—	1	—	—	2
Newton, Mass.	41,345	6	1	—	—	—	—	—	—	—
New York, N. Y.	5,737,492	1,021	131	7	84	2	31	1	211	116
Niagara Falls, N. Y.	35,466	9	9	1	15	—	2	6	—	1
North Adams, Mass.	122,019	3	—	—	—	—	—	—	—	—
Northampton, Mass.	29,006	7	—	—	—	—	1	—	—	—
North Attleboro, Mass.	11,248	1	1	—	—	—	—	—	—	1
North Little Rock, Ark.	15,515	6	—	—	—	—	—	—	—	2
North Tonawanda, N. Y.	14,060	—	—	—	—	—	—	—	1	—
Norwalk, Conn.	27,332	—	—	—	—	—	—	—	—	—
Norwich, Conn.	21,923	—	—	—	—	—	—	—	1	—
Norwood, Ohio	23,269	2	—	—	—	—	—	—	2	1
Oakland, Calif.	206,495	37	8	1	—	—	3	—	6	1
Oak Park, Ill.	27,816	15	1	—	2	—	—	—	—	—
Ogdensburg, N. Y.	16,845	2	—	—	—	—	—	—	—	—
Oklahoma City, Okla.	97,538	15	1	—	—	—	1	—	—	—
Olean, N. Y.	16,927	3	—	—	—	—	—	—	—	1
Omaha, Nebr.	177,777	22	—	—	—	—	—	—	—	3
Orange, Conn.	14,333	7	—	—	—	—	—	—	—	4
Orange, N. J.	33,635	7	1	—	1	—	—	—	3	1
Oshkosh, Wis.	36,549	8	—	—	—	—	—	—	1	1
Parkersburg, W. Va.	21,059	4	—	—	—	—	—	—	1	2
Parsons, Kans.	15,952	—	—	—	—	—	—	—	2	—
Pasadena, Calif.	49,620	8	—	—	—	—	—	—	2	1
Passaic, N. J.	74,478	8	—	—	—	—	—	—	1	—
Paterson, N. J.	149,512	—	1	—	—	—	3	—	5	—
Pawtucket, R. I.	69,666	18	—	—	—	—	2	—	—	1
Peoria, Ill.	72,184	18	1	—	1	—	—	—	3	1
Perth Amboy, N. J.	42,646	8	—	—	—	—	—	—	2	—
Philadelphia, Pa.	1,735,514	353	52	3	63	2	15	1	86	42
Phillipsburg, N. J.	15,879	4	1	—	—	—	—	—	—	—
Pine Bluff, Ark.	17,777	—	1	—	—	—	—	—	—	—
Piqua, Ohio	14,275	2	—	—	—	—	—	—	1	1
Pittsburgh, Pa.	586,196	—	16	—	6	—	7	—	21	—
Pittsfield, Mass.	39,678	7	—	—	2	—	—	—	2	1
Plainfield, N. J.	24,330	5	—	—	5	—	—	—	2	2
Plattsburg, N. Y.	13,111	2	—	—	—	—	—	—	—	1
Plymouth, Mass.	14,031	4	—	—	—	—	—	—	—	—
Pocatello, Idaho	12,895	—	—	—	—	—	2	—	—	—
Pontiac, Mich.	18,096	7	—	—	8	—	—	—	4	—
Port Chester, N. Y.	16,727	3	—	—	1	—	—	—	1	—
Portland, Me.	64,720	15	—	—	—	—	5	—	—	1
Portland, Ore.	308,399	44	3	1	1	—	4	—	12	1
Portsmouth, Ohio	29,356	—	—	—	—	—	—	—	1	—
Portsmouth, Va.	49,633	27	—	—	1	—	—	—	2	1
Poughkeepsie, N. Y.	39,786	—	—	—	—	—	—	—	3	1
Providence, R. I.	253,895	41	5	—	1	—	5	—	—	2
Quincy, Mass.	39,022	5	—	—	—	—	3	—	—	1
Racine, Wis.	47,455	10	—	—	2	—	—	—	3	2
Rahway, N. J.	10,361	1	—	—	—	—	—	—	—	—
Redlands, Calif.	14,573	3	—	—	—	—	—	—	—	—
Reno, Nev.	15,514	2	—	—	—	—	—	—	—	—
Richmond, Va.	158,702	53	4	1	3	—	—	—	11	8
Riverside, Calif.	20,496	4	4	—	—	—	—	—	—	1
Roanoke, Va.	46,282	16	—	—	—	—	—	—	—	—
Rochester, N. Y.	264,714	63	3	—	2	—	5	—	9	3
Rockford, Ill.	56,739	15	2	—	—	—	1	—	—	3
Rock Island, Ill.	29,452	6	—	—	—	—	—	—	—	—

1 Population April 15, 1910.

1917

August 15, 1919.

DIPHTHERIA, MEASLES, SCARLET FEVER, AND TUBERCULOSIS—Contd.

City Reports for Week Ended July 26, 1919—Continued.

City.	Population as of July 1, 1917 (estimated by U. S. Census Bureau).	Total deaths from all causes.	Diphtheria.		Measles.		Scarlet fever.		Tuberculosis.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Rocky Mount, N. C.	12,673	5							1	1
Rome, Ga.	15,607		2							
Rome, N. Y.	24,259								1	
Rutland, Vt.	15,038	1								
Sacramento, Calif.	68,984	14	2						6	3
Saint Cloud, Minn.	12,013								1	
Saint Joseph, Mo.	86,498	25	3							3
Saint Louis, Mo.	768,630	168	43	4	27		5		26	10
Saint Paul, Minn.	252,465	41	16	1	2		2		12	6
Salem, Mass.	49,346	12	1				1			1
Salt Lake City, Utah	121,623	31	2							2
San Bernardino, Calif.	17,616	4								2
San Diego, Calif.	56,412	24							1	2
Sandusky, Ohio	20,226	10			2					
Sanford, Me.	11,217	1							2	
San Francisco, Calif.	471,023	143	2		6		7		20	14
Santa Cruz, Calif.	15,150	4								
Saratoga Springs, N. Y.	13,839	3							1	1
Saugus, Mass.	10,210	2			3		1			
Sault Ste. Marie, Mich.	14,130		1							
Savannah, Ga.	69,250	36	3	2			1		4	4
Schenectady, N. Y.	103,774	14	2		2		1		3	1
Seattle, Wash.	366,445		5		6		11			
Somerville, Mass.	88,618	13	2				1		5	1
South Bend, Ind.	70,967	15								2
Southbridge, Mass.	14,465	1								
Spartanburg, S. C.	21,985	6	2							
Spokane, Wash.	157,656				7		5			
Springfield, Ill.	62,623	20					2			
Springfield, Mass.	108,668	38	1				1		3	3
Springfield, Mo.	41,169	7								
Springfield, Ohio	52,206	3							2	
Steubenville, Ohio	28,259	4	1							
Stockton, Calif.	36,209	14					1			
Superior, Wis.	47,167	2					1			
Syracuse, N. Y.	158,559	25			1		2		1	2
Tacoma, Wash.	117,446		6				1			
Taunton, Mass.	36,610	9							1	2
Terre Haute, Ind.	67,361	12								1
Toledo, Ohio	202,010	56	2	1	55	1	7		15	4
Topeka, Kans.	49,538	13	1						5	2
Trenton, N. J.	113,974	42			11				7	4
Troy, N. Y.	78,094	25	1						8	3
Tulsa, Okla.	32,507								1	
Tuscaloosa, Ala.	10,824	2	1							2
Vallejo, Calif.	13,803	2								
Waco, Tex.	34,015	7								2
Wakefield, Mass.	12,947	1								
Waltham, Mass.	31,011	6	3		6					
Washington, D. C.	366,282		7	1	1		1		17	8
Watertown, N. Y.	30,404		4				2		2	
Wausau, Wis.	19,666	3							4	1
Webster, Mass.	13,484	3								
Westfield, Mass.	18,769	3	3						1	
West Hoboken, N. J.	44,388	2								1
West New York, N. J.	19,612	2								
West Orange, N. J.	13,984	1	1				1			
Wheeling, W. Va.	43,657	12								
Wichita, Kans.	73,597	16	2						2	
Wilmington, Del.	95,369									
Wilmington, N. C.	33,400	3			1				1	
Winchester, Mass.	19,812	1								
Winona, Minn.	18,583		1							
Winston-Salem, N. C.	33,136	10							1	
Winthrop, Mass.	13,195						1			
Woburn, Mass.	16,076	5								
Worcester, Mass.	166,106	41	3				1		14	5
Yakima, Wash.	22,058				7					
Yonkers, N. Y.	103,066	22	6		6		1		4	1
Youngstown, Ohio	112,282	31	2		4		2		5	3
Zanesville, Ohio	31,320	6							2	

FOREIGN.

AUSTRALIA.

Influenza—Brisbane—Newcastle—South Australia.

Influenza has been further¹ reported in Australia as follows:

Brisbane.—At Brisbane, State of Queensland, during the four weeks ended June 28, 1919, 3,619 new cases of influenza were reported. (Population, 143,000.)

Newcastle.—At Newcastle (with eight suburbs), State of New South Wales, during the period from June 8 to July 5, 1919, 287 new cases of pneumonic influenza were reported. (Population, 49,858.)

State of South Australia.—In the State of South Australia during the period from June 1 to 21, 1919, 1,149 cases of influenza with 94 fatalities were reported. (Population, estimated, 436,640.)

Influenza—State of Western Australia Declared Infected—Quarantine.²

Under date of June 19, 1919, the Governor General of the Commonwealth of Australia declared the State of Western Australia infected with influenza and ordered quarantine measures to be observed against the infected area as regarded persons and merchandise.

BRAZIL.

Influenza—Bahia.

Influenza was reported present at Bahia, Brazil, in January, 1919, with 9 cases, in February with 3 cases, and in March, April, May, and June, 1919, with a total of 9 cases.

Influenza—State of São Paulo.

During the period from January to May, inclusive, 1919, 257 fatal cases of influenza were reported in the State of São Paulo, Brazil. The cases were distributed as follows: São Paulo, capital, 98 deaths (population, 528,295); Campinas, 69 (population, 105,160); Santos, 13 (population, 95,365); Ribeirão Preto, 47 (population, 56,000); São Carlos, 14 (population, 70,209); Guaratinguetá, 8 (population, 51,655); Botucatu, 8 (population, 33,545).

There were reported during the month of November, 1918, 5,304 cases of influenza, and during the month of December, 1918, 597 cases, occurring in the cities of São Paulo, Campinas, Ribeirão Preto, and Santos.³

¹ Public Health Reports, July 11, 1919, p. 1570, and July 25, 1919, p. 1664.

² Idem, July 18, 1919, p. 1612.

³ Idem, Mar. 28, 1919, p. 667.

1919

August 15, 1919.

CHINA.**Cerebrospinal Meningitis—Hongkong.**

Cerebrospinal meningitis has been continuously reported present at Hongkong, China, since the beginning of the current year. During the month of January, exclusive of the week ended January 11, for which data have not been received, 22 cases of cerebrospinal meningitis were reported; during the month of February, 27 cases. In March, 1919, a marked increase was observed, 77 cases being notified. In April, 50 cases were notified; in May, 33 cases. Ten cases were notified from June 1 to 21, 1919.

Quarantine Against Hongkong—Cerebrospinal Meningitis.

During the months of April and May, 1919, quarantine measures were ordered to be enforced against arrivals from Hongkong, on account of cerebrospinal meningitis, as follows: At Amoy, Canton, Chefoo, Newchwang, Ningpo, Shanghai, Swatow, and Tientsin, in China; in the Philippine Islands; at Bangkok, Siam; and at Singapore, Straits Settlements.

Examination of Rats—Hongkong.

During the month of April, 1919, out of 8,215 rats taken, 20 were found plague infected and during the month of May, 1919, out of 10,961 rats taken, 58 were found plague infected.

Mortality—Comparative Statistics, 1916–1918—Hongkong.

During the period 1916–1918, mortality from all causes was reported at Hongkong as follows: 1916, 10,558 deaths; 1917, 10,433; 1918, 13,714 (estimated population, 561,500).

The table following shows mortality from certain diseases during the periods under report:

	1916	1917	1918
Influenza.....			405
Cerebrospinal meningitis.....			973
Smallpox.....	542	549	26
Typhoid fever.....	146	116	179
Diphtheria.....	51	61	98
Cholera.....	10		
Dysentery.....	306	190	225
Plague.....	39	35	251
Malaria.....	378	416	398
Syphilis.....	394	336	416
General tuberculosis.....	561	438	475
Beriberi.....	520	654	804
Pneumonia.....	1,577	1,532	2,251
Phtisis and pulmonary tuberculosis.....	931	854	1,055
Enteritis and gastroenteritis.....	770	775	820

Influenza—Canton.

Influenza was reported prevalent at Canton, China, June 30, 1919.

August 15, 1919.

1920

JAPAN.

Cerebrospinal Meningitis—Kobe—Nov.—Dec., 1918—Jan.—July, 1919.

During the month of November, 1918, cerebrospinal meningitis was present at Kobe, Japan, with 12 reported cases, and from December 1 to 28, with 19 cases. From December 29, 1918, to January 25, 1919, 40 cases were reported; January 26 to February 15, 26 cases; March 2 to 29, 45 cases; March 30 to April 26, 54 cases; April 27 to May 31, 72 cases; June 1 to July 5, 60 cases.

Cholera—Taihoku and Keelung, Island of Taiwan.

Cholera was reported present, August 5, 1919, in the Island of Taiwan (Formosa), at Taihoku, and in Keelung and vicinity.

Influenza—Mortality, 1918-1919.

From October 1, 1918, to April 30, 1919, 250,333 fatal cases of influenza were reported in Japan. (Quoted from the Bulletin of the Quarantine Service, Commonwealth of Australia, June 6, 1919.) Population of Japan, 50,751,900.

NEW ZEALAND.

Further Relative to Influenza.¹

During the period from May 19 to June 2, 1919, 89 cases of influenza were reported in New Zealand. (Population, 1,029,400.)

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER.

Reports Received During Week Ended Aug. 15, 1919.²

CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.
China:				
Canton.....	June 22-28.....	3	2	
Do.....	August 8.....			Present. In foreign section, Isl-
				and of Sha-mien.
India:				
Calcutta.....	June 8-14.....		81	
Indo-China:				
Cochin China—				
Saigon.....	June 9-22.....	112	80	
Japan:				
Taiwan Island—				
Keelung.....	August 5.....			Present, vicinity of Keelung.
Taihoku.....	do.....			Present.
Philippine Islands:				
Manila.....	June 15-21.....	2	1	
Provinces.....				June 15-21, 1919: Cases, 163;
Batangas.....	June 15-21.....	6	5	deaths, 113.
Bohol.....	do.....	8	4	
Bulacan.....	do.....	17	11	
Cavite.....	do.....	9	6	
Ilocos Sur.....	do.....	1		
Laguna.....	do.....	11	7	
Nueva Ecija.....	do.....	14	10	
Pampanga.....	do.....	26	19	
Pangasinan.....	do.....	49	34	
Tayabas.....	do.....	22	17	

¹ Public Health Reports, June 13, p. 1343.

² From medical officers of the Public Health Service, American consuls and other sources.

1921

August 15, 1919.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued.

Reports Received During Week Ended Aug. 15, 1919—Continued.

CHOLERA—Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Philippine Island—Continued.				
Mindanao Provinces	June 22-28	4	3	
Bantangas	June 22-28	48	37	
Bohol	do	3	4	
Bulacan	do	14	7	
Cavite	do	6	6	
Cebu	do	24	11	
Nueva Ecija	do	36	22	
Pampanga	do	31	22	
Pangasinan	do	55	42	
Tayabas	do	64	47	
Union	do	7	7	

PLAQUE.

China:				
Canton	June 22-28	1	1	
Hongkong	June 8-14	27	20	
India:				
Calcutta	do		3	
Indo-China:				
Cochin China—Saigon	June 9-22	4	2	Including district of Saigon.

SMALLPOX.

Brazil:				
Bahia	May 25-June 7	2		
Pernambuco	May 4-25	5		Jan. 12-May 3, 1919: Cases, 10.
Canada:				
New Brunswick—St. John	July 27-Aug. 2	1		
Nova Scotia—Halifax	July 20-26	10		Counties: Pictou, East Hants, and in Inverness County, Cape Breton Island.
Ontario—Hamilton	July 27-Aug. 2	1		
Quebec—Restigouche	June 15-July 31	40		Estimated. Occurring on Indian Reserve.
China:				
Chungking	June 15-28			Present.
Foochow	June 15-July 5			Do.
Nanking	June 22-28			Do.
France:				
Ha're	May 23-30	1		
Paris	June 22-28	5		
Gibraltar	June 28-July 5		2	One from Bay.
Greece:				
Saloniki	June 1-7		10	
India:				
Calcutta	June 8-14		47	
Italy:				
Leghorn	June 24-29	1		
Turin	June 23-29	1		
Japan:				
Ko'e	May 4-July 5	117	57	
Taiwan Island	June 25-July 1	1	1	
Mexico:				
Vera Cruz	June 13-19	2		
Newfoundland				July 19-25, 1919: Outports, 4 cases.
Portugal:				
Lisbon	June 22-28	20		

August 15, 1919.

1922

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued.
Reports Receivd During Week Ended Aug. 15, 1919—Continued.
TYPHUS FEVER.

Place.	Date.	Cases.	Deaths.	Remarks.
Greece: Saloniki.....	June 1-7.....		3	
Italy: Trieste.....	July 6-12.....	1	
Japan: Nagasaki.....	June 30-July 13.....	4	
Mexico: Mexico City.....	July 6-12.....	32	
Portugal: Lisbon.....	June 22-28.....	1	

YELLOW FEVER.

Brazil: Bahia.....	May 18-June 14....	26	11	Jan. 12-May 17, 1919: Cases, 43; deaths, 25.
-----------------------	--------------------	----	----	---

Reports Received from June 28 to Aug. 8, 1919.
CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.
Ceylon: Colombo.....	Apr. 23-26.....	10	
China: Canton.....	June 8-21.....	10	3	
Foochow.....	July 3.....	
Swatow.....	June 2-21.....	118	Present.
India: Bombay.....	Apr. 28-June 7....	29	25	
Calcutta.....	May 4-31.....	476	
Madras.....	May 18-24.....	10	8	
Rangoon.....	Apr. 23-June 7....	73	53	
Indo-China: Cochin-China — Saigon.....	Apr. 21-June 8....	218	152	City and district.
Japan: Pescadores Islands.....	July 14.....	40	In one vilage.
Java: East Java.....	Apr. 23-May 20....	83	66	Apr. 2-May 20, 1919: Cases, 553; deaths, 459.
Surabaya.....	Mar. 28-Apr. 24, 1919: Cases, 1,595; deaths, 1,225.
Mid-Java: Samarang.....	Mar. 28-Apr. 24....	75	74	May 2-June 9, 1919: Cases, 70; deaths, 43.
West Java: Batavia.....	May 2-June 5....	12	5	
Persia: Ardebil.....	May 2.....	Present.
Enzeli.....	Apr. 23.....	1	
Khorram-Ahab.....	May 3.....	Outbreak.
Mianedge.....	Apr. 28.....	Do.
Zindjan.....	Apr. 21-May 4....	49	
Philippine Islands: Manila.....	Apr. 26-May 31....	7	2	May 4-24, 1919: Cases, 567; deaths, 383.
Provinces: Batangas.....	May 4-24.....	25	23	
Bulacan.....	do.....	48	25	
Cebu.....	do.....	162	84	
Laguna.....	do.....	20	15	
Mindoro.....	do.....	19	14	
Misamis.....	do.....	9	2	
Pampanga.....	do.....	166	131	
Tayabas.....	do.....	118	89	
Provinces: Batangas.....	June 1-14.....	25	19	June 1-14, 1919: Cases, 164; deaths, 117.
Bulacan.....	do.....	32	19	
Cavite.....	June 8-14.....	7	2	
Laguna.....	do.....	5	6	
Nueva Ecija.....	June 1-14.....	10	7	
Pampanga.....	do.....	48	38	
Pangasinan.....	June 8-14.....	9	5	
Tayabas.....	do.....	22	17	
Siam: Bangkok.....	Apr. 13-May 17.....	603	

1923

August 15, 1919.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued.
Reports Received from June 28 to Aug. 8, 1919—Continued.
PLAGUE.

Place.	Date.	Cases.	Deaths.	Remarks.
China:				
Canton.....	May 25-June 21.....	Present, Apr. 27-May 10, 1919; Cases, 3; present May 24-June 7, 1919.
Foochow.....	May 18-24.....	Do.
Hongkong.....	June 15-28.....	42	33	
Ecuador:				
Guayaquil.....	June 16-30.....	2	1	Bathing place 65 kilometers from Guayaquil.
Posorja.....	June 1-30.....	3	1	
Egypt.....				
Cities—				
Cairo.....	May 15.....	1	
Kantarah.....	June 19-20.....	4	2	Two European. Septicemic.
Port Said.....	May 1-4.....	1	2	
Suez.....	June 5-11.....	3	3	
Provinces—				
Assiout.....	May 17-June 24.....	80	41	
Beni-Souef.....	May 19-June 21.....	6	5	
Fayoum.....	May 18-June 21.....	8	7	
Girgeh.....	May 15-June 25.....	3	4	
Menoufia.....	June 8-24.....	5	1	
Mimieh.....	May 24-June 23.....	29	11	
Great Britain:				
Liverpool.....	July 30.....	1	1	In dock laborer.
Hawaii:				
Paauhau.....	July 19.....	1	
India:				
Bombay.....	Apr. 28-June 7.....	259	182	Apr. 27-June 7, 1919: Cases, 7,807; deaths, 6,232.
Calcutta.....	May 18-31.....	30	
Karachi.....	May 18-June 21.....	144	131	
Rangoon.....	Apr. 28-June 7.....	46	35	
Indo-China:				
Cochin China—				
Saigon.....	Apr. 21-May 25.....	22	16	City and district.
Japan:				
Yokohama.....	June 9-15.....	1	1	
Java:				
East Java.....	Apr. 23-May 20.....	6	6	Apr. 8-May 20, 1919: Cases, 77; deaths, 77.
Surabaya.....	
Mesopotamia:				
Bagdad.....	Apr. 19-June 6.....	317	246	
Basra.....	May 3-10.....	108	89	Including suburb of Ashar. Total from date of outbreak to May 19, 1919, 288 cases.
Siam:				
Bangkok.....	Apr. 27-May 17.....	2	2	
Straits Settlements:				
Singapore.....	Apr. 13-26.....	2	1	
On vessel:				
S. S. City of Sparta.....	Apr. 19-21.....	1	1	From Bombay Apr. 3, 1919: case a soldier: at sea.
Do.....	May 13-17.....	1	1	At Liverpool: case, a native member of the crew. (Public Health Reports, June 27, 1919, p. 1463.)

SMALLPOX.

Arabia:				
Aden.....	May 13-19.....	1	
Austria:				
Salzburg.....	Mar. 9-Apr. 5.....	50	Mar. 9-Apr. 5, 1919: Cases, 92.
Vienna.....	do.....	17	
Azores:				
St. Michaels.....	June 7-20.....	1	
Brazil:				
Bahia.....	Apr. 20-May 3.....	2	
Canada:				
British Columbia—				
Vancouver.....	June 15-July 5.....	4	
New Brunswick—				
Campbellton.....	June 15-21.....	1	
Moncton.....	July 6-12.....	1	

August 15, 1919.

1924

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued.

Reports Received from June 28 to Aug. 8, 1919—Continued.

SMALLPOX—Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Canada—Continued.				
Novia Scotia—				
Cities—				
Halifax.....	June 15-July 19.....	101	
Sydney.....	June 8-21.....	3	
Counties—				
Antigonish.....	June 28.....			Present.
Cumberland.....	do.....			Do.
Guysborough.....	do.....			Do.
Halifax.....	do.....			Do.
Hants.....	do.....			Do.
Lunenburg.....	do.....			Do.
Ontario—				
Province—				
Hamilton.....	June 29-July 5.....	1	May 1-June 30, 1919: Cases, 166;
Harwich.....	May 1-31.....	14	2	deaths, 4; Township in Kent County.
Ottawa.....	June 15-July 5.....	4	
Peterborough.....	June 15-21.....	4	
Waipole Island—	May 1-31.....	42	Kent County. Island in Lake St. Clair. Among Indians.
Prince Edward Island—				
Charlottetown.....	July 16-19.....	6	
Quebec—				
Montreal.....	June 8-28.....	18	June 8-14, 1919: 10 cases. On incoming vessels.
Quebec.....	June 29-July 19.....	8	Jan. 1-Apr. 30, 1919: Cases, 41; deaths, 1.
Saskatchewan—				
Regina.....				
Ceylon:				
Colombo.....	May 1-31.....	4	
China:				
Amoy.....	May 20-June 16.....		13	
Canton.....	May 18-June 21.....			Present.
Chefoo.....	June 8-21.....			Do.
Chungking.....	May 4-June 14.....			Do.
Foochow.....	May 18-31.....			Do.
Hongkong.....	May 18-June 7.....	4	4	Do.
Nanking.....	May 25-June 21.....			Do.
Chosen (Korea):				
Chemulpo.....	Apr. 1-May 31.....	19	4	
Fusan.....	do.....	294	81	
Seoul.....	do.....	3	1	
Czecho-Slovakia:				
Prague.....	May 18-June 21.....	11	2	
Egypt:				
Alexandria.....	May 14-June 24.....	233	95	
Cairo.....	Jan. 2-Mar. 4.....	81	7	
Finland—				Apr. 16-May 31, 1919: Cases, 357.
Provinces—				
Abo Och Bjorneborg.....	Apr. 16-May 31.....	8	
Kuopio.....	do.....	45	
Nyland.....	do.....	9	
St. Michael.....	do.....	51	
Tavastehus.....	do.....	30	
Vasa.....	do.....	7	
Viborg.....	do.....	196	
France:				
Marseille.....	May 1-31.....		2	
Paris.....	May 11-June 21.....	12	6	
Great Britain:				
Cardiff.....	June 15-July 5.....	4	
Dundee.....	June 1-7.....	1	
Glasgow.....	June 8-21.....	5	
London.....	May 25-July 5.....	12	
Greece:				
Saloniki.....	May 15-21.....		18	
India:				
Bombay.....	Apr. 28-June 7.....	639	221	
Calcutta.....	May 4-31.....		353	
Karachi.....	May 4-June 21.....	28	17	
Madras.....	May 18-24.....	23	11	
Rangoon.....	Apr. 28-June 7.....	162	75	
Indo-China:				
Cochin-China—				
Saigon.....	Apr. 21-May 18.....	11	4	City and district.

1925

August 15, 1919.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued.
Reports Received from June 28 to Aug. 8, 1919—Continued.
SMALLPOX—Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Italy:				
Leghorn.....	June 16-22.....	1		
Messina.....	June 1-21.....	13		
Milan.....	Mar. 1-Apr. 30.....	20	5	
Milazzo.....	June 1-7.....	1	1	
Naples.....	June 2-22.....	96	79	
Palermo.....	May 2-June 23.....	39	5	
Turin.....	May 18-June 22.....	4	1	
Venice.....	May 26-June 1.....	2		
Japan:				
Kobe.....	May 4-31.....	48	17	
Nagoya.....	June 1-7.....	1	1	
Taiwan.....	May 21-June 17.....	9	4	Entire island.
Tokyo.....	May 1-June 5.....	2		
Yokohama.....	May 26-June 1.....	1		
Java:				
East Java.....				
West Java.....				
Batavia.....	Apr. 18-June 5.....	4	1	
Manchuria:				
Dairen.....	May 13-June 2.....	3	2	
Mesopotamia:				
Bagdad.....	May 29-30.....	1		
Mexico:				
Mexico City.....	June 1-July 5.....	20	1	
Piedras Negras.....	June 22-28.....	2	2	
San Jeronimo.....	June 17-30.....	5		
Vera Cruz.....	July 6-12.....	2		
Newfoundland:				
St. Johns.....	June 13-July 4.....	3		
Philippine Islands:				
Manila.....	May 11-17.....	1		
Portugal:				
Oporto.....	June 2-14.....	17	9	
Portuguese East Africa:				
Laurenco Marques.....	Apr. 1-May 31.....	2	1	
Siberia:				
Vladivostok.....	June 8-15.....	23		
Spain:				
Almeria.....	May 18-June 16.....	48	5	
Barcelona.....	May 15-June 19.....	3	6	
Bilbao.....	May 1-10.....	1		
Cadiz.....	Apr. 1-May 31.....		5	
Madrid.....	May 1-31.....	3		
Valencia.....	May 11-June 7.....	174	12	
Vigo.....	Apr. 12.....	2		
Straits Settlements:				
Singapore.....	Mar. 24-May 10.....	4	2	
Tunis:				
Tunis.....	June 15-28.....	2	1	
On vessels:				
S. S. Eastern.....	Apr. 23-26.....	2	1	Death at sea. Second case landed at Woodman's Quarantine Station, Fremantle, Australia, Apr. 29. Vessel from England via Egypt and Colombo.
S. S. Karoa.....	Apr. 19.....	1		Landed at Colombo. Vessel from the United Kingdom via Egypt and Colombo.
S. S. Khyber.....	Apr. 10-May 4.....	4		From Liverpool, via Port Said, Suez, and Colombo. One case landed at Port Said Apr. 10, 2 cases at Colombo Apr. 22, 1 at quarantine, Fremantle, Australia, May 4, 1919.

TYPHUS FEVER.

Algeria:				
Algiers.....	May 1-31.....	76	8	
Austria:				
Vienna.....	Mar. 23-Apr. 5.....	9		Mar. 23-Apr. 5, 1919: Cases, 118.
China:				
Changsha.....	May 11-17.....	1	1	

August 15, 1919.

1926

CHOLEPA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued.
Reports Received from June 28 to Aug. 8, 1919—Continued.
TYPHUS FEVER—Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Chosen (Korea):				
Chemulpo.....	Apr. 1-May 31.....	54	8	
Fusan.....	May 1-31.....	4	1	
Seoul.....	Apr. 1-May 31.....	79	14	
Czecho-Slovakia:				
Prague.....	May 18-24.....	1		
Egypt:				
Alexandria.....	May 14-June 24.....	425	236	
Cairo.....	Jan. 2-Mar. 4.....	150	66	
Port Said.....	Jan. 9-Mar. 4.....	6	4	
Finland:				
Provinces—				
Abo Och Bjorneborg.....	May 15.....	1		
Nyland.....	Apr. 16-May 31.....	4		
St. Michael.....	Apr. 16-May 15.....	8		
Viborg.....	do.....	3		
Germany:				
Jan. 12-Feb. 22.....	344			Military.
Do.....	Feb. 23-Mar. 22.....	220		Civil.
Do.....	Mar. 23-Apr. 12.....	333		Civil, military, prisoners of war, deserters.
Great Britain:				
Glasgow.....	June 8-July 5.....	13	2	
Dundee.....	June 30-July 5.....	3		
Greece:				
Saloniki.....	May 15-21.....		2	
Hungary:				
Budapest.....	Feb. 24-May 9.....	124	6	Feb. 24-May 9, 1919: Cases, 258.
Debraczin.....	do.....	42		
Italy:				
Genoa.....	June 25-July 1.....	62		
Naples.....	May 12-June 22.....	50	16	
Venice.....	Apr. 27-June 14.....	58	9	
Japan:				
Nagasaki.....	June 16-29.....	3		
Mesopotamia:				
Bagdad.....	Apr. 19-June 6.....	34	22	
Mexico:				
Mexico City.....	May 4-July 5.....	211		
Newfoundland:				
St. Johns.....	June 21-27.....	1		From vessel.
Palestine:				
Jaffa.....				Oct. 22-Dec. 22, 1918: Cases, 8; deaths, 3.
Portugal:				
Oporto.....	June 1-15.....	52		
Siberia:				
Vladivostok.....	June 9-15.....	90		
Spain:				
Barcelona.....	May 15-21.....		1	
Madrid.....	May 1-31.....		1	
Tunis:				
Tunis.....	May 24-June 21.....	3	1	

YELLOW FEVER.

Brazil:				
Bahia.....	Apr. 12-May 17....	22	15	July 29, 1919, reported seriously prevalent in States of Bahia and Pernambuco.
Ecuador:				
Guayaquil.....	May 1-31.....	1	1	
Naranjito.....	May 1-June 15....	2	1	
Mexico:				
Merida.....	June 30-July 26....	17	7	State of Yucatan.
Peru:				
Paita.....	July 10-22.....	8	5	Department of Piura.
Piura.....	do.....	46	10	Do.
El Salvador:				
La Union.....	July 6.....	2		
St. Miguel.....	June 24-July 6....	4		
San Salvador.....	do.....	1	1	75 miles from city of San Salvador.

